Practical synthesis of DOPA derivative for biosynthetic production of potent antitumor natural products, saframycins and ecteinascidin 743

Torikai, Kohei

Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, University of Shizuoka

Saruwatari, Takayoshi

Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, University of Shizuoka

Kitano, Tatsuya

Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, University of Shizuoka

Sano, Takuya

Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, University of Shizuoka

他

<http://hdl.handle.net/2324/25642>

出版情報 : Letters in Organic Chemistry. 8 (10), pp.686-689, 2011-12. Bentham Science Publishers

バージョン :

権利関係 : (C) 2011 Bentham Science Publishers

Practical Synthesis of DOPA Derivative for Biosynthetic Production of Potent Antitumor Natural Products, Saframycins and Ecteinasidin 743

Kohei Torikai,^{†,*} Takayoshi Saruwatari,[‡] Tatsuya Kitano,[‡] Takuya Sano, Atsuko Nakane, Hiroshi Noguchi, and Kenji Watanabe
 Department of Pharmacognosy, Graduate School of Pharmaceutical Sciences, University of Shizuoka,
 52-1 Yada, Suruga-ku, Shizuoka 422-8526, Japan

(E-mail: torikai@chem.kyushu-univ.jp)

Abstract A practical synthetic route of DOPA derivative **2**, which should be useful for direct biosynthetic production of potent antitumor natural products, saframycins and ecteinasidin 743 was established. The developed strategy features i) easy-to-handle reactions without special care upon both dryness and inert atmosphere, and ii) the facile HPLC-free purification of **2** via recrystallization enabling scalable synthesis of **2**.

Introduction Saframycins (SMs), produced by *Streptomyces* and various soil bacteria as well as marine vertebrates such as ascidians and sponges, are potent antitumor antibiotics [1]. In particular, a highly potent SM analog, ecteinasidin 743 [2] (ET-743), has recently been in use as an anticancer drug against soft-tissue sarcoma [3]. ETs share the central pentacyclic tetrahydroisoquinoline core with SMs, except for the oxidation state of their terminal rings and the additional ten-membered lactone bridge found in ET-743. Due to the short supply from natural resources, the production of ET-743 should depend on a semi-synthesis including 21 synthetic steps [4]. In order to facilitate the direct biosynthetic production of SMs including ETs, unremitting bioinformatic analyses were carried out, and it was found that SMs are biosynthesized from L-alanine, glycine and two molecules of 3,4-dihydroxyphenylalanine (DOPA) derivative **1** [5] through dual Pictet-Spengler (PS) mechanism [6] (Scheme 1). Briefly, tetrahydroisoquinoline core is constructed by the following three steps; i) Schiff base is formed between DOPA derivative **1** and dipeptidic aldehyde **3**, generated from glycine and L-alanine by the aid of non-ribosomal polypeptide synthetases (NRPSs); ii) PS cyclization occurs to give **4**; and iii) enzymatic region is reductively eliminated to afford aldehyde **5**, which is involved in the same sequence to furnish SMs and ETs through intermediates **6** and **7**. Thus, to develop an engineered perpetual SM-producing system, we cloned necessary biosynthetic gene clusters for SMs and expressed them in model creatures, however, it was unsuccessful to detect the production of SMs even by mass spectrometric analysis [7]. We envisaged that one of the reasons would be insufficient amount of endogenous non-natural amino acid **1**, and if **2** were fed from outside the system, enough supply of SMs would be realized. Herein, we report practical and scalable synthesis of amino acid **2** to tolerate the feeding experiments.

Results and Discussion Synthesis of DOPA derivative **2** commenced with *N*-*t*-butoxycarbonyl (*N*-Boc) tyrosine **8** according to the Schmidt's report [8], one of the most expeditious and easy-to-handle methods to date [9] (Scheme 2). Aldehyde **9** was prepared by Reimer-Tiemann formylation [10a] and the subsequent esterification of *N*-Boc tyrosine **8** as reported previously [8,10b]. Although we attempted the

Scheme 1. Hypothetical Biosynthesis of Saframycin A and Ecteinasidin 743.

conversion of **9** into iodobenzene **10** by the action of I_2 and H_2O_2 , the reaction could not be reproduced even in refluxing ethanol. Hence, we decided to set a robust and reproducible route for the two-steps-introduction of iodine as follows; (i) $NaBH_4$ reduction of aldehyde **9** into the corresponding alcohol [9a], and (ii) iodination of the resulting alcohol with I_2/H_2O_2 combination. As a result, probably due to the fact that electron-withdrawing formyl group was converted to electron-donating hydroxymethyl group, electrophilic iodination proceeded smoothly to afford **11** in good yield (69% for two steps).

Our next task was to oxidize alcohol **11** into Schmidt's intermediate **10** with MnO_2 under argon atmosphere, however, our endeavor was wasteful only to observe the decomposition of substrate **11**. We postulated that *o*-hydroxybenzaldehyde structure of **10** would be unstable even to mild heterogeneous oxidant under inert atmosphere, therefore, methylation of phenolic hydroxy group was first performed. After selective methylation of **11** using methyl iodide and potassium carbonate

Conclusion In conclusion, by improving the Schmidt's method, we have established a practical synthetic route of DOPA derivative **2**, which should be useful for biosynthetic production of saframycins and ecteinascidins, potent antitumor antibiotics. The developed route features higher reproducibility, more facile handling, and much better scalability than the original method to tolerate the feeding experiments. Moreover, this improved route without needs of setting dry, inert, and cold ($< 0\text{ }^{\circ}\text{C}$) conditions should open the door to access amino acid **2** for researchers in biosynthetic community who are not

- [1] For instance, see: a) Arai, T.; Takahashi, K.; Kubo, A. *J. Antibiot.* **1977**, *30*, 1015. b) Scott, J. D.; Williams, R. M. *Chem. Rev.* **2002**, *102*, 1669.
- [2] Rinehart, K. L.; Holt, T. G.; Fregeau, N. L.; Stroh, J. G.; Keifer, P. A.; Sun, F.; Li, L. H.; Martin, D. G. *J. Org. Chem.* **1990**, *55*, 4512.
- [3] Cuevas, C.; Francesch, A. *Nat. Prod. Rep.* **2009**, *26*, 322.
- [4] Cuevas, C.; Pérez, M.; Martín, M. J.; Chicharro, J. L.; Fernández-Rivas, C.; Flores, M.; Francesch, A.; Gallego, P.; Zarzuelo, M.; de la Calle, F.; García, J.; Polanco, C.; Rodríguez, I.; Manzanares, I. *Org. Lett.* **2000**, *2*, 2545.
- [5] a) Mikami, Y.; Takahashi, K.; Yazawa, K.; Namikoshi, M.; Iwasaki, S.; Okuda, S. *J. Biol. Chem.* **1985**, *260*, 344. b) Velasco, A.; Acebo, P.; Gomez, A.; Schellssner, C.; Rodríguez, P.; Aparicio, T.; Conde, S.; Muñoz, R.; de la Calle, F.; Garcia, J. L.; Sánchez-Puelles, J. M. *Mol. Microbiol.* **2005**, *56*, 144. c) -Yu, F. C.; Tang, M.-C.; Peng, C.; Li, L.; He, Y.-L.; Liu, W.; Tang, G.-L. *J. Microbiol. Biotechnol.* **2009**, *19*, 439.
- [6] Koketsu, K.; Watanabe, K.; Suda, H.; Oguri, Oikawa, H. *Nat. Chem. Biol.* **2010**, *6*, 408.
- [7] Details will be published elsewhere.
- [8] Schmidt, E. W.; Nelson, J. T.; Fillmore, J. P. *Tetrahedron Lett.* **2004**, *45*, 3921.
- [9] For recent reports of alternative efficient syntheses of L-3-hydroxy-4-methoxy-5-methyl-phenylalanine derivatives from L-tyrosine, see: a) Chen, R.; Zhu, D.; Hu, Z.; Zheng, Z.; Chen, X. *Tetrahedron: Asymmetry* **2010**, *21*, 39. b) Demoulin, N.; Zhu, J. *Synlett* **2009**, 466.
- [10] For Reimer-Tiemann reaction under slightly hydrated solid-liquid medium conditions, see: a) Thöer, A.; Denis, G.; Delmas, M.; Gaset, A. *Synth. Commun.* **1988**, *18*, 2095. For the application of the conditions in reference [10a] to the synthesis of DOPA derivatives, see: b) Jung, M. E.; Lazarova, T. I. *J. Org. Chem.* **1997**, *62*, 1553.
- [11] Data of 2: Mp. 233–236 °C; $[\alpha]_D^{28}$ -6.1 (c 0.13, H₂O); ¹H NMR (400 MHz, CD₃OD) δ 7.55 (1H, s), 7.34 (1H, s), 3.82 (4H, m), 3.14 (2H, m), 2.32 (3H, s); ¹³C NMR (100 MHz, CD₃OD) δ 173.2, 170.2, 158.7, 136.8, 134.4.

132.4, 130.9, 127.2, 61.9, 57.1, 37.3, 24.8, 16.2 ppm; IR (KBr) 3431, 2989, 2361, 2341, 1695, 1610, 1481, 1396, 1265, 1220, 1139, 1011, 897, 807 cm⁻¹

† Present address: Department of Chemistry, Graduate Faculty of Sciences, Kyushu University, 6-10-1 Hakozaki, Higashi-ku, Fukuoka 812-8581, Japan.

‡ These authors contributed equally to this work.