

Photosensitizing catalysis of the B₁₂ complex without an additional photosensitizer

Shimakoshi, Hisashi

Department of Chemistry and Biochemistry, Graduate School of Engineering, Kyushu University

Li, Li

Department of Chemistry and Biochemistry, Graduate School of Engineering, Kyushu University

Nishi, Masashi

Department of Chemistry and Biochemistry, Graduate School of Engineering, Kyushu University

Hisaeda, Yoshio

Department of Chemistry and Biochemistry, Graduate School of Engineering, Kyushu University

<https://hdl.handle.net/2324/25653>

出版情報 : Chemical Communications. 47 (39), pp.10921-10923, 2011-06-13. RSC Publishing
バージョン :
権利関係 : (C) The Royal Society of Chemistry 2011

Photosensitizing catalysis of B₁₂ complex without additional photosensitizer[†]


Hisashi Shimakoshi, Li Li, Masashi Nishi and Yoshio Hisaeda*

A cobalamin derivative, heptamethyl cobrinate perchlorate, was activated by UV light irradiation to form a Co(I) species in the presence of triethanolamine and used for a dechlorination reaction, and this photochemical reaction was accelerated in an ionic liquid.

The cobalamin derivative (B₁₂) is a cobalt complex with a tetrapyrrole ring system (corrin ring) that has emerged in a variety of enzymes such as methylmalonyl CoA mutase, methionine synthase and reductive dehalogenase.¹ Inspired by the unique functions of these enzymes, various catalytic reactions have already been reported using B₁₂ model complex such as the 1,2-migration of a functional group, dechlorination of organic chlorides, methylation of heavy metals and thiols, etc.² Among the B₁₂ model complexes, the corrinoid compound was mostly developed in the B₁₂ mimic reaction due to its structure and physicochemical properties similar to natural cobalamin.

Most of the catalytic reactions by the corrinoid compound were achieved using the supernucleophilicity of the reduced form of the Co(I) species.³ Therefore, all of the catalytic reaction were coupled with a reducing system such as a chemical reductant (NaBH₄, Zn, Na amalgam and so on), electrochemical reduction and combined use with a photosensitizer.² Recently, we reported the dechlorination of 1,1-bis(4-chlorophenyl)-2,2,2-trichloroethane (DDT) catalyzed by the B₁₂ derivative, heptamethyl cobrinate perchlorate (Co(II) form), with a [Ru(II)(bpy)₃]Cl₂ photosensitizer by irradiation with visible light.⁴ The B₁₂ complex was reduced to form the reactive Co(I) species by electron transfer from the ruthenium photosensitizer in the presence of a sacrificial reductant. Although the B₁₂ complex showed a high catalytic efficiency in the reaction, a photosensitizer was required for the reaction. If the B₁₂ complex itself shows dual properties, photosensitization and catalysis, the catalytic reaction was achieved without any additional photosensitizer. To alleviate this problem, we started to explore the photosensitizing property of the cobalamin derivative and determined the unique photosensitizing property of the B₁₂ complex, especially in an ionic liquid. The ionic liquid is a unique ion pair solvent having various properties such as high polarity, good conductivity and negligible vapor pressure.⁵ In this paper, the unique photosensitizing property of the B₁₂ complex in the ionic liquid and its application for DDT degradation without any photosensitizer are reported (Figure 1).

Fig. 1 Catalytic cycle of B₁₂ complex by UV light irradiation.



The heptamethyl cobrinate perchlorate was first synthesized by Eschenmoser's group⁶ and used by many groups.⁷ It shows a high solubility toward not only a variety of organic solvents, but also ionic liquids such as *N*-methyl-*N*-propylpyrrolidinium bis(trifluoromethanesulfonyl)amide ([P13][TFSA]) and 1-butyl-3-methyl imidazolium bis(trifluoromethanesulfonyl)amide ([bmim][TFSA]) (Chart 1).




Chart 1

When heptamethyl cobrinate perchlorate was dissolved in an ionic liquid ([P13][TFSA]) or methanol containing 0.1 M triethanolamine (TEOA) and subsequently irradiated by UV light ($\lambda_{\text{max}} = 365 \text{ nm}$)[†] under anaerobic conditions, the color of the solution changed from brown to dark green. This photoreaction was well monitored by UV-vis spectroscopy. The UV-vis spectrum of the starting Co(II) form of the B₁₂ complex having absorption maxima at 314 nm and 468 nm was changed to a new

spectrum with absorption maxima at 390 and 560 nm which is typical for the Co(I) state of the B₁₂ complex⁸ by UV light irradiation as shown in Figures 2a (in [P13][TFSA]) and 2b (in MeOH).⁸




Fig. 2 UV-vis spectral change of B₁₂ complex (2.9×10^{-5} M) in the presence of TEOA (0.1 M) by UV light irradiation ($\lambda_{\text{max}} = 365$ nm) under anaerobic condition; (a) in [P13][TFSA], (b) in MeOH.

From these UV-vis spectral changes, the rate constant for the Co(I) formation was determined to be $k = 2 \times 10^{-4} \text{ s}^{-1}$ and $3 \times 10^{-5} \text{ s}^{-1}$ in [P13][TFSA] and MeOH, respectively (ESI[†]). This photochemical reaction was ca. 7 times enhanced in the ionic liquid. In [bmim][TFSA], the B₁₂ complex showed a poor UV-vis change (Figure S1). Because the imidazolium-type ionic liquid has a relatively acidic C-H bond at the imidazole ring,⁹ it should quench the Co(I) species of the B₁₂ complex.¹⁰

The photoreduction of the B₁₂ complex was also investigated by ESR spectroscopy. The heptamethyl cobrinat perchlorate showed the typical Co(II) low-spin signal ($g_1=2.51$, $g_2=2.25$, $g_3=2.00$, $A_2^{\text{Co}}=63$ G, $A_3^{\text{Co}}=134$ G) in [P13][TFSA] containing 0.1 M TEOA as shown in Figure 3a.¹¹ After UV light irradiation, this signal almost disappeared to form the diamagnetic Co(I) species (Figure 3b). This ESR spectral change also indicated that the B₁₂ complex was reduced to the Co(I) form by the UV light irradiation.

Two plausible mechanisms are shown in Figure 4. Once the Co(II) form of the B₁₂ complex is excited, a ligand (corrin ring) to the metal charge transfer state may be formed. While the oxidation state of B₁₂ was Co(III), an ultrafast electronic relaxation of the excited state vitamin B₁₂ (cyanocobalamin) in the gas phase was reported by Shafizadeh et al.¹² The decay is interpreted as resulting from a ring to metal charge transfer. A similar excited state is predicted in the present case, and such a charge-separated (CS) state (Co(I) cation radical) could be

stabilized in an ion-pair solvent, ionic liquid having a high polarity, and subsequent electron transfer from TEOA to form the reduced form of the B₁₂, Co(I) state (Figure 4a). Another possibility is forming an encounter complex between the excited B₁₂ complex and TEOA. In this model, electron transfer from TEOA to the B₁₂ complex in the encounter complex should form the CS state (Co(I)-TEOA^{•+}). This charge-separated state could be stabilized in a polar ionic liquid to prevent a back-electron transfer process (Figure 4b). To further understand the photosensitizing property of the B₁₂ complex, ultrafast transition spectroscopy for the Co(II) state of corrinoid compound is needed.




Fig. 3 ESR spectra of B₁₂ complex (5.0×10^{-4} M) in the presence of 0.1 M TEOA in [P13][TFSA] at 100 K under anaerobic condition. (a) Before UV light irradiation, (b) after 2 hr UV light irradiation.




Fig. 4 Proposed photosensitizing mechanisms of B₁₂ complex by UV light irradiation in the presence of TEOA.

The catalytic reaction was carried out using DDT as a substrate.⁸ The results are shown in Table 1. After a 2 hr UV light irradiation, DDT was converted to a monodechlorinated compound, 1,1-bis(4-chlorophenyl)-2,2-dichloroethane (DDD) in 72 % yield (Entry 1 in Table 1). The reaction did not proceed in

the dark (Entry 2 in Table 1). In MeOH, the yield of DDD was only 9 % which reflects the Co(I) formation efficiency. As for the mechanism, nucleophilic attack of the Co(I) species on DDT could form an alkylated complex as an intermediate as shown in Figure 1.^{4,13} This alkylated complex homolysis by UV light irradiation or thermolysis forms a substrate radical. Hydrogen abstraction from the bulk formed DDD as a dechlorinated product. After the photoreaction with workup, most of the B₁₂ catalyst remained in the ionic liquid and was quantitatively recovered as confirmed by UV-vis spectra (Figure S2). Therefore, the B₁₂ complex could be reused for the successive reaction. In fact, the DDT dechlorination reaction proceeded with almost the same efficiency in the second and third runs (yields of DDD, 2nd 70 % and 3rd 68 %).

Table 1 Photocatalytic dechlorination of DDT catalyzed by B₁₂ complex.^a

Entry	Solvent	Irradiation	Yields (%) ^b	TON ^c
1	[P13][TfSA]	UV	72	35
2	[P13][TfSA]	dark	0	0
3	MeOH	UV	9	4

^a Condition: [B₁₂] = 4.4 × 10⁻⁵ M, [DDT] = 2.2 × 10⁻³ M, [triethanolamine] = 0.1 M. λ_{max} = 365 nm under degassed condition at room temperature. Reaction time, 2 hr.

^b Yields were based on initial concentration of the substrate.

^c Turnover numbers (TON) were based on [B₁₂].

In summary, the photosensitizing property of the B₁₂ complex under anaerobic conditions was investigated. The reductive quenching of the excited state of the B₁₂ complex by a sacrificial reductant provided the reduced form of the B₁₂ complex, Co(I) species. This photosensitizing property of the B₁₂ complex was used for dechlorination of the pollutant, DDT. Therefore, the results reported here will open a door to the new use of the corrinoid compound in photocatalytic chemistry.

This work was partially supported by a Grant-in-Aid for Scientific Research on Priority Areas No. 452, "Science of Ionic Liquids", Innovative Areas No. 2204, "Molecular Activation toward Straightforward Synthesis" and the Global COE Program "Science for Future Molecular Systems" from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan and a Grant-in-Aid for Scientific Research (A), No. 21245016 from the Japan Society for the Promotion of Science (JSPS).

Notes and references

Department of Chemistry and Biochemistry, Graduate School of Engineering, Kyushu University, Motoooka, Fukuoka 819-0395, Japan.

Fax: +81-92-802-2827; Tel: +81-92-802-2826; E-mail:

yhisatcm@mail.cstm.kyushu-u.ac.jp

[†]Electronic supplementary information (ESI) available: Details of experimental procedures. See: DOI: #####

[‡]Black light (UVP, XX-15BLB) was used for the UV light irradiation (1.76 mW cm² at 12-cm distance).

[§]The photochemical reaction did not proceed under visible irradiation using a 200-W tungsten lamp with a 420-nm cut-off filter (Sigma Koki, 42L) and a heat cut-off filter (Sigma Koki, 30H).

^{||}In MeOH, the photochemical reaction slowly proceeded without TEOA.

[¶]Therefore, MeOH also acts as a weak sacrificial reductant.

^{‡‡}General procedure: A 4 mL [P13][TfSA] solution of the B₁₂ complex (4.4 × 10⁻⁵ M), DDT (2.2 × 10⁻³ M) and triethanolamine (0.1 M) was degassed by freeze-pump-thaw cycles. The solution was then stirred at room temperature under irradiation by 365 nm UV light. After 2 hr, the product was extracted with ether-hexane (1:3 v/v). The product, DDD, was identified by NMR and GC-MS comparison with the purchased authentic sample. The B₁₂ complex remaining in the ionic liquid was recycled after being dried under reduced pressure for 24 hr.

- (a) W. Buckel and B. T. Golding, *Chem. Soc. Rev.*, 1996, **25**, 329; (b) *Vitamin B₁₂ and B₁₂-Protein*, (Eds.: B. Kräutler, D. Arigoni and B. T. Golding), Wiley-VCH, Weinheim, 1998; (c) *Chemistry and Biochemistry of B₁₂*, (Eds.: R. Banerjee), Wiley-Interscience, New York, 1999; (d) R. Banerjee and S. W. Ragsdale, *Annu. Rev. Biochem.*, 2003, **72**, 209; (e) T. Toraya, *Chem. Rev.*, 2003, **103**, 2095; (f) B. Kräutler and S. Ostermann, in *The Porphyrin Handbook* (Eds.: K. M. Kadish, K. M. Smith and R. Guilard), Academic Press, 2003, **11**, 229-276.
- (a) K. L. Brown, *Chem. Rev.*, 2005, **105**, 2075; (b) Y. Hisaeda and H. Shimakoshi, in *Handbook of Porphyrin Science* (Eds.: K. M. Kadish, K. M. Smith and R. Guilard), World Scientific, 2010, **10**, 313-370.
- (a) G. N. Schrauzer and E. Deutsch, *J. Am. Chem. Soc.*, 1969, **91**, 3341; (b) A. Fischli and J. J. Daly, *Helv. Chim. Acta.*, 1980, **63**, 1628.
- (a) H. Shimakoshi, M. Tokunaga, T. Baba and Y. Hisaeda, *Chem. Commun.*, 2004, 1806; (b) H. Shimakoshi, S. Kudo and Y. Hisaeda, *Chem. Lett.*, 2005, **34**, 1096.
- (a) T. Welton, *Chem. Rev.*, 1999, **99**, 2071; (b) J. Dupont, R. F. de Souza and P. A. Z. Suarez, *Chem. Rev.*, 2002, **102**, 3667; (c) J. P. Hallett and T. Welton, *Chem. Rev.*, DOI: 10.1021/cr1003248.
- (a) L. Werthemann, R. Keese and A. Eschenmoser, unpublished results. (b) see L. Werthemann, Dissertation, ETH Zürich (No. 4097), Juris Druck und Verlag, Zürich, 1968.
- (a) Y. Murakami, Y. Hisaeda and A. Kajihara, *Bull. Chem. Soc. Jpn.*, 1983, **56**, 3642; (b) C. W.-Exl, T. Darbre and R. Keese, *Org. Biomol. Chem.*, 2007, **5**, 2119; (c) C. Männel-Croisé, B. Probst and F. Zelder, *Anal. Chem.*, 2009, **81**, 9493; (d) H. A. Hassanin, M. F. Ei-Shahat and M. S. A. Hamza, *J. Coord. Chem.*, 2010, **63**, 2431.
- H. Shimakoshi, E. Sakumori, K. Kaneko and Y. Hisaeda, *Chem. Lett.*, 2009, **38**, 468.
- (a) A. G. Avent, P. A. Chaloner, M. P. Day, K. R. Seddon and T. Welton, *J. Chem. Soc., Dalton Trans.*, 1994, 3405; (b) E. Baciocchi, C. Chiappe, T. D. Giacco, C. Fasciani, O. Lanzalunga, A. Lapi and B. Melai, *Org. Chem.*, 2009, **11**, 1413.
- G. N. Schrauzer, *Angew. Chem. Int. Ed. Engl.*, 1976, **15**, 417.
- (a) J. R. Pilbrow, EPR of B₁₂-Dependent Enzyme Reactions and Related Systems, in *B₁₂* (Ed.: D. Dolphin), Wiley, New York, 1982, **1**, 431; (b) S. V. Doorslaser, G. Jeschke, B. Epel, D. Goldfarb, R.-A. Eichel, B. Kräutler and A. Schweiger, *J. Am. Chem. Soc.*, 2003, **125**, 5915; (c) H. Shimakoshi, A. Nakazato, M. Tokunaga, K. Katagiri, K. Ariga, J. Kikuchi and Y. Hisaeda, *Dalton Trans.*, 2003, 2308.
- N. Shafizadeh, L. Poisson and B. Soep, *Chem. Phys.*, 2008, **350**, 2.
- H. Shimakoshi, M. Tokunaga and Y. Hisaeda, *Dalton Trans.*, 2004, 878.

Photosensitizing catalysis of B_{12} complex without additional photosensitizer

