

Variable and boundary selection for functional data via multiclass logistic regression modeling

Matsui, Hidetoshi
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/26121>

出版情報 : Computational Statistics & Data Analysis. 78, pp.176-185, 2014-10. Elsevier
バージョン :
権利関係 :



MI Preprint Series

**Kyushu University
The Global COE Program
Math-for-Industry Education & Research Hub**

Variable and boundary selection for functional data via multiclass logistic regression modeling

Hidetoshi Matsui

MI 2013-7

(Received March 14, 2013)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Variable and boundary selection for functional data via multiclass logistic regression modeling

Hidetoshi Matsui

*Faculty of Mathematics, Kyushu University
744 Motooka, Nishi-ku, Fukuoka 819-0395, Japan.*

hmatsui@math.kyushu-u.ac.jp

Abstract: ℓ_1 penalties such as the lasso provide solutions with some coefficients to be exactly zeros, which lead to variable selection in regression settings. They also can select variables which affect the classification by being applied to the logistic regression model. We focus on the form of ℓ_1 penalties in logistic regression models for functional data, especially in the case for classifying the functions into three or more groups. We provide penalties that appropriately select variables in the functional multinomial regression modeling. Simulation and real data analysis show that we should select the form of the penalty in accordance with the purpose of the analysis.

Key Words and Phrases: Functional Data Analysis, Lasso, Logistic regression model, Model selection, Regularization.

1 Introduction

Variable selection is one of crucial issues in regression analysis. The lasso by Tibshirani (1996) and its offshoots provide us an unified approach for estimation and variable selection problems, and therefore they are broadly applied in several fields (see, e.g. Hastie et al., 2009). We can also select variables which affect to classification problems by applying these types of penalties to logistic regression models (Krishnapuram et al., 2005; Park and Hastie, 2007; Friedman et al., 2010). The logistic regression model is one of the most useful tools for classifying data by providing posterior probabilities which group the data belong to.

In this paper we consider the cases for classifying data into three or more groups using the multiclass logistic regression model, which contains multiple parameters in each variable, as well as a multivariate linear model. Several works have been provided for multivariate linear or multiclass logistic regression models. Turlach et al. (2005) proposed a new penalty to estimate multivariate linear models. They imposed an ℓ_1 sum of maximum absolute values (ℓ_∞ norm) of coefficients with respect to multiple responses, and also generalized it to the ℓ_1 sum of ℓ_q ($q \geq 1$) penalties. Then afterwards Yuan et al. (2007) and Obozinski et al. (2011) let the penalty denote ℓ_1/ℓ_q and investigated theoretical properties of it. Furthermore, Obozinski et al. (2010) proposed a new algorithm for estimating the multinomial logistic regression model with the ℓ_1/ℓ_q regularization for $q = 1, 2$.

When the data to be classified are repeatedly measured over time, it is natural that they are represented by some functional forms. Ramsay and Silverman (2005) established this type of analysis and call it the Functional data analysis (FDA). FDA is one of the most useful methods for effectively analyzing such discretely observed data and has received considerable attentions in various fields (Ramsay and Silverman, 2002; Ferraty and Vieu, 2006; Ferraty, 2011). The basic idea behind FDA is to express repeated measurement data as a smooth function and then draw information from the collection of functional data. FDA includes extensions of traditional analyses such as principal component, discriminant and regression analysis (James et al., 2000; James, 2002).

In this paper we consider the variable selection problem for classifying functional data using the logistic regression model via the sparse regularization. Repeated measurement data are represented by basis expansions, and then the functional logistic regression model is estimated by the penalized maximum likelihood method with the help of sparsity penalties. We propose a new penalty, denoted by $\ell_1\ell_2/\ell_q$ with $q = 1, 2$ here, for appropriately estimating and selecting variables or boundaries for the multiclass functional logistic regression model. Furthermore, the estimated model is evaluated by the model selection criterion since the model evaluation problem is a crucial issue. We examine the proposed method through Monte Carlo simulations and real data examples.

This paper is organized as follows. Section 2 provides a multiclass logistic regression model for functional data. Section 3 shows the method for estimating and evaluating the model. We apply the proposed method to the analysis of simulated data and real data in Section 4 and Section 5 respectively. Concluding remarks are given in Section 6.

2 Multinomial logistic regression model for functional data

Suppose that we have n sets of functional data and a class label $\{(\mathbf{x}_\alpha(t), g_\alpha); \alpha = 1, \dots, n\}$, where $\mathbf{x}_\alpha(t) = (x_{\alpha 1}(t), \dots, x_{\alpha p}(t))^T$ are predictors given as functions and $g_\alpha \in \{1, \dots, L\}$ are classes which \mathbf{x}_α belongs to. In the classification setting, we apply the Bayes rule which assigns \mathbf{x}_α to class $g_\alpha = l$ with the maximum posterior probability given \mathbf{x}_α , denoted by $\Pr(g_\alpha = l | \mathbf{x}_\alpha)$. Then the logistic regression model is given by the log-odds of posterior probabilities:

$$\log \left\{ \frac{\Pr(g_\alpha = l | \mathbf{x}_\alpha)}{\Pr(g_\alpha = L | \mathbf{x}_\alpha)} \right\} = \beta_{l0} + \sum_{j=1}^p \int x_{\alpha j}(t) \beta_{lj}(t) dt, \quad (1)$$

where β_{l0} is an intercept and $\beta_{lj}(t)$ are coefficient functions. We consider that $x_{\alpha j}(t)$ is expressed by basis expansions by

$$x_{\alpha j}(t) = \sum_{m=1}^{M_j} w_{\alpha jm} \phi_{jm}(t) = \mathbf{w}_{\alpha j}^T \boldsymbol{\phi}_j(t), \quad (2)$$

where $\boldsymbol{\phi}_j(t) = (\phi_{j1}(t), \dots, \phi_{jM_j}(t))^T$ are vectors of basis functions such as B -splines or radial basis functions, $\mathbf{w}_{\alpha j} = (w_{\alpha j1}, \dots, w_{\alpha jM_j})$ are coefficient vectors. Since the data are originally observed at discrete time points, we obtain functional data $x_{\alpha j}(t)$ using a smoothing method with basis expansions in advance. In other words, $w_{\alpha j}$ are obtained before constructing the functional logistic regression model (1). Details of the smoothing method are described in Araki et al. (2009b). Furthermore, $\beta_{lj}(t)$ are also expressed by basis expansions

$$\beta_{lj}(t) = \sum_{m=1}^{M_j} b_{ljm} \phi_{jm}(t) = \mathbf{b}_{lj}^T \boldsymbol{\phi}_j(t), \quad (3)$$

where $\mathbf{b}_{lj} = (b_{lj1}, \dots, b_{ljM_j})^T$ are vectors of coefficient parameters.

Using a notation $\pi_l(\mathbf{x}_\alpha; \mathbf{b}) = \Pr(g_\alpha = l | \mathbf{x}_\alpha)$ with $\mathbf{b} = (\mathbf{b}_1^T, \dots, \mathbf{b}_{(L-1)}^T)^T$ and $\mathbf{b}_l = (\beta_{l0}, \mathbf{b}_{l1}^T, \dots, \mathbf{b}_{lp}^T)^T$ since it is controlled by \mathbf{b} , we can re-express the functional logistic regression model (1) as

$$\log \left\{ \frac{\pi_l(\mathbf{x}_\alpha; \mathbf{b})}{\pi_L(\mathbf{x}_\alpha; \mathbf{b})} \right\} = \beta_{l0} + \sum_{j=1}^p \mathbf{w}_{\alpha j}^T \Phi_j \mathbf{b}_{lj} = \mathbf{z}_\alpha^T \mathbf{b}_l, \quad (4)$$

where $\mathbf{z}_\alpha = (1, \mathbf{w}_{\alpha 1}^T \Phi_1, \dots, \mathbf{w}_{\alpha p}^T \Phi_p)$ and $\Phi_j = \int \boldsymbol{\phi}_j^T(t) \boldsymbol{\phi}_j(t)$. It follows from (1) that the posterior probability is given by

$$\begin{aligned} \pi_l(\mathbf{x}_\alpha; \mathbf{b}) &= \frac{\exp(\mathbf{z}_\alpha^T \mathbf{b}_l)}{1 + \sum_{h=1}^{L-1} \exp(\mathbf{z}_\alpha^T \mathbf{b}_h)} \quad (l = 1, \dots, L-1), \\ \pi_L(\mathbf{x}_\alpha; \mathbf{b}) &= \frac{1}{1 + \sum_{h=1}^{L-1} \exp(\mathbf{z}_\alpha^T \mathbf{b}_h)}. \end{aligned}$$

We define vectors of response variables \mathbf{y}_α which indicate class labels as

$$\mathbf{y}_\alpha = (y_{\alpha 1}, \dots, y_{\alpha(L-1)})^T = \begin{cases} (0, \dots, 0, \overset{(l)}{1}, 0, \dots, 0)^T & \text{if } g_\alpha = l, \quad l = 1, \dots, L-1, \\ (0, \dots, 0)^T & \text{if } g_\alpha = L. \end{cases}$$

Then the functional logistic regression model has the following probability function

$$f(\mathbf{y}_\alpha | \mathbf{x}_\alpha; \mathbf{b}) = \prod_{l=1}^{L-1} \pi_l(\mathbf{x}_\alpha; \mathbf{b})^{y_{\alpha l}} \pi_L(\mathbf{x}_\alpha; \mathbf{b})^{1 - \sum_{h=1}^{L-1} y_{\alpha h}}.$$

3 Estimation by the sparse regularization

In order to construct the statistical model, we estimate the functional logistic model (1) by the penalized likelihood method, which maximizes the penalized log-likelihood function given in the form of

$$l_\lambda(\mathbf{b}) = l(\mathbf{b}) - n\lambda P(\mathbf{b}), \quad (5)$$

where $l(\mathbf{b}) = \sum_{\alpha=1}^n \log f(\mathbf{y}_\alpha | \mathbf{x}_\alpha; \mathbf{b})$ is a log-likelihood function, λ is a regularization parameter which controls the effect of the penalty and $P(\cdot)$ is a penalty function. We can also select variables which actually have an influence on the classification by employing one of the ℓ_1 type penalties on $P(\cdot)$. Here we adopt the lasso as the ℓ_1 type penalty in this paper. Now we consider two types of penalties for $P(\cdot)$:

$$P_{(1)}(\mathbf{b}) = \sqrt{M_j} \sum_{j=1}^p \sum_{l=1}^{L-1} \|\mathbf{b}_{lj}\|_2, \quad P_{(2)}(\mathbf{b}) = \sqrt{M_j(L-1)} \sum_{j=1}^p \left\{ \sum_{l=1}^{L-1} \|\mathbf{b}_{lj}\|_2^2 \right\}^{\frac{1}{2}}. \quad (6)$$

In order to select variables appropriately we need to treat $w_{\alpha j1}, \dots, w_{\alpha jM_j}$ as grouped variables and thus apply the group lasso (Yuan and Lin, 2006) to the corresponding coefficients $b_{lj1}, \dots, b_{ljM_j}$, otherwise we fail to select functional variables (Matsui and Konishi, 2011). In addition, $P_{(1)}$ imposes ℓ_1 norms of coefficient parameters for each class. On the other hand, $P_{(2)}$ treats $L-1$ parameters as grouped parameters gain. The former penalty shrinks each coefficient towards exactly zero, whereas the latter shrinks all the $L-1$ parameter vectors in the j -th variable towards zero simultaneously. We denote $P_{(1)}$ and $P_{(2)}$ as $\ell_1\ell_2/\ell_1$ and $\ell_1\ell_2/\ell_2$ penalties respectively. If we consider the case $M_j = 1$ for all j , they correspond to ℓ_1/ℓ_q penalties by Obozinski et al. (2010, 2011) with $q = 1, 2$ respectively, and therefore they are regarded as natural extensions of the ℓ_1/ℓ_q penalties. We can also express the $\ell_1\ell_2/\ell_q$ norm in the following form.

$$P_{(q)}(\mathbf{b}) = C \sum_{j=1}^p \left\{ \sum_{l=1}^{L-1} \|\mathbf{b}_{lj}\|_2^q \right\}^{\frac{1}{q}},$$

here we denoted a constant independent of \mathbf{b} as C .

Since the penalized log-likelihood function involves the ℓ_1 norm of coefficients it is difficult to derive estimates analytically. We apply the local quadratic approximation to the penalty (Tibshirani, 1996; Fan and Li, 2001). Then parameters are updated in the following form:

$$\mathbf{b}^{(k+1)} = \mathbf{b}^{(k)} - \left\{ \frac{\partial^2 l(\mathbf{b})}{\partial \mathbf{b} \partial \mathbf{b}^T} \Big|_{\mathbf{b}^{(k)}} - n \Sigma(\mathbf{b}^{(k)}) \right\}^{-1} \left\{ \frac{\partial l(\mathbf{b})}{\partial \mathbf{b}} \Big|_{\mathbf{b}^{(k)}} - n \Sigma(\mathbf{b}^{(k)}) \mathbf{b}^{(k)} \right\},$$

where $\Sigma(\mathbf{b})$ is given by

$$\Sigma(\mathbf{b}) = \text{diag} \{ \Sigma(\mathbf{b}_1), \dots, \Sigma(\mathbf{b}_{(L-1)}) \},$$

$$\Sigma(\mathbf{b}_l) = \begin{cases} \text{diag} \left\{ \frac{P'_{(1)}(|b_{l1}|)}{|b_{l1}|}, \dots, \frac{P'_{(1)}(|b_{lp}|)}{|b_{lp}|} \right\} & \text{for } P_{(1)}, \\ \text{diag} \left\{ \frac{P'_{(2)}(\|\mathbf{b}_{(1)}\|)}{\|\mathbf{b}_{(1)}\|}, \dots, \frac{P'_{(2)}(\|\mathbf{b}_{(p)}\|)}{\|\mathbf{b}_{(p)}\|} \right\} & \text{for } P_{(2)} \end{cases} \quad (\mathbf{b}_{(j)} = (b_{1j}, \dots, b_{(L-1)j})^T).$$

This update is continued until convergence, and then we obtain a estimated coefficient vector $\hat{\mathbf{b}}$.

Statistical models estimated by the penalized likelihood method depend on regularization parameters and it is a crucial issue to select appropriate values of them. In this case we need to select an appropriate value of λ in the log-likelihood function (5). We use a model selection criterion BIC originally proposed by Schwarz (1978). For the sparse regularization problem, Wang et al. (2007) proved that the BIC select the true model consistently for the SCAD regularization setting. The BIC for evaluating models estimated by the penalized maximum likelihood method with the ℓ_1 penalty is given by

$$\begin{aligned} \text{BIC} &= -2l(\hat{\mathbf{b}}) + \tilde{df} \log n \\ &= \sum_{\alpha=1}^n \sum_{l=1}^{L-1} \left\{ y_{\alpha l} \log \pi_l(\mathbf{z}_{\alpha}; \hat{\mathbf{b}}) + \left(1 - \sum_{h=1}^{L-1} y_{\alpha h} \right) \log \pi_L(\mathbf{z}_{\alpha}; \hat{\mathbf{b}}) \right\} + \tilde{df} \log n, \end{aligned}$$

where \tilde{df} is an effective degrees of freedom. Although Zou et al. (2007) derived a degrees of freedom for the model estimated by the lasso-type regularization, it is not given in the logistic settings. Konishi et al. (2004) proposed a model selection criterion GBIC for evaluating models estimated by the regularization method, though, it needs a second derivative of the penalized likelihood function. On the other hand, they also derived AIC or BIC type criterion with effective degrees of freedoms in the logistic regression model. Using this idea, the degrees of freedom here is given by

$$\tilde{df} = \text{tr}\{W \tilde{Z}_{\mathcal{A}} (\tilde{Z}_{\mathcal{A}}^T W \tilde{Z}_{\mathcal{A}} + n \Sigma(\hat{\mathbf{b}}_{\mathcal{A}}))^{-1} \tilde{Z}_{\mathcal{A}}^T\},$$

where

$$\begin{aligned} \tilde{Z} &= I_{L-1} \otimes Z, \quad Z = (\mathbf{z}_1^T, \dots, \mathbf{z}_n^T), \\ W^{(k)} &= \left(W_{hl}^{(k)} \right)_{hl}, \\ W_{hl}^{(k)} &= \begin{cases} \text{diag}\{\pi_l(\mathbf{z}_1; \hat{\mathbf{b}})(1 - \pi_l(\mathbf{z}_1; \hat{\mathbf{b}})), \\ \dots, \pi_l(\mathbf{z}_n; \hat{\mathbf{b}})(1 - \pi_l(\mathbf{z}_n; \hat{\mathbf{b}}))\} & (h = l), \\ \text{diag}\{-\pi_h(\mathbf{z}_1; \hat{\mathbf{b}})\pi_l(\mathbf{z}_1; \hat{\mathbf{b}}), \\ \dots, -\pi_h(\mathbf{z}_n; \hat{\mathbf{b}})\pi_l(\mathbf{z}_n; \hat{\mathbf{b}})\} & (h \neq l) \end{cases} \end{aligned}$$

and the suffix \mathcal{A} denotes an active set of \mathbf{b} . We select the λ which minimizes the BIC and then adopt the corresponding model as an optimal model.

4 Simulation

In order to investigate the effectiveness and the behavior of the proposed method, we conducted Monte Carlo simulations. We simulated multiple predictors, essentially given as functions having three classes. In this example we aim to examine whether the proposed method appropriately select variables which affect to the classification.

First, we generated $2n$ sets of p predictors each of which were repeatedly measured at several time points; $\{(x_{\alpha i 1}, \dots, x_{\alpha i p}); \alpha = 1, \dots, 2n, i = 1, \dots, n_i\}$, where $x_{\alpha i j}$ ($j = 1, \dots, p$) are assumed to be obtained from $x_{\alpha i j} = u_{\alpha j}(t_i) + \varepsilon_{\alpha i j}$ with uniformly distributed n_i observational points t_i and $\varepsilon_{\alpha i j} \sim N(0, (\sigma R_{\alpha j})^2)$, $R_{\alpha j} = \max_i(u_{\alpha j}(t_i)) - \min_i(u_{\alpha j}(t_i))$. We divided the data with sample size $2n$ in half in order to use them as training and test data. Here we consider a three-class classification problem, and therefore we trisected the data and then assigned each class with them. We set true functions $u_{\alpha j}(t)$ as follows:

$$\begin{aligned} u_{\alpha 1}(t) &= t^3 + a_1 t^2 + a_2 t + a_3, & u_{\alpha 2}(t) &= b_1 \sin(b_2 \pi(t - b_3)), \\ u_{\alpha 3}(t) &= c_1 \exp(c_2 t)(t - c_3)^2, & u_{\alpha 4}(t) &= d_1 \sin(\pi t) + d_2, \end{aligned} \quad t \in [-1, 1],$$

where they are controlled by following random numbers for classes $g = 1, 2, 3$ respectively.

		Class 1	Class 2	Class 3
$u_1 :$	$a_1 \sim$	$N(3.0, 0.1^2)$,	$N(2.0, 0.1^2)$,	$N(4.0, 0.1^2)$,
	$a_2 \sim$	$N(0.5, 0.2^2)$,	$N(0.3, 0.2^2)$,	$N(0, 0.2^2)$,
	$a_3 \sim$	$N(1.0, 0.5^2)$,	$N(1.0, 0.5^2)$,	$N(2.0, 0.5^2)$,
$u_2 :$	$b_1 \sim$	$U(0.4, 0.7)$,	$U(0.8, 1.1)$,	$U(0.4, 0.7)$,
	$b_2 \sim$	$U(0.9, 1.2)$,	$U(0.4, 0.7)$,	$U(0.9, 0.7)$,
	$b_3 \sim$	$N(0.2, 0.1^2)$,	$N(0.5, 0.1^2)$,	$N(0.2, 0.7^2)$,
$u_3 :$	$c_1 \sim$	$U(0.7, 1.1)$,	$U(0.6, 0.9)$,	$U(0.6, 0.9)$,
	$c_2 \sim$	$U(0.7, 1.2)$,	$U(0.8, 1.4)$,	$U(0.8, 1.4)$,
	$c_3 \sim$	$N(0.8, 1.4)$,	$N(0.5, 0.1^2)$,	$N(0.5, 0.1^2)$,
$u_4 :$	$d_1 \sim$	$U(0.5, 3.0)$,	$U(0.5, 3.0)$,	$U(0.5, 3.0)$,
	$d_2 \sim$	$N(0.2, 0.5)$,	$N(0.2, 0.5)$,	$N(0.2, 0.5)$.

Note that there are no classification boundaries for Class 1 and 3 on u_2 , Class 2 and 3 on u_3 and all classes on u_4 , as seen in Table 1 since they respectively have same settings for the random number generation. Especially u_4 does not affect the classification. These facts show that the coefficient functions in the model (1) should be $\beta_{12}(t) = 0$, $\beta_{23}(t) = 0$ and $\beta_{14}(t) = \beta_{24}(t) = 0$.

As the first step of the analysis, we converted longitudinal predictors $x_{\alpha i j}$ into functions. Since they contain additive noises we applied the smoothing method with basis expansions, then obtained functional data sets $x_{\alpha j}(t)$. We used radial basis functions with the idea of B -splines by Kawano and Konishi (2007) for basis functions $\phi_j(t)$ in (2) and (3). In order to reduce computational burden, numbers of basis functions are supposed to be 6 for all variables. Then after constructing the functional logistic regression model, we estimated it by the maximum penalized likelihood method with penalties $P_{(1)}$ and $P_{(2)}$ in (6) and evaluated the model by BIC. We repeated them 100 times for $n = 75, 150, 300$ and $\sigma = 0.05, 0.2$ and then investigated about results of errors and variable selections for two penalties.

Table 1 shows numbers selected by the above method with penalties $P_{(1)}$ and $P_{(2)}$ for 100 repetition, where " $\beta_{1j}\beta_{2j}$ " in the table denotes numbers which both β_{1j} and β_{2j}




Figure 1: True functions for u_j ($j = 1, \dots, 4$ from top left to bottom right) used in the simulation study. Solid, dashed and dotted lines represent functions for Class 1, 2 and 3 respectively.

are estimated to be non-zeros, " β_{lj} " for $l = 1, 2$ denotes numbers which either β_{1j} or β_{2j} is estimated to be non-zero respectively and " ϕ " is numbers which both coefficients are estimated to be zeros. We can find that the penalty $P_{(1)}$ shrinks some of coefficient functions to be zeros appropriately, especially for variable 1 and 3. Thus it selects which classification boundary is important or not for each variable. On the other hand, $P_{(2)}$ selects all or no coefficients for each variable owing to the form of the penalty and hence it cannot select boundaries individually like $P_{(1)}$. However, it excludes variable 4 completely more than $P_{(1)}$, which indicates that $P_{(2)}$ is more suitable for the purpose of only excluding variables themselves which affect the classification. We also obtained training and test errors and selected values of regularization parameters in Table 2. It indicates that the penalty $P_{(1)}$ gives smaller test errors than $P_{(2)}$ in all cases.

Note that we did not mention a classification boundary between Class 1 and 2. We can directly derive the coefficients for the boundary as $\hat{\beta}_{1j} - \hat{\beta}_{2j}$, and if there are no boundaries for Class 1 and 2 there should be $\hat{\beta}_{1j} = \hat{\beta}_{2j}$. However, the above method does not estimate like it and therefore we cannot estimate appropriately, which reminds to be




Figure 2: Yeast cell cycle gene expression profiles for each synchronization. Each plot consists of 5 genes from 5 classes. G1 (solid), G2/M (dashed), M/G1 (dotted), S (dot-dashed), S/G2 (long dashed).

a future work.

5 Real data example

We applied the functional logistic regression modeling to the analysis of yeast cell cycle gene expression data. Spellman et al. (1998) measured expression profiles for 6,178 genome-wide genes in the yeast genome using cDNA microarrays over about two cell cycles. The data contain 77 microarrays consisting of several types of time course synchronization; "cln3" (2 points), "clb2" (2 points), " α -factor" (18 points), "cdc15" (24 points), "cdc28" (17 points) and "elu" (14 points). Spellman et al. (1998) classified 800 genes into 5 groups, G1, G2/M, M/G1, S and S/G2, by the clustering method from the above 77 experiments. Figure 2 shows examples for each synchronization. Araki et al. (2009a) classified genes by using the "cdc15" experiments as functional data and examined the misclassified data from the posterior probabilities. Here we aim to confirm that these experiments actually affect the classification.

Since there are some missing values in expression profiles we excluded some genes from data by following two rules: (1) Genes with at least one missing value for "cln3" and "clb2" respectively are excluded. (2) Those with more than 10 missing values in all for

" α -factor", "cdc15", "cdc28" and "elu" are also excluded. Although the number of genes with no missing values are only 72, we can easily apply the regression model even if there are some (not excessively many) missing values by converting them into functional data. The resulting 657 genes are used for this analysis. First we converted time-course data except for "cln3" and "clb2" into functions. They are expressed by basis expansions with 4 basis functions, which was selected in the functionalization step. Remaining variables "cln3" and "clb2", each of which have only 2 time points, were treated as vector data rather than functional data, and then we treated variables corresponding to 2 time points for each variable as grouped variables. Next we constructed a functional logistic model

$$\log \left\{ \frac{\Pr(g_\alpha = l | \mathbf{x}_\alpha)}{\Pr(g_\alpha = L | \mathbf{x}_\alpha)} \right\} = \beta_{l0} + \sum_{j=1}^2 \sum_{j'=1}^2 x_{\alpha j j'} \beta_{l j j'} + \sum_{j=3}^6 \int x_{\alpha j}(t) \beta_{l j}(t) dt, \quad (7)$$

which is a special case of (1), where X_j ($j = 1, \dots, 6$) respectively correspond to "cln3", "clb2", " α -factor", "cdc15", "cdc28" and "elu". The model was estimated by the penalized likelihood method and evaluated by BIC. Furthermore we altered the class label L in the left hand side of (7) and repeatedly estimated in order to investigate all coefficients for classification boundaries. We repeated this process for 100 bootstrap samples, and then investigated which variable or boundary affects the classification.

Table 3 shows the numbers of selected variables in 100 bootstrap samples for the penalty $P_{(1)}$. We can find that most coefficients are estimated to be non-zeros. However, the coefficient for the boundary between M/G1 and S/G2 is hardly selected for "clb2", and similarly that between M/G1 and S is often excluded. It reveals that the variable "clb2" does not affect the classification. On the other hand, the penalty $P_{(2)}$ selected all variables for 100 repetitions. It indicates the variables themselves are actually relevant to the classification.

6 Concluding remarks

We have proposed a form of penalty for constructing functional multinomial logistic regression models. We derived the estimation and evaluation procedures for the model with the $\ell_1 \ell_2 / \ell_q$ penalty for $q = 1, 2$. The model was fitted by the penalized maximum likelihood method, and the regularization parameter involved in the model was selected by the model selection criterion. Monte Carlo simulations were conducted in order to investigate the effects for the accuracy of prediction and variable selection. Results show that $\ell_1 \ell_2 / \ell_1$ and $\ell_1 \ell_2 / \ell_2$ penalties give different result and therefore we should select these penalties for different uses.

As described in the end of Section 4, there are cases that this method does not estimate coefficients appropriately at one time. It will be a topic for future research. Furthermore we will focus on the theoretical investigation for the functional regression models with the sparse regularization.

References

- Araki, Y., Konishi, S., Kawano, S., and Matsui, H. (2009a), “Functional logistic discrimination via regularized basis expansions,” *Communications in Statistics–Theory and Methods*, 38, 2944–2957.
- (2009b), “Functional regression modeling via regularized Gaussian basis expansions,” *Annals of the Institute of Statistical Mathematics*, 61, 811–833.
- Fan, J. and Li, R. (2001), “Variable selection via nonconcave penalized likelihood and its oracle properties,” *Journal of the American Statistical Association*, 96, 1348–1360.
- Ferraty, F. (2011), *Recent Advances in Functional Data Analysis and Related Topics*, Physica Verlag.
- Ferraty, F. and Vieu, P. (2006), *Nonparametric functional data analysis: theory and practice*, Springer Verlag.
- Friedman, J., Hastie, T., and Tibshirani, R. (2010), “Regularization paths for generalized linear models via coordinate descent,” *Journal of statistical software*, 33, 1.
- Hastie, T., Tibshirani, R., and Friedman, J. (2009), *The elements of statistical learning 2nd ed.*, vol. 1, Springer Series in Statistics.
- James, G. (2002), “Generalized linear models with functional predictors,” *Journal of the Royal Statistical Society: Series B*, 64, 411–432.
- James, G., Hastie, T., and Sugar, C. (2000), “Principal component models for sparse functional data,” *Biometrika*, 87, 587–602.
- Kawano, S. and Konishi, S. (2007), “Nonlinear regression modeling via regularized Gaussian basis functions,” *Bulletin of informatics and cybernetics*, 39, 83–96.
- Konishi, S., Ando, T., and Imoto, S. (2004), “Bayesian information criteria and smoothing parameter selection in radial basis function networks,” *Biometrika*, 91, 27–43.
- Krishnapuram, B., Carin, L., Figueiredo, M., and Hartemink, A. (2005), “Sparse multinomial logistic regression: Fast algorithms and generalization bounds,” *Pattern Analysis and Machine Intelligence, IEEE Transactions on*, 27, 957–968.
- Matsui, H. and Konishi, S. (2011), “Variable selection for functional regression models via the L1 regularization,” *Computational Statistics & Data Analysis*, 55, 3304–3310.
- Obozinski, G., Taskar, B., and Jordan, M. (2010), “Joint covariate selection and joint subspace selection for multiple classification problems,” *Statistics and Computing*, 20, 231–252.

- Obozinski, G., Wainwright, M., and Jordan, M. (2011), “Support union recovery in high-dimensional multivariate regression,” *The Annals of Statistics*, 39, 1–47.
- Park, M. and Hastie, T. (2007), “L1-regularization path algorithm for generalized linear models,” *Journal of the Royal Statistical Society: Series B*, 69, 659–677.
- Ramsay, J. and Silverman, B. (2002), *Applied functional data analysis: methods and case studies*, vol. 77, Springer, New York.
- (2005), *Functional data analysis*, Wiley Online Library.
- Schwarz, G. (1978), “Estimating the dimension of a model,” *The annals of statistics*, 6, 461–464.
- Spellman, P., Sherlock, G., Zhang, M., Iyer, V., Anders, K., Eisen, M., Brown, P., Botstein, D., and Futcher, B. (1998), “Comprehensive identification of cell cycle-regulated genes of the yeast *Saccharomyces cerevisiae* by microarray hybridization,” *Molecular biology of the cell*, 9, 3273–3297.
- Tibshirani, R. (1996), “Regression shrinkage and selection via the lasso,” *Journal of the Royal Statistical Society. Series B (Methodological)*, 58, 267–288.
- Turlach, B., Venables, W., and Wright, S. (2005), “Simultaneous variable selection,” *Technometrics*, 47, 349–363.
- Wang, H., Li, R., and Tsai, C. (2007), “Tuning parameter selectors for the smoothly clipped absolute deviation method,” *Biometrika*, 94, 553–568.
- Yuan, M., Ekici, A., Lu, Z., and Monteiro, R. (2007), “Dimension reduction and coefficient estimation in multivariate linear regression,” *Journal of the Royal Statistical Society: Series B*, 69, 329–346.
- Yuan, M. and Lin, Y. (2006), “Model selection and estimation in regression with grouped variables,” *Journal of the Royal Statistical Society: Series B*, 68, 49–67.
- Zou, H., Hastie, T., and Tibshirani, R. (2007), “On the “degrees of freedom” of the lasso,” *The Annals of Statistics*, 35, 2173–2192.

Table 1: Numbers of selected variables. Top: $P_{(1)}$, bottom: $P_{(2)}$.

$P_{(1)}$	σ	0.05				0.2			
n	j	$\beta_{1j}\beta_{2j}$	β_{1j}	β_{2j}	ϕ	$\beta_{1j}\beta_{2j}$	β_{1j}	β_{2j}	ϕ
75	1	100	0	0	0	100	0	0	0
	2	37	0	63	0	33	0	67	0
	3	0	100	0	0	1	99	0	0
	4	0	25	6	69	12	45	16	27
150	1	100	0	0	0	100	0	0	0
	2	32	0	68	0	49	0	51	0
	3	0	100	0	0	1	99	0	0
	4	1	33	2	64	15	42	14	29
300	1	100	0	0	0	100	0	0	0
	2	45	0	55	0	34	0	66	0
	3	0	100	0	0	1	99	0	0
	4	2	32	5	61	26	40	15	19

$P_{(2)}$	σ	0.05				0.2			
n	j	$\beta_{1j}\beta_{j2}$	β_{1j}	β_{2j}	ϕ	$\beta_{1j}\beta_{2j}$	β_{1j}	β_{2j}	ϕ
75	1	100	0	0	0	100	0	0	0
	2	98	0	0	2	91	0	0	9
	3	100	0	0	0	99	0	0	1
	4	18	0	0	82	54	0	0	46
150	1	100	0	0	0	100	0	0	0
	2	100	0	0	0	100	0	0	0
	3	100	0	0	0	100	0	0	0
	4	22	0	0	78	51	0	0	49
300	1	100	0	0	0	100	0	0	0
	2	100	0	0	0	100	0	0	0
	3	100	0	0	0	100	0	0	0
	4	26	0	0	74	71	0	0	29

Table 2: Averages of 100 training, test errors (%) and the regularization parameter λ (and its standard deviations in parenthesis). The top is for $P_{(1)}$ and the bottom is for $P_{(2)}$.

σ	0.05			0.2		
n	Train	Test	$\lambda \times 10^3$	Train	Test	$\lambda \times 10^3$
75	0.04	0.41	5.81 (2.74)	0.15	1.45	9.64 (4.74)
150	0.10	0.33	3.79 (2.02)	0.35	1.12	6.51 (3.20)
300	0.10	0.28	2.22 (1.13)	0.28	0.92	3.68 (1.64)

σ	0.05			0.2		
n	Train	Test	$\lambda \times 10^3$	Train	Test	$\lambda \times 10^3$
75	0.08	0.80	6.28 (2.94)	0.27	1.93	12.69 (6.28)
150	0.15	0.45	4.04 (2.20)	0.49	1.56	7.96 (3.77)
300	0.12	0.38	2.37 (1.27)	0.40	1.17	4.29 (1.58)

Table 3: Numbers of selected variables for $P_{(1)}$. Signs i/j below indicate classification boundaries between i and j for $i, j = 1, \dots, 5$ with 1:G1, 2:G2/M, 3:M/G1, 4:S, 5:S/G2.

	cln3	clb2	α	cdc15	cdc28	elu
1/2	100	100	100	100	100	98
1/3	100	100	100	99	100	100
1/4	94	99	100	100	100	100
1/5	100	100	100	100	100	100
2/3	100	76	100	100	100	100
2/4	100	100	99	100	100	100
2/5	100	100	100	100	100	100
3/4	100	47	100	100	100	100
3/5	100	16	100	100	100	100
4/5	100	82	100	100	98	98

List of MI Preprint Series, Kyushu University

The Global COE Program

Math-for-Industry Education & Research Hub

MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO
On the L^2 a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI
Variable selection for functional regression model via the L_1 regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA
Hypergeometric q -functions of the q -Painlevé system of type $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI
Large time behavior of the semigroup on L^p spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO
On very accurate enclosure of the optimal constant in the a priori error estimates for H_0^2 -projection
- MI2009-26 Manabu YOSHIDA
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA
Projective reduction of the discrete Painlevé system of type $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra

- MI2010-25 Toshimitsu TAKAESU
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI
Spin-spin correlation functions of the q -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO
Hypergeometric τ functions of the q -Painlevé systems of type $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUOKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA
CAP representations of inner forms of $Sp(4)$ with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA
Solutions to a q -analog of Painlevé III equation of type $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE
RIPless Theory for Compressed Sensing

MI2013-6 Hiroshi INOUE

Improved bounds on Restricted isometry for compressed sensing

MI2013-7 Hidetoshi MATSUI

Variable and boundary selection for functional data via multiclass logistic regression modeling