

Induction of Polyploid *Phalaenopsis amabilis* by N₂ Treatment

Wongprichachan, Prapassorn
Department of Horticultural Science, National Chiayi University

Huang, Kuang-Liang
Department of Horticultural Science, National Chiayi University

Hsu, Shan-Te
Department of Horticultural Science, National Chiayi University

Chou, Yi-Mei
Department of Horticultural Science, National Chiayi University

他

<https://doi.org/10.5109/26158>

出版情報：九州大学大学院農学研究院紀要. 58 (1), pp.33-36, 2013-02. Faculty of Agriculture,
Kyushu University

バージョン：

権利関係：

Induction of Polyploid *Phalaenopsis amabilis* by N₂O Treatment

Prapassorn WONGPRICHACHAN¹, Kuang-Liang HUANG^{1*}, Shan-Te HSU¹,
Yi-Mei CHOU¹, Tzu-Yu LIU¹ and Hiroshi OKUBO

Laboratory of Horticultural Science, Division of Agronomy and Environmental Sciences,
Department of Agro-environmental Sciences, Faculty of Agriculture,
Kyushu University, Fukuoka 812-8581, Japan

(Received October 31, 2012 and accepted November 8, 2012)

Ploidy levels of the progenies obtained by the crossing of untreated *Phalaenopsis amabilis* (2n=2x=38) with the same species treated with N₂O for 24 or 48 hrs were determined by flow cytometry and chromosome counting. The treatment for 24 hrs resulted in 35.6% triploid and 6.7% tetraploid induction. Only 5% tetraploid progenies were obtained by the treatment for 48 hrs.

Key words: Flow cytometry, nitrous oxide, polyploidy, *Phalaenopsis*

INTRODUCTION

Polyploids have played an important role in higher plant evolution, as most flowering plants are polyploids (Leitch and Bennett, 1997; Rubuluza *et al.*, 2007). Polyploidy has been used in horticulture as a breeding tool to enhance characteristics such as flower size, flower number, plant size, increased width-to-length ratio of leaves and leaf thickness (Shao *et al.*, 2003; Rubuluza *et al.*, 2007).

Phalaenopsis is one of the most popular orchids in the world market (Wang, 2004). The basic chromosome number of the genus is 2n=2x=38. The most popular commercial varieties/ hybrids are usually tetraploid, while the wild species are diploid. Therefore, it is hard to transfer decent genes from wild species to the most advanced varieties or hybrids. To develop a conversion procedure to double the ploidy level of the wild species of *Phalaenopsis* enables hybridization with the commercial varieties (Chen *et al.*, 2009). Chromosome doubling to produce different ploidy levels in plants is usually achieved by anti-microtubule agents such as colchicines, oryzalin, trifluralin and nitrous oxide gas (N₂O). We showed in the previous report the effectiveness of N₂O application to induce unreduced gamete of *Phalaenopsis amabilis* (Wongprichachan *et al.*, 2013)

Flow cytometry is a rapid and exact method for estimating nuclear DNA content (Galbraith *et al.*, 1983). It can be efficiently used for ploidy determination in plants growing in the field, greenhouse and *in vitro* (Doležel *et al.*, 1989; Śliwińska and Steen, 1995; Joachimiak *et al.*, 2001; Thiem and Śliwińska, 2003). This technique has been proven to be faster and precise than conventional methods such as chromosome counting or measuring stomata length (Sgorbati *et al.*, 1986; Ollittraut-Sammarcelli *et al.*, 1994; Tosca *et al.*, 1995; O'Brien *et al.*, 1996; Ozaki *et al.*, 1998; Pinheiro *et al.*, 2000; Vainola, 2000; Thao *et al.*, 2003).

The aim of this study was to produce polyploids of *P. amabilis* by using unreduced gametes induced by N₂O treatments.

MATERIALS AND METHODS

Plant materials

Phalaenopsis amabilis (2n=2x=38) were used in this study. Plants were cultivated in 10 cm diameter pots in a greenhouse with a 16/8 h (day/night) photoperiod at the Horticulture Technology Center, National Chiayi University, Chiayi, Taiwan (R.O.C.).

Pollen grains of *P. amabilis* treated with N₂O for 0, 24 or 48 hrs were hand pollinated to untreated plants of *P. amabilis*. Ten flowers were used for each crossing. Capsules were harvested 130 days after pollination.

Three capsules per crossing were surface-sterilized for 20 min in 20% (v/v) Clorox solution and 0.05% Tween20 and subsequently rinsed 3 times with sterilized distilled water. Seeds were germinated aseptically in 90×15 mm petri dishes containing 25 mL basal medium supplemented with 2.5 g/L Hyponex #1 (7N-6P₂O₅-19K₂O), 200 mg/L myo-inositol, 2 g/L peptone, 1 g/L ammonium nitrate, 60 g/L Potato, 1 mg/L naphthalene acetic acid (NAA), 0.5 mg/L kinetin, 0.5 mg/L indole-3-butyric acid, 1 g/L activated charcoal, 20 g/L sucrose and 10 g/L agar. The pH value of the media was adjusted to 5.4 with 1 N KOH or HCl prior to autoclaving for 15 min at 121°C. After 3 months, the developed protocorms were transferred to new medium supplemented with 2.5 g/L Hyponex #1, 200 mg/L myo-inositol, 2 g/L peptone, 25 g/L Potato, 30 g/L Banana, 0.5 mg/L NAA, 1 g/L activated charcoal, 20 g/L sucrose and 15 g/L agar. Plantlets were subcultured on the medium of the same constituents every 2 months. All cultures were incubated at 25±2°C in a 16/8 hrs photoperiod at light intensity of 54 μmol m⁻² s⁻¹ with cool-white fluorescent light lamps.

Sample preparation and Giemsa staining for chromosome observation

Growing root tips were pretreated with 2 mM 8-hydroxyquinoline (Sigma-Aldrich, St. Louis) on a

¹ Department of Horticultural Science, National Chiayi University, Chiayi 60004, Taiwan, Republic of China

* Corresponding author (E-mail: kluang@mail.ncyu.edu.tw)

rotary shaker (100 rpm) for 5 hrs at 20°C, and fixed in 95% ethanol–acetic acid (3:1 by vol.) for 24 hrs at 4°C. They were digested with 1% (w/v) cellulase (Sigma–Aldrich) and 0.5% (w/v) pectinase (Fluka) in 10 mM of citrate buffer (Pedrosa *et al.*, 2001). The digested samples were mounted on a slide with a droplet of 10% Giemsa (Sigma–Aldrich), covered with a cover slide, and microscopically analyzed (Axio version Rel. 4.5, Germany) with a CCD camera (Cannon A620, Japan).

Preparation of nuclear suspension for determination of ploidy levels by flow cytometry

Nuclear DNA content in young leaves (0.3–0.5 cm²) of the plantlets were analyzed by flow cytometry (Beckman, Germany). Samples were prepared according to the manual of the Partec Cystain Cystain PI Absolut P kit, which included the extraction and staining buffers. A sharp razor blade was used to chop fresh tissue into pieces with a size of <1 mm in a 6 cm glass petri dish containing 500 μ L of the extracting buffer. After 30–90 sec of incubation in the Extraction Buffer the samples were filtered through a Partec 50 μ m CellTrics disposable filter. Then 2 ml of staining solution Propidium Iodide (with PI and Rnase) was added. The process of nuclear extraction and staining was carried out on ice.

Determination of ploidy levels of plantlet by flow cytometry

To determine the ploidy levels of the samples of each treatment, young leaves of *P. aphrodite* var. *formosana* with diploid level ($2n=2x=38$) were used as reference for the 2C DNA value which was set to the channel at 50 (G_0/G_1 peak) of the machine using linear scale. The ploidy levels of the samples were determined by comparison with these reference 2C DNA values. During analyses, samples of *P. aphrodite* var. *formosana* (diploid) were regularly run, in order to check the stability of peak position corresponding to the diploid level. Results were displayed in the histograms showing number of nuclei according to relative fluorescence intensity, which is proportional to DNA content. Diploids progenies showed 2C peak at 50 (G_0/G_1 peak) of the machine using linear scale and tetraploid progenies will show first peak position on 100 of the machine. In each sample the DNA content of at least 5,000 nuclei was analyzed and percentage of ploidy levels of progenies was calculated based on at least 20 observations of each treatment.

RESULTS

Some progenies from the crossing of untreated parents with N_2O –treated parents and *P. aphrodite* var. *formosana* as internal standards exhibited G_0/G_1 peak at the same channel, indicating that these progenies are diploid (Fig. 1). The flow cytometry histograms of the diploid progenies (Figs. 2B and 3B) were clearly different from those of other progenies, and they were considered to be triploid (Fig. 2C) and tetraploid (Figs. 2D and 3C). The ratio of genome values in the first peak of diploid: triploid: tetraploid progenies was about 1 : 1.5 : 2.0, which indicates that the genome sizes of triploid and tetraploid progenies were about 1.5– and 2–fold bigger than those of diploid progenies, respectively (Figs. 2 and 3).

N_2O treatment for 24 hrs induced triploid (36%) and tetraploid (7%), whereas that for 48 hrs induced only 5% tetraploid (Table 1).

The chromosome observation confirmed the ploidy levels of the progenies. Chromosome numbers of the diploid, triploid and tetraploid progenies were $2n=2x=38$, $2n=3x=57$ and $2n=4x=76$, respectively (Fig. 4).




Fig. 1. Flow cytometry histograms of a diploid *P. aphrodite* var. *formosana* (A) and progenies of untreated *P. amabilis* crossed with 24 h N_2O –treated *P. amabilis* (B).

Table 1. Effects of N_2O on ploidy levels in the progenies of untreated \times treated *P. amabilis*

N_2O treatment (hr)	No. of plantlets examined	% ploidy (No.)		
		Diploid	Triploid	Tetraploid
0	20	100 (20)		
24	45	57.8 (26)	35.6 (16)	6.7 (3)
48	20	95.0 (19)		5.0 (1)




Fig. 2. Flow cytometry histograms of the progenies of untreated *P. amabilis* crossed with 24 hrs N_2O -treated *P. amabilis*. (A) Untreated parent, (B) diploid progenies, (C) triploid progenies and (D) tetraploid progenies.




Fig. 3. Flow cytometry histograms of the progenies of untreated *P. amabilis* crossed with 48 hrs N_2O -treated *P. amabilis*. (A) Untreated parent, (B) diploid progenies and (C) tetraploid progenies.




Fig. 4. Chromosomes of root tip cells of (A) diploid ($2n=2x=38$) (B) triploid ($2n=3x=57$) and (C) tetraploid ($2n=4x=76$) progenies. bar= $10\mu m$.

DISCUSSION

Polyploidy induction has been successfully applied in ornamental, vegetable and medicinal plants by polyploidizing agents such as colchicines, oryzalin, trifluralin and N_2O (Petersen *et al.*, 2003; Thao *et al.*, 2003, 2004; Rubuluza *et al.*, 2007; Praça *et al.*, 2009). N_2O has been used previously in tulips (Okazaki *et al.*, 2005), lilies (Barba-Gonzalez *et al.*, 2006; Akutsu *et al.*, 2007) and *Begonia* (Dewitte *et al.*, 2010) for polyploidy breeding by inducing unreduced gametes and to restore fertility in sterile genotypes.

Our results showed that N_2O treatments are effective methods to induce polyploidy in *P. amabilis*. The histograms in Figs. 2C, 2D and 3C are representative fluorescence profiles for nuclei from the diploid, triploid and tetraploid plants, which are similar to the results of Petersen *et al.* (2003), Thao *et al.* (2003, 2004) and Rubuluza *et al.* (2007). Our results indicate that the procedure was adequate for induction polyploidy *P. amabilis* plantlets. Induction of tetraploid progenies in this study suggests that the plant also produced $2n$ eggs in the crossings without N_2O treatment. The agreement of the results of chromosome count in root tips and flow cytometric values demonstrates the reliability of flow

cytometry as a practical and rapid tool for screening the ploidy level of the progenies as shown by Galbraith *et al.* (1983) and Thao *et al.* (2004), since a plant can be analyzed when the first leaf appears *in vitro* before acclimatization to greenhouses.

In conclusion, successful induction of polyploids in *P. amabilis* confirmed the effectiveness by N₂O as a polyploidizing agent. The polyploids that we obtained, will be used in further breeding works.

ACKNOWLEDGEMENTS

The authors would like to thank Tainan District Agricultural Research and Extension Station, COA, Executive Yuan, Taiwan (R.O.C.) for kindly providing support in this research.

REFERENCES

- Alexander, M. P. 1980 A versatile staining for pollen, fungi yeast and bacteria. *Stain Technol.*, **55**(1): 13–18
- Akutsu, M., S. Kitamura, R. Toda, I. Miyajima and K. Okazaki 2007 Production of 2n pollen of Asiatic hybrid lilies by nitrous oxide treatment. *Euphytica*, **155**: 143–152
- Barba-Gonzalez, R. M., C. T. Miller, M. S. Ramanna and J. M. van Tuyl 2006 Nitrous oxide N₂O induces 2n gametes in sterile F₁ hybrids of Oriental × Asiatic lilies (*Lilium*) and leads to inter-genomic recombination. *Euphytica*, **148**: 303–309
- Chen, W. H., C. Y. Tang and Y. L. Kao 2009 Ploidy doubling by *in vitro* culture of excised protocorms or protocorm-like bodies in *Phalaenopsis* species. *Plant Cell, Tiss. Org. Cult.*, **98**: 229–238
- Dewitte, A., T. Eeckhaut, J. van Huylenbroeck and E. van Bockstaele 2010 Induction of 2n pollen formation in *Begonia* by trifluralin and N₂O treatment. *Euphytica*, **171**: 283–293
- Doležel, J., P. Binarova and S. Lucretti 1989 Analysis of nuclear DNA content in plant cells by flow cytometry. *Biol. Plant.*, **31**: 113–120
- Galbraith, D. W., K. R. Harkins, J. R. Maddox, N. M. Ayres, D. P. Sharma and E. Firoozabady 1983 Rapid flow cytometric analysis of the cell cycle in intact plant tissues. *Science*, **220**: 1049–1051
- Joachimiak, A., A. Kula, E. Śliwińska and A. Sobieszczkańska 2001 C-banding and nuclear DNA amount in six *Bromus* species. *Acta Biol. Crac. Ser. Bot.*, **43**: 105–115
- Leitch, J. and M. D. Bennett 1997 Polyploidy in angiosperms. *Trends Plant Sci.*, **2**: 470–475
- O'Brien, I. E. W., D. R. Smith, R. C. Gardner and B. G. Murray 1996 Flow cytometric determination of genome size in *Pinus*. *Plant Sci.*, **115**: 91–99
- Okazaki, K., K. Kurimoto, I. Miyajima, A. Enami, H. Mizuochi, Y. Matumoto and H. Ohya 2005 Induction of 2n pollen in tulips by arresting the meiotic process with nitrous oxide gas. *Euphytica*, **143**: 101–114
- Ollitrault-Sammarcelli, F., J. M. Legave, N. Michaux-Ferriere and A. M. Hirsch 1994 Use of flow cytometry for rapid determination of ploidy level in the genus *Actinidia*. *Scientia Hort.*, **57**: 303–313
- Ozaki, Y., K. Narikyo, M. Hiramatsu, K. Ureshino and H. Okubo 1998 Application of flow cytometry for rapid determination of ploidy levels in asparagus (*Asparagus officinalis* L.). *J. Fac. Agr., Kyushu Univ.*, **43**: 83–88
- Pedrosa, A., M. F. Jantsch, E. A. Moscone, P. F. Ambros and D. Schweizer 2001 Characterisation of pericentromeric and sticky intercalary heterochromatin in *Ornithogalum longibracteatum* (Hyacinthaceae). *Chromosoma*, **110**: 203–213
- Petersen, K. K., P. Hagberg and K. Kristiansen 2003 Colchicine and oryzalin mediated chromosome doubling in different genotypes of *Miscanthus sinensis*. *Plant Cell, Tiss. Org. Cult.*, **73**: 137–146
- Pinheiro, A. A., M. T. Pozzobon and V. T. C. Carneiro 2000 Duplication of the chromosome number of diploid *Brachiaria brizantha* plants using colchicine. *Plant Cell Rep.*, **19**: 274–278
- Praça, M. M., C. R. Carvalho and W. R. Clarindo 2009 A practical and reliable procedure for *in vitro* induction of tetraploid tomato. *Scientia Hort.*, **122**: 501–505
- Rubuluza, T., R. V. Nikolova, M. T. Smith and K. Hannweg 2007 *In vitro* induction of tetraploids in *Colophospermum mopane* by colchicines. *S. Afr. J. Bot.*, **73**: 259–261
- Shao, J., C. Chen and X. Deng 2003 *In vitro* induction of tetraploid in pomegranate (*Punica granatum*). *Plant Cell, Tiss. Org. Cult.*, **75**: 241–246
- Sgorbati, S., M. Levi, E. Sparvoli, F. Trezzi and G. Lucchini 1986 Cytometry and flow cytometry of 4',6-diamidino-2-phenylidole (DAPI) stained suspensions of nuclei released from fresh and fixed tissues of plants. *Physiol. Plant.*, **68**: 471–476
- Śliwińska, E. and P. Steen 1995 Flow cytometric estimation of ploidy in anisoploid sugar-beet populations. *J. Appl. Genet.*, **36**: 111–118
- Thao, N. T. P., K. Ureshino, I. Miyajima, Y. Ozaki and H. Okubo 2003 Induction of tetraploids in ornamental *Alocasia* through colchicine and oryzalin treatments. *Plant Cell, Tiss. Org. Cult.*, **72**: 19–25
- Thao, N. T. P., Y. Ozaki and H. Okubo 2004 Colchicine and oryzalin induced tetraploids in Ornamental *Alocasia* × *amazonica* hort. *J. Japan. Soc. Hort. Sci.*, **73**: 63–65
- Thiem, B. and E. Śliwińska 2003 Flow cytometric analysis of nuclear DNA content in cloudberry (*Rubus chamaemorus* L.) *in vitro* cultures. *Plant Sci.*, **164**: 129–134
- Tosca, A., R. Pandolfi, S. Citterio, A. Fasoli and S. Sgorbati 1995 Determination of flow cytometry of the chromosome doubling capacity of oryzalin and colchicine in gynogenetic haploids of gerbera. *Plant Cell Rep.*, **14**: 455–458
- Vainola, A. 2000 *Polyplodization* and early screening of *Rhododendron* hybrids. *Euphytica*, **112**: 239–244
- Wang, Y. T. 2004 Flourishing market for potted orchids. *Flower Tech.*, **7**: 2–5
- Wongprichachan, P., K.-L. Huang, Y.-M. Chou, S.-T. Hsu, T.-Y. Liu and H. Okubo 2013 Induction of unreduced gamete in *Phalaenopsis* by N₂O treatments. *J. Fac. Agr., Kyushu Univ.*, **58**: 27–31