

Protein disulfide isomerase like 1-1 participates in the maturation of proglutelin within the endoplasmic reticulum in rice endosperm

Satoh-Cruz, Mio

Faculty of Agriculture, Kyushu University | Institute of Biological Chemistry, Washington State University

Crofts, Andrew J.

International Liberal Arts Program, Akita International University | Institute of Biological Chemistry, Washington State University

Takemoto-Kuno, Yoko

National Institute of Crop Science | Faculty of Agriculture, Kyushu University

Sugino, Aya

Faculty of Agriculture, Kyushu University | Institute of Biological Chemistry, Washington State University

他

<https://hdl.handle.net/2324/26299>

出版情報 : Plant and Cell Physiology. 51 (9), pp.1581-1593, 2010-09. Oxford University Press
バージョン :
権利関係 : (C) The Author 2010. Published by Oxford University Press on behalf of Japanese Society of Plant Physiologists.

Supplementary Fig. S1

cDNA and deduced amino acid sequence of PDIL1-1 in Kinmaze.

The deduced amino acid sequence is shown below the nucleotide sequence using the single-letter code. The stop codon is indicated with an asterisk. The putative signal peptide is highlighted. Thioredoxin active sites (-CXXC-), N-linked glycosylation site (-NXF/T-) and the C-terminal ER-retention signal (KDEL) are indicated.

Supplementary Fig. S2

Spatial expression of the various rice PDI homologs as assessed by RT-PCR.