

# Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations

Okita, Masatoshi  
Graduate School of Mathematics, Kyushu University

<https://hdl.handle.net/2324/27216>

---

出版情報 : MI Preprint Series. 2013-11, 2013-08-31. 九州大学大学院数理学研究院  
バージョン :  
権利関係 :



# **MI Preprint Series**

**Kyushu University  
The Global COE Program  
Math-for-Industry Education & Research Hub**

## **Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations**

**Masatoshi OKITA**

**MI 2013-11**

( Received August 31, 2013 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations

Masatoshi Okita

Graduate School of Mathematics,  
Kyushu University,  
Fukuoka 819-0395, Japan  
Email: m-okita@math.kyushu-u.ac.jp

**Abstract:** In this paper we are concerned with the convergence rates of the global strong solution to motionless state with constant density for the compressible Navier-Stokes equations in the whole space  $\mathbb{R}^n$  for  $n \geq 3$ . It is proved that the perturbations decay in critical spaces, if the initial perturbations of density and velocity are small in  $B_{2,1}^{\frac{n}{2}}(\mathbb{R}^n) \cap \dot{B}_{1,\infty}^0(\mathbb{R}^n)$  and  $B_{2,1}^{\frac{n}{2}-1}(\mathbb{R}^n) \cap \dot{B}_{1,\infty}^0(\mathbb{R}^n)$ , respectively.

**Key Words:** compressible Navier- Stokes equations; convergence rate.

2010 Mathematics Subject Classification Numbers. 35Q30, 76N15.

## 1 Introduction

This paper studies the initial value problem for the compressible Navier-Stokes equation in  $\mathbb{R}^n$  :

$$\begin{cases} \partial_t \rho + \nabla \cdot (\rho u) = 0, \\ \partial_t u + (u \cdot \nabla)u + \frac{\nabla P(\rho)}{\rho} = \frac{\mu}{\rho} \Delta u + \frac{\mu + \mu'}{\rho} \nabla(\nabla \cdot u), \\ (\rho, u)(0, x) = (\rho_0, u_0)(x). \end{cases} \quad (1)$$

Here  $t > 0$ ,  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ ; the unknown functions  $\rho = \rho(t, x) > 0$  and  $u = u(t, x) = (u_1(t, x), u_2(t, x), \dots, u_n(t, x))$  denote the density and velocity, respectively;  $P = P(\rho)$  is the pressure that are assumed to be a function of the density  $\rho$ ;  $\mu$  and  $\mu'$  are the viscosity coefficients satisfying the conditions  $\mu > 0$  and  $\mu' + \frac{2}{n}\mu \geq 0$ ; and  $\nabla \cdot$ ,  $\nabla$  and  $\Delta$  denote the usual divergence, gradient and Laplacian with respect to  $x$ , respectively.

We assume that  $P(\rho)$  is smooth in a neighborhood of  $\bar{\rho}$  with  $P'(\bar{\rho}) > 0$ , where  $\bar{\rho}$  is a given positive constant.

In this paper we derive the convergence rate of solution of problem (1) to the constant stationary solution  $(\bar{\rho}, 0)$  as  $t \rightarrow \infty$  when the initial perturbation  $(\rho_0 - \bar{\rho}, u_0)$  is sufficiently small in critical spaces and  $\dot{B}_{1,\infty}^0$  for  $n \geq 3$ .

Matsumura-Nishida [8] showed the global in time existence of solution of (1) for  $n = 3$ , provided that the initial perturbation  $(\rho_0 - \bar{\rho}, u_0)$  is sufficiently small in  $H^3(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$ . Furthermore, the following decay estimates was obtained in [8];

$$\|\nabla^k(\rho - \bar{\rho}, u)(t)\|_{L^2} \leq C(1+t)^{-\frac{3}{4}-\frac{k}{2}} \quad k = 0, 1. \quad (2)$$

These results were proved by combining the energy method and the decay estimates of the semigroup  $E(t)$  generated by the linearized operator  $A$  at the constant state  $(\bar{\rho}, 0)$ .

On the other hand, Kawashita [6] showed the global existence of solution for initial perturbations sufficiently small in  $H^{s_0}(\mathbb{R}^n)$  with  $s_0 = [\frac{n}{2}] + 1$ ,  $n \geq 2$ . (Note that  $s_0 = 2$  for  $n = 3$ ). Wang-Tan [11] then considered the case  $n = 3$  when the initial perturbation  $(\rho_0 - \bar{\rho}, u_0)$  is sufficiently small in  $H^2(\mathbb{R}^3) \cap L^1(\mathbb{R}^3)$ , and proved the decay estimates (2). Okita [10] showed that if  $n \geq 2$  then the following estimates hold for the solution  $(\rho, u)$  of (1) :

$$\|\nabla^k(\rho - \bar{\rho}, u)(t)\|_{L^2} \leq C(1+t)^{-\frac{n}{4}-\frac{k}{2}} \quad k = 0, \dots, s_0,$$

provided that  $(\rho_0 - \bar{\rho}, u_0)$  is sufficiently small in  $H^{s_0}(\mathbb{R}^n) \cap L^1(\mathbb{R}^n)$  with  $s_0 = [\frac{n}{2}] + 1$ . This result was shown by decomposition of the perturbation into low and high frequency parts. Moreover Liang Li-Zhang [7] showed the density and momentum converge at the rates  $(1+t)^{\frac{3}{4}-\frac{s}{2}}$  in the  $L^2$ -norm, when initial perturbation sufficiently small in  $H^l(\mathbb{R}^3) \cap \dot{B}_{1,\infty}^{-s}(\mathbb{R}^3)$  with  $l \geq 4$  and  $s \in [0, 1]$ . Note that  $L^1$  is included in  $\dot{B}_{1,\infty}^0$ .

Danchin [2] proved the global existence in critical homogeneous Besov space, which is stated as follows.

**Proposition 1.1** (Danchin [2]). *Let  $n \geq 2$ . There are two positive constants  $\epsilon_1$  and  $M$  such that for all  $(\rho_0, u_0)$  with  $(\rho_0 - \bar{\rho}) \in \dot{B}_{2,1}^{\frac{n}{2}} \cap \dot{B}_{2,1}^{\frac{n}{2}-1}$ ,  $u_0 \in \dot{B}_{2,1}^{\frac{n}{2}-1}$  and*

$$\|\rho_0 - \bar{\rho}\|_{\dot{B}_{2,1}^{\frac{n}{2}} \cap \dot{B}_{2,1}^{\frac{n}{2}-1}} + \|u_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq \epsilon_1, \quad (3)$$

*then problem (1) has a unique global solution  $(\rho, u) \in C(\mathbb{R}^+; \dot{B}_{2,1}^{\frac{n}{2}} \cap \dot{B}_{2,1}^{\frac{n}{2}-1}) \times (L^1(\mathbb{R}^+; \dot{B}_{2,1}^{\frac{n}{2}+1}) \cap C(\mathbb{R}^+; \dot{B}_{2,1}^{\frac{n}{2}-1}))$  that satisfies the estimate*

$$\|\rho - \bar{\rho}\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \|u\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \int_0^\infty \|u\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} dt \leq M(\|\rho_0 - \bar{\rho}\|_{\dot{B}_{2,1}^{\frac{n}{2}} \cap \dot{B}_{2,1}^{\frac{n}{2}-1}} + \|u_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}).$$

For critical nonhomogeneous Besov space, Haspot [4] proved the local well-posedness.

**Proposition 1.2** (Haspot [4]). *Let  $n \geq 2$ . Let  $u_0 \in B_{2,1}^{\frac{n}{2}-1}$  and  $(\rho_0 - \bar{\rho}) \in B_{2,1}^{\frac{n}{2}}$  with  $\rho_0 - \bar{\rho} > 0$ . Then there exist a constant  $T > 0$  such that the problem (1) has a unique local solution  $(\rho, u)$  on  $[0, T]$  with  $\rho - \bar{\rho} > 0$  and:*

$$\rho - \bar{\rho} \in C([0, T]; B_{2,1}^{\frac{n}{2}}), \quad u \in (C([0, T]; B_{2,1}^{\frac{n}{2}-1}) \cap L^1(0, T; B_{2,1}^{\frac{n}{2}+1})).$$

We now state our main result of this paper which gives the optimal  $L^2$  decay rate for strong solutions in critical Besov spaces.

**Theorem 1.3.** *Assume that  $n \geq 3$ . There exists  $\epsilon > 0$  such that if  $u_0 \in B_{2,1}^{\frac{n}{2}-1} \cap \dot{B}_{1,\infty}^0$ ,  $(\rho_0 - \bar{\rho}) \in B_{2,1}^{\frac{n}{2}} \cap \dot{B}_{1,\infty}^0$  and*

$$\|\rho_0 - \bar{\rho}\|_{B_{2,1}^{\frac{n}{2}} \cap \dot{B}_{1,\infty}^0} + \|u_0\|_{B_{2,1}^{\frac{n}{2}-1} \cap \dot{B}_{1,\infty}^0} \leq \epsilon,$$

*then problem (1) has a unique global solution  $(\rho - \bar{\rho}, u) \in C(\mathbb{R}^+; B_{2,1}^{\frac{n}{2}}) \times (C(\mathbb{R}^+; B_{2,1}^{\frac{n}{2}-1}) \cap L^1(\mathbb{R}^+; \dot{B}_{2,1}^{\frac{n}{2}+1}))$ . Furthermore, there exists constant  $C_0 > 0$ , we have*

$$\|(\rho - \bar{\rho}, u)\|_{B_{2,1}^{\frac{n}{2}-1}} \leq C_0(1+t)^{-\frac{n}{4}} \quad (4)$$

*for  $t \geq 0$ .*

**Remark 1.4.** *If  $(\rho_0, u_0)$  satisfies the assumption of Theorem 1.3, then it also satisfies the assumption of Proposition 1.1. Therefore, we have estimate of (3).*

**Remark 1.5.** *We will derive the a priori estimate with time weight for  $\|(\rho(t) - \bar{\rho}, u(t))\|_{B_{2,1}^{\frac{n}{2}} \times B_{2,1}^{\frac{n}{2}-1}}$  which, together with Proposition 1.2, proves the global existence in nonhomogeneous critical Besov spaces and  $\dot{B}_{1,\infty}^0$ .*

**Remark 1.6.** *The convergence rates for the problem of (1) given in (2) are optimal. By  $B_{2,1}^{\frac{n}{2}-1} \subset L^2$ , the convergence rate of (4) are optimal.*

To prove Theorem 1.3, as in [5], we introduce a decomposition of the perturbation  $U(t) = (\rho - \bar{\rho}, u)(t)$  associated with the spectral properties of the linearized operator  $A$ . In the case of our problem, we simply decompose the perturbation  $U(t)$  into low and high frequency parts. As for the low frequency part, we apply the decay estimates for the low frequency part of  $E(t)$ ; while the high frequency part is estimated by using the energy method. One of the points of our approach is that by restricting the use of the decay estimates for  $E(t)$  to its low frequency part, one can avoid the derivative loss due to the convective term of the transport equation  $(1)_1$ . On the other hand, the convective term of  $(1)_1$  can be controlled by the energy method and commutator estimate which we apply to the high frequency part. Another point is that we have  $\int_0^\infty \|u\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} dt < C\epsilon$ , established in Proposition 1.1. We need this estimate when we estimate the nonlinear terms.

The paper is organized as follows. In Section 2 we introduce the notation and some properties of Besov spaces. In Section 3 we rewrite the system into the one for the perturbation and introduce auxiliary Lemmas used in this paper. In Section 4 we give the proof of Theorem 1.3.

## 2 Preliminaries

In this section we first introduce the notation which will be used throughout this paper. We then introduce Besov spaces and some properties of Besov spaces.

## 2.1 Notation

Let  $L^p(1 \leq p \leq \infty)$  denote the usual  $L^p$ -Lebesgue space on  $\mathbb{R}^n$ . For nonnegative integer  $m$ , we denote by  $H^m$  the usual  $L^2$ -Sobolev space of order  $m$ . The inner-product of  $L^2$  is denoted by  $(\cdot, \cdot)$ . If  $S$  is any nonempty set, sequence space  $l^p(S)$  denote the usual  $l^p$  sequence space on  $S$ .

For any integer  $l \geq 0$ ,  $\nabla^l f$  denotes all of  $l$ -th derivatives of  $f$ .

For a function  $f$ , we denote its Fourier transform by  $\mathfrak{F}[f] = \hat{f}$ :

$$\mathfrak{F}[f](\xi) = \hat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{-ix \cdot \xi} dx.$$

The inverse of  $\mathfrak{F}$  is denoted by  $\mathfrak{F}^{-1}[f] = \check{f}$ ,

$$\mathfrak{F}^{-1}[f](x) = \check{f}(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} f(\xi) e^{i\xi \cdot x} d\xi.$$

## 2.2 Besov spaces

Let us now define the homogeneous and nonhomogeneous Besov spaces. First we introduce the dyadic partition of unity. We can use for instance any  $(\phi, \chi) \in C^\infty$ , such that  $\phi$  is supported in  $\{\xi \in \mathbb{R}^n | \frac{3}{4} \leq |\xi| \leq \frac{8}{3}\}$ ,  $\chi$  is supported in  $\{\xi \in \mathbb{R}^n | |\xi| \leq \frac{4}{3}\}$  such that

$$\begin{aligned} \forall \xi \in \mathbb{R}^n, \quad \chi(\xi) + \sum_{j \geq 0} \phi(2^{-j}\xi) &= 1, \\ \forall \xi \in \mathbb{R}^n \setminus \{0\}, \quad \sum_{j \in \mathbb{Z}} \phi(2^{-j}\xi) &= 1, \end{aligned}$$

$$\begin{aligned} |j - j'| \geq 2 &\Rightarrow \text{Supp } \phi(2^{-j}\cdot) \cap \text{Supp } \phi(2^{-j'}\cdot) = \emptyset, \\ j \geq 1 &\Rightarrow \text{Supp } \chi \cap \text{Supp } \phi(2^{-j}\cdot) = \emptyset. \end{aligned}$$

Denoting  $h = \mathfrak{F}^{-1}\phi$  and  $\tilde{h} = \mathfrak{F}^{-1}\chi$ , we then define the dyadic blocks by

$$\begin{aligned} \Delta_{-1}u &= \chi(D)u = \tilde{h} * u, \\ \Delta_j u &= \phi(2^{-j}D)u = 2^{jn} \int_{\mathbb{R}^n} h(2^j y) u(x - y) dy \quad \text{if } j \geq 0, \\ \dot{\Delta}_j u &= \phi(2^{-j}D)u = 2^{jn} \int_{\mathbb{R}^n} h(2^j y) u(x - y) dy \quad \text{if } j \in \mathbb{Z}. \end{aligned}$$

The low-frequency cut-off operator is defined by

$$S_j u = \sum_{-1 \leq k \leq j-1} \Delta_k u, \quad \dot{S}_j u = \sum_{k \leq j-1} \dot{\Delta}_k u.$$

Obviously we can write that:  $Id = \sum_j \Delta_j$ . The high-frequency cut-off operator  $\tilde{S}_j$  is defined by

$$\tilde{S}_j = Id - S_j = \sum_{k \geq j} \Delta_k u,$$

To begin, we define Besov spaces.

**Definition 1.** For  $s \in \mathbb{R}$  and  $1 \leq p, r \leq \infty$ , and  $u \in \mathcal{S}'$  we set

$$\|u\|_{B_{p,r}^s} := \|2^{js} \|\Delta_j u\|_{L^p}\|_{l^r(\{k \geq -1\})},$$

$$\|u\|_{\dot{B}_{p,r}^s} := \|2^{js} \|\dot{\Delta}_j u\|_{L^p}\|_{l^r(\mathbb{Z})}.$$

The nonhomogeneous Besov space  $B_{p,r}^s$  and the homogeneous Besov space  $\dot{B}_{p,r}^s$  are set of function  $u \in \mathcal{S}'$  such that  $\|u\|_{B_{p,r}^s}, \|u\|_{\dot{B}_{p,r}^s} < \infty$  respectively.

Let us state some basic Lemmas for Besov spaces.

**Lemma 2.1.** *The following properties hold:*

- (i)  $\|\nabla \Delta_{-1} u\|_{L^2} \leq C \|\Delta_{-1} u\|_{L^2}.$
- (ii)  $C^{-1} 2^j \|\dot{\Delta}_j u\|_{L^2} \leq \|\nabla \dot{\Delta}_j u\|_{L^2} \leq C 2^j \|\dot{\Delta}_j u\|_{L^2} \quad (j \in \mathbb{Z}).$
- (iii)  $\|\nabla S_j u\|_{L^2} \leq C 2^j \|S_j u\|_{L^2} \quad (j \geq 0).$
- (iv)  $\|\tilde{S}_j u\|_{L^2} \leq C 2^j \|\nabla \tilde{S}_j u\|_{L^2} \quad (j \geq 0).$

The assertions (i), (ii), (iii) and (iv) easily follow from the Plancherel theorem.

**Lemma 2.2.** *The following properties hold:*

- (i)  $C^{-1} \|u\|_{\dot{B}_{p,r}^s} \leq \|\nabla u\|_{\dot{B}_{p,r}^{s-1}} \leq C \|u\|_{\dot{B}_{p,r}^s}.$
- (ii)  $\|\nabla u\|_{B_{p,r}^{s-1}} \leq C \|u\|_{B_{p,r}^s}.$
- (iii) *If  $s' > s$  or if  $s' = s$  and  $r_1 \leq r$  then  $B_{p,r_1}^{s'} \subset B_{p,r}^s$ .*
- (iv) *If  $r_1 \leq r$  then  $\dot{B}_{p,r_1}^s \subset \dot{B}_{p,r}^s$ .*
- (v) *Let  $\Lambda := \sqrt{-\Delta}$  and  $t \in \mathbb{R}$ . Then the operator  $\Lambda^t$  is an isomorphism from  $\dot{B}_{2,1}^s$  to  $\dot{B}_{2,1}^{s-t}$ .*

See, e.g., [2] [3] and [4] for the proof.

**Lemma 2.3.** *The following properties hold:*

- (i)  $\|u\|_{L^\infty} \leq C \|u\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \quad (\dot{B}_{2,1}^{\frac{n}{2}} \subset L^\infty).$
- (ii)  $\dot{B}_{1,1}^0 \subset L^1 \subset \dot{B}_{1,\infty}^0.$
- (iii)  $B_{2,2}^s \approx H^s.$
- (iv)  $B_{p,r}^s \subset \dot{B}_{p,r}^s \quad (s > 0).$

See, e.g., [2] [3] and [4] for the proof.

### 3 Reformulation of the problem

In this section we first rewrite system (1) into the one for the perturbation. We then introduce some auxiliary lemmas which will be useful in the proof of the main result.

Let us rewrite the problem (1). We define  $\mu_1, \mu_2$  and  $\gamma$  by

$$\mu_1 = \frac{\mu}{\bar{\rho}}, \quad \mu_2 = \frac{\mu + \mu'}{\bar{\rho}}, \quad \gamma = \sqrt{P'(\bar{\rho})}.$$

By using the new unknown function

$$\sigma(t, x) = \frac{\rho(t, x) - \bar{\rho}}{\bar{\rho}}, \quad w(t, x) = \frac{1}{\gamma} u(t, x),$$

the initial value problem (1) is reformulated as

$$\begin{cases} \partial_t \sigma + \gamma \nabla \cdot w = F_1(U), \\ \partial_t w - \mu_1 \Delta w - \mu_2 \nabla(\nabla \cdot w) + \gamma \nabla \sigma = F_2(U), \\ (\sigma, w)(0, x) = (\sigma_0, w_0)(x), \end{cases} \quad (5)$$

where,  $U = \begin{pmatrix} \sigma \\ w \end{pmatrix}$ ,

$$F_1(U) = -\gamma(w \cdot \nabla \sigma + \sigma \nabla \cdot w),$$

$$\begin{aligned} F_2(U) &= -\gamma(w \cdot \nabla)w - \mu_1 \frac{\sigma}{\sigma + 1} \Delta w - \mu_2 \frac{\sigma}{\sigma + 1} \nabla(\nabla \cdot w) \\ &\quad + \left( \frac{\bar{\rho}\gamma}{\sigma + 1} - \frac{\bar{\rho}}{\gamma} \frac{\int_0^1 P''(s\bar{\rho}\sigma + \bar{\rho})ds}{\sigma + 1} \right) \sigma \nabla \sigma. \end{aligned}$$

We set

$$A = \begin{pmatrix} 0 & -\gamma \nabla \cdot \\ -\gamma \nabla & \mu_1 \Delta + \mu_2 \nabla \nabla \cdot \end{pmatrix}.$$

By using operator  $A$ , problem (5) is written as

$$\partial_t U - AU = F(U), \quad U|_{t=0} = U_0, \quad (6)$$

where

$$F(U) = \begin{pmatrix} F_1(U) \\ F_2(U) \end{pmatrix}, \quad U_0 = \begin{pmatrix} \sigma_0 \\ w_0 \end{pmatrix}.$$

We introduce a semigroup associated with  $A$ . We set

$$E(t)u := \mathfrak{F}^{-1}[e^{\hat{A}(\xi)t}\hat{u}] \quad \text{for } u \in L^2,$$

where

$$\hat{A}(\xi) = \begin{pmatrix} 0 & -i\gamma\xi^t \\ -i\gamma\xi & -\mu_1|\xi|^2 I_n - \mu_2\xi\xi^t \end{pmatrix}.$$

Here and in what follows the superscript  $\cdot^t$  means the transposition.

We next state some basic Lemmas.



**Lemma 3.1.** *Let  $s_1, s_2 \leq \frac{n}{2}$  such that  $s_1 + s_2 > 0$ ,  $u \in \dot{B}_{2,1}^{s_1}$  and  $v \in \dot{B}_{2,1}^{s_2}$ . Then  $uv \in \dot{B}_{2,1}^{s_1+s_2-\frac{n}{2}}$  and*

$$\|uv\|_{\dot{B}_{2,1}^{s_1+s_2-\frac{n}{2}}} \leq C\|u\|_{\dot{B}_{2,1}^{s_1}}\|v\|_{\dot{B}_{2,1}^{s_2}}.$$

See, e.g., [1], for the proof.

**Lemma 3.2.** *Let  $s > 0$  and  $u \in \dot{B}_{2,1}^s \cap L^\infty$ . Let  $F \in W_{loc}^{[s]+2,\infty}(\mathbb{R}^n)$  such that  $F(0) = 0$ . Then  $F(u) \in \dot{B}_{2,1}^s$ . Moreover, there exists a function  $C_1$  of one variable depending only on  $s, n$  and  $F$ , and such that*

$$\|F(u)\|_{\dot{B}_{2,1}^s} \leq C_1(\|u\|_{L^\infty})\|u\|_{\dot{B}_{2,1}^s}.$$

See, e.g., [2], for the proof.

**Lemma 3.3.** (i) *Let  $a, b > 0$  satisfying  $\max\{a, b\} > 1$ . Then*

$$\int_0^t (1+s)^{-a}(1+t-s)^{-b}ds \leq C(1+t)^{-\min\{a,b\}}, \quad t \geq 0.$$

(ii) *Let  $a, b > 0$  and  $f \in L^1(0, \infty)$ . Then*

$$\int_0^t (1+t)^{-a}(1+t-s)^{-b}f(s)ds \leq C(1+t)^{-\min\{a,b\}} \int_0^t |f(s)|ds, \quad t \geq 0.$$

For the proof of (i), see [9]. Proof of (ii) can be proved using Hölder inequality.

## 4 Proof of main result

In this section we prove Theorem 1.3. In subsections 4.1 and 4.2 we establish the necessary estimates for  $\Delta_{-1}U(t)$  and  $\Delta_j U(t)$  for  $j \geq 0$ , respectively. In subsection 4.3 we derive the a priori estimate to complete the proof of Theorem 1.3.

**Proposition 4.1.** *Let  $T > 0$  and let  $(\sigma, w)$  be a solution of problem (6) on  $[0, T]$  such that*

$$\sigma \in C([0, T]; B_{2,1}^{\frac{n}{2}}), w \in C([0, T]; B_{2,1}^{\frac{n}{2}}) \cap L^1(0, T; B_{2,1}^{\frac{n}{2}+1}), \quad (7)$$

*Then,  $\Delta_j U(t) = (\Delta_j \sigma, \Delta_j w)^t$  for  $j \geq -1$  satisfy*

$$\partial_t \Delta_j U - A \Delta_j U = \Delta_j F(U), \quad (8)$$

$$\Delta_j U|_{t=0} = \Delta_j U_0. \quad (9)$$

*Moreover,  $\Delta_j U(t)$  for  $j \geq -1$  satisfy*

$$\Delta_j U(t) = E(t) \Delta_j U_0 + \int_0^t E(t-s) \Delta_j F(U)(s) ds. \quad (10)$$

**Proof.** Let  $U(t) = (\sigma, w)^t$  be a solution of (6) satisfying (7). Since  $\Delta_j AU = A\Delta_j U$ , applying  $\Delta_j$  to (6), we obtain

$$\begin{cases} \partial_t \Delta_j U - A\Delta_j U = \Delta_j F(U), \\ \Delta_j U|_{t=0} = \Delta_j U_0. \end{cases} \quad (11)$$

It then follows that

$$\Delta_j U(t) = E(t)\Delta_j U_0 + \int_0^t E(t-s)\Delta_j F(U)(s)ds.$$

This completes the proof. □

Set

$$\begin{aligned} M_1(t) &:= \sup_{0 \leq \tau \leq t} (1 + \tau)^{\frac{n}{4}} \|\Delta_{-1} U(\tau)\|_{L^2}, \\ M_\infty(t) &:= \sup_{0 \leq \tau \leq t} (1 + \tau)^{\frac{n}{4}} \sum_{j=0}^{\infty} 2^{(\frac{n}{2}-1)j} \{ \|\Delta_j U(\tau)\|_{L^2} + 2^j \|\Delta_j \sigma\|_{L^2} \}, \\ M(t) &:= M_1(t) + M_\infty(t). \end{aligned}$$

## 4.1 Estimate of low frequency parts

In this subsection we derive the estimate of  $\Delta_{-1} U(t)$ , in other words, we estimate  $M_1(t)$ .

**Lemma 4.2** (Matsumura-Nishida [9]). *(i) The set of all eigenvalues of  $\hat{A}(\xi)$  consists of  $\lambda_i(\xi)$  ( $i = 1, 2, 3$ ), where*

$$\begin{cases} \lambda_1(\xi) = \frac{-(\mu_1 + \mu_2)|\xi|^2 + i|\xi|\sqrt{4\gamma^2 - (\mu_1 + \mu_2)|\xi|^2}}{2}, \\ \lambda_2(\xi) = \overline{\lambda_1(\xi)}, \\ \lambda_3(\xi) = -\mu_1|\xi|^2, \end{cases}$$

for all  $\xi \in \mathbb{R}^n$ . Here  $\overline{\lambda_1(\xi)}$  denotes the complex conjugate of  $\lambda_1(\xi)$ .

*(ii)  $e^{t\hat{A}(\xi)}$  has the spectral resolution*

$$e^{t\hat{A}(\xi)} = \sum_{j=1}^3 e^{t\lambda_j(\xi)} P_j(\xi),$$

for all  $|\xi|$  except at most points of  $|\xi| > 0$ , where  $P_j(\xi)$  is the eigenprojection for  $\lambda_j(\xi)$  and  $P_j(\xi)$  satisfies

$$\|P_j(\xi)\| \leq C \quad (|\xi| \leq r).$$

where  $r = \frac{\gamma}{\sqrt{\mu_1 + \mu_2}}$ .

Moreover it has the estimate

$$\|e^{t\hat{A}(\xi)}\| \leq Ce^{-\beta t},$$

for all  $|\xi| \geq r$  and a positive constant  $\beta$ .

**Remark 4.3.** For each  $M > 0$  there exist  $C_2 = C_2(M) > 0$  and  $\beta_2 = \beta_2(M) > 0$  such that the estimate

$$\|e^{t\hat{A}(\xi)}\| \leq C_2 e^{-\beta_2|\xi|^2 t}$$

holds for  $|\xi| \leq M$  and  $t > 0$ .

**Lemma 4.4.**  $E(t)$  satisfies the estimate,

$$\|E(t)\Delta_{-1}U_0\|_{L^2} \leq C(1+t)^{-\frac{n}{4}}\|U_0\|_{\dot{B}_{1,\infty}^0}$$

for  $t \geq 0$ .

**Proof.** By Plancherel's theorem and Lemma 4.2 (ii), we have

$$\begin{aligned} \|E(t)\Delta_{-1}U_0(t)\|_{L^2} &\leq C\left(\int_{|\xi|\leq 2} |e^{\hat{A}(\xi)t}\hat{U}_0|^2 d\xi\right)^{\frac{1}{2}} \\ &\leq C\left(\int_{0<|\xi|\leq 2} e^{-\beta|\xi|^2 t} |\hat{U}_0(\xi)|^2 d\xi\right)^{\frac{1}{2}} \\ &\leq C\sum_{j\leq 2} (\|\Delta_j U_0\|_{L^1} \|\tilde{\Delta}_j U_0\|_{L^1} \int_{2^{j-1}<|\xi|\leq 2^{j+1}} e^{-\beta|\xi|^2 t} d\xi)^{\frac{1}{2}} \\ &\leq Ct^{-\frac{n}{4}}\|U_0\|_{\dot{B}_{1,\infty}^0}, \end{aligned} \tag{12}$$

where  $\tilde{\Delta}_j := \Delta_{j-1} + \Delta_j + \Delta_{j+1}$ .

We also find that

$$\begin{aligned} \|E(t)\Delta_{-1}U_0\|_{L^2} &\leq C\sum_{j\leq 2} (\|\Delta_j U_0\|_{L^1} \|\tilde{\Delta}_j U_0\|_{L^1} \int_{2^{j-1}<|\xi|\leq 2^{j+1}} e^{-\beta|\xi|^2 t} d\xi)^{\frac{1}{2}} \\ &\leq C\|U_0\|_{\dot{B}_{1,\infty}^0}. \end{aligned} \tag{13}$$

The estimate of Lemma 4.4 follows from (12) and (13).  $\square$

As for  $M_1(t)$ , we show the following estimate.

**Proposition 4.5.** Let  $n \geq 3$ . There exists a  $\epsilon > 0$  such that if

$$\|w_0\|_{B_{2,1}^{\frac{n}{2}-1}} + \|\sigma_0\|_{B_{2,1}^{\frac{n}{2}}} \leq \epsilon,$$

then there exists a constant  $C > 0$  independent of  $T$  such that

$$M_1(t) \leq C\|U_0\|_{\dot{B}_{1,\infty}^0} + C\epsilon M(t) + CM^2(t)$$

for  $t \in [0, T]$

To prove Proposition 4.5, we will use the following estimates on  $F(U)$ .

**Lemma 4.6.** *There exists a  $\epsilon > 0$  such that if*

$$\|w_0\|_{B_{2,1}^{\frac{n}{2}-1}} + \|\sigma_0\|_{B_{2,1}^{\frac{n}{2}}} \leq \epsilon,$$

*then there exists a constant  $C > 0$  independent of  $T$  such that*

$$\|F(U)\|_{\dot{B}_{1,\infty}^0} \leq C(1+t)^{-\frac{n}{4}}M(t)f(t) + C(1+t)^{-\frac{n}{2}}M^2(t)$$

*for  $t \in [0, T]$ , where  $0 \leq f \leq \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} \in L^1(0, \infty)$ .*

We will prove Lemma 4.6 later. Now we prove Proposition 4.5.

**Proof of Proposition 4.5.** By Lemma 4.4 and (10), we see that

$$\begin{aligned} \|\Delta_{-1}U(\tau)\|_{L^2} &\leq \|E(\tau)\Delta_{-1}U_0\|_{L^2} + \int_0^\tau \|E(\tau-s)\Delta_{-1}F(U(s))\|_{L^2}ds \\ &\leq C(1+\tau)^{-\frac{n}{4}}\|U_0\|_{\dot{B}_{1,\infty}^0} + \int_0^\tau (1+\tau-s)^{-\frac{n}{4}}\|F(U(s))\|_{\dot{B}_{1,\infty}^0}ds. \end{aligned} \quad (14)$$

Using Lemma 4.6, we have

$$\begin{aligned} &\int_0^\tau (1+\tau-s)^{-\frac{n}{4}}\|F(U(s))\|_{\dot{B}_{1,\infty}^0}ds \\ &\leq C \int_0^\tau (1+\tau-s)^{-\frac{n}{4}}\{(1+s)^{-\frac{n}{4}}f(s)M(t) + (1+s)^{-\frac{n}{2}}M^2(t)\}ds \\ &\leq CM(t) \int_0^\tau (1+\tau-s)^{-\frac{n}{4}}(1+s)^{-\frac{n}{4}}f(s)ds \\ &\quad + CM^2(t) \int_0^\tau (1+\tau-s)^{-\frac{n}{4}}(1+s)^{-\frac{n}{2}}ds \\ &\leq C(1+\tau)^{-\frac{n}{4}}\epsilon M(t) + C(1+\tau)^{-\frac{n}{4}}M^2(t). \end{aligned} \quad (15)$$

Here we used Lemma 3.3, in other words, we used  $\frac{n}{2} > 1$  for  $n \geq 3$  and  $\int_0^\tau f(s)ds \leq M\epsilon$ . By (14) and (15), we obtain

$$\|\Delta_{-1}U(\tau)\|_{L^2} \leq C(1+\tau)^{-\frac{n}{4}}\|U_0\|_{\dot{B}_{1,\infty}^0} + C\epsilon(1+\tau)^{-\frac{n}{4}}M(t) + C(1+\tau)^{-\frac{n}{4}}M^2(t),$$

and hence,

$$(1+\tau)^{\frac{n}{4}}\|\Delta_{-1}U(\tau)\|_2 \leq C\|U_0\|_{\dot{B}_{1,\infty}^0} + C\epsilon M(t) + CM^2(t).$$

Taking the supremum in  $\tau \in [0, t]$ , we obtain the desired estimate. □

It remains to prove Lemma 4.6.

**Proof of Lemma 4.6.** Since  $L^1 \subset \dot{B}_{1,\infty}^0$ , it suffices to estimate  $\|F(U)\|_{L^1}$ . By the Hölder inequality, we have

$$\begin{aligned} \|F(U)\|_{L^1} \leq & C\{\|w\|_{L^2}\|\nabla\sigma\|_{L^2} + \|\sigma\|_{L^2}\|\nabla w\|_{L^2} + \|w\|_{L^2}\|\nabla w\|_{L^2} \\ & + \|\sigma\|_{L^2}\|\nabla^2 w\|_{L^2} + \|\sigma\|_{L^2}\|\nabla\sigma\|_{L^2}\}, \end{aligned}$$

We see from  $B_{2,1}^s \subset L^2$  ( $s \geq 0$ ) that

$$\|w\|_{L^2}\|\nabla\sigma\|_{L^2} \leq C\|w\|_{B_{2,1}^{\frac{n}{2}-1}}\|\nabla\sigma\|_{B_{2,1}^{\frac{n}{2}-1}} \leq C(1+s)^{-\frac{n}{2}}M^2(t),$$

$$\begin{aligned} \|\sigma\|_{L^2}\|\nabla w\|_{L^2} & \leq C\|\sigma\|_{B_{2,1}^{\frac{n}{2}-1}}\{\|\Delta_{-1}w\|_{L^2} + \|\tilde{S}_0w\|_{B_{2,1}^{\frac{n}{2}}}\} \\ & \leq C\|\sigma\|_{B_{2,1}^{\frac{n}{2}-1}}\{\|w\|_{B_{2,1}^{\frac{n}{2}-1}} + \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}}\} \\ & \leq C(1+s)^{-\frac{n}{2}}M^2(t) + C(1+s)^{-\frac{n}{4}}M(t)f(t), \end{aligned}$$

where  $0 \leq f(t) \leq \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}}$ . As for  $\|\nabla^2 w\|_{L^2}$ , we get

$$\begin{aligned} \|\nabla^2 w\|_{L^2} & \leq C\{\|\Delta_{-1}w\|_{L^2} + \|\tilde{S}_0w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}}\} \\ & \leq C\{\|w\|_{B_{2,1}^{\frac{n}{2}-1}} + \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}}\}. \end{aligned}$$

The other terms are estimated similarly. Hence we have

$$\|F(U)\|_{\dot{B}_{1,\infty}^0} \leq C(1+s)^{-\frac{n}{2}}M^2(t) + C(1+s)^{-\frac{n}{4}}M(t)f(t).$$

This completes the proof. □

## 4.2 Estimate of high frequency parts

We next derive estimates for  $M_\infty$ . The system (8) is written as

$$\begin{cases} \partial_t \Delta_j \sigma + \gamma \nabla \cdot \Delta_j w = \Delta_j F_1(U), \\ \partial_t \Delta_j w - \mu_1 \Delta \Delta_j w - \mu_2 \nabla \cdot (\nabla \Delta_j w) + \gamma \nabla \Delta_j \sigma = \Delta_j F_2(U). \end{cases} \quad (16)$$

**Proposition 4.7.** *Let  $j \geq 0$ . There holds*

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \|\Delta_j U(t)\|_{L^2}^2 + \mu_1 \|\nabla \Delta_j w(t)\|_{L^2}^2 + \mu_2 \|\nabla \cdot \Delta_j w(t)\|_{L^2}^2 \\ & = (\Delta_j F_1(U), \Delta_j \sigma) + (\Delta_j F_2(U), \Delta_j w) \end{aligned} \quad (17)$$

for a.e.  $t \in [0, T]$ .

**Proof.** We take the inner product of  $(16)_1$  and  $(16)_2$  with  $\Delta_j \sigma$  and  $\Delta_j w$  respectively, integrating by parts and then adding them together, we obtain our proposition.  $\square$

For  $s \in \mathbb{R}$ , we denote  $\Lambda^s z := \mathfrak{F}^{-1}[|\xi|^s \hat{z}]$ . Let  $d = \Lambda^{-1} \nabla \cdot w$  be the "compressible part" of the velocity. Applying  $\Lambda^{-1} \nabla \cdot$  to  $(16)_2$ , system (16) writes

$$\begin{cases} \partial_t \Delta_j \sigma + \gamma \Lambda \Delta_j d = \Delta_j F_1(U), \\ \partial_t \Delta_j d - \nu \Delta \Delta_j d - \gamma \Lambda \Delta_j \sigma = \Lambda^{-1} \nabla \cdot \Delta_j F_2(U), \end{cases} \quad (18)$$

where we denote  $\nu = \mu_1 + \mu_2$ .

**Proposition 4.8.** *Let  $j \geq 0$ . There holds*

$$\begin{aligned} & \frac{1}{2} \frac{\nu}{\gamma} \frac{d}{dt} \|\Lambda \Delta_j \sigma\|_{L^2}^2 - \frac{d}{dt} (\Lambda \Delta_j \sigma, \Delta_j d) + \|\Lambda \Delta_j \sigma\|_{L^2}^2 = \gamma \|\Lambda \Delta_j d\|_{L^2}^2 \\ & - (\Lambda \Delta_j F_1(U), \Delta_j d) - (\Lambda^{-1} \nabla \cdot \Delta_j F_2(U), \Lambda \Delta_j \sigma) + \frac{\nu}{\gamma} (\Lambda \Delta_j F_1(U), \Lambda \Delta_j \sigma) \end{aligned} \quad (19)$$

for a.e.  $t \in [0, T]$ .

**Proof.** We apply  $\Lambda$  to the equation  $(18)_1$  and then take  $L^2$  inner product with  $\Delta_j d$ . We take  $L^2$  inner product of  $(18)_2$  with  $\Lambda \Delta_j \sigma$ . We also apply  $\Lambda$  to the equation  $(18)_1$  and take  $L^2$  inner product with  $\frac{\nu}{\gamma} \Delta_j \sigma$ . By a suitable linear combination of them, we obtain the desired identity of the proposition.  $\square$

We introduce a lemma for estimates of the right hande side.

**Lemma 4.9.** *The following inequalities hold*

$$|(\Lambda \Delta_j (w \cdot \nabla \sigma), \Lambda \Delta_j \sigma)| \leq C \alpha_j 2^{-(\frac{n}{2}-1)j} \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|\Lambda \Delta_j \sigma\|_{L^2},$$

$$\begin{aligned} & |(\Lambda \Delta_j (w \cdot \nabla \sigma), \Lambda \Delta_j d) + (\Delta_j (w \cdot \nabla d), \Lambda \Delta_j \sigma)| \\ & \leq C \alpha_j 2^{-(\frac{n}{2}-1)j} \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} (\|\Lambda \Delta_j \sigma\|_{L^2} \|d\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|\Delta_j d\|_{L^2}), \end{aligned}$$

where  $C$  depends on  $j$  and  $\|\{\alpha_j\}\|_{l^1} \leq 1$ .

See, e.g., [2], for the proof.

**Proposition 4.10.** *There exists a  $\epsilon > 0$  such that if*

$$\|w_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \|\sigma_0\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \leq \epsilon$$

then there holds

$$\begin{aligned} \frac{d}{dt} E_j(t) + c_0 E_j(t) &\leq C \{ \alpha_j (1+t)^{-\frac{n}{4}} M(t) f(t) \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j(\sigma \nabla \cdot w)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j(w \cdot \nabla d)\|_{L^2} \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_1(U)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_2(U)\|_{L^2} \}, \end{aligned} \quad (20)$$

for  $t \in [0, T]$  and  $j \geq 1$ , where  $\sum_{j \in \mathbb{Z}} \alpha_j \leq 1$ ,  $\int_0^\infty f(t) dt \leq \int_0^\infty \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} < C\epsilon$  and  $c_0$  is not depend on  $j$ . Here,  $E_j(t)$  is equivalent to  $2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2} + 2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2}$ . That is, there exists a  $D_1$  such that

$$\frac{1}{D_1} (2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2} + 2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2}) \leq E_j \leq D_1 (2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2} + 2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2}).$$

**Proof.** We add (17) to  $\kappa \times (19)$  with a constant  $\kappa > 0$  to be determined later. Then, we obtain

$$\begin{aligned} &\frac{d}{dt} \left\{ \frac{1}{2} \|\Delta_j U\|_{L^2}^2 + \frac{\kappa \nu}{2\gamma} \|\Lambda \Delta_j \sigma\|_{L^2}^2 - \kappa (\Lambda \Delta_j \sigma, \Delta_j d) \right\} \\ &\quad + \mu_1 \|\nabla \Delta_j w\|_{L^2}^2 + \mu_2 \|\nabla \cdot \Delta_j w\|_{L^2}^2 + \kappa \|\Lambda \Delta_j \sigma\|_{L^2}^2 \\ &= \gamma \kappa \|\Lambda \Delta_j w\|_{L^2}^2 + (\Delta_j F_1(U), \Delta_j \sigma) + (\Delta_j F_1(U), \Delta_j w) + \kappa \frac{\nu}{\gamma} (\Lambda \Delta_j F_1(U), \Lambda \Delta_j \sigma) \\ &\quad - \kappa (\Lambda \Delta_j F_1(U), \Delta_j d) - \kappa (\Lambda^{-1} \nabla \cdot \Delta_j F_2(U), \Lambda \Delta_j \sigma). \end{aligned} \quad (21)$$

We set

$$E_j^2(t) = 2^{2(\frac{n}{2}-1)j} \left\{ \frac{1}{2} \|\Delta_j U\|_{L^2}^2 + \frac{\kappa \nu}{2\gamma} \|\Lambda \Delta_j \sigma\|_{L^2}^2 - \kappa (\Lambda \Delta_j \sigma, \Delta_j d) \right\}.$$

For each  $\kappa \leq 1$ , there exists a  $D_1 > 3$  such that

$$E_j^2 \leq D_1^2 (2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2} + 2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2})^2.$$

By Cauchy's inequality with  $\delta$ , we have

$$\begin{aligned} &(2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2} + 2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2})^2 + D_1^2 \kappa 2^{2(\frac{n}{2}-1)j} (\Delta_j d, \Lambda \Delta_j \sigma) \\ &\leq 2 \{ (2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2})^2 + (2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2})^2 \} \\ &\quad + D_1^2 \kappa \delta (2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j \sigma\|_{L^2})^2 + D_1^2 \kappa \frac{1}{4\delta} (2^{(\frac{n}{2}-1)j} \|\Delta_j w\|_{L^2})^2. \end{aligned}$$

We select  $\delta = \frac{\nu}{4\gamma D_1}$  and  $\kappa$  is fixed in such a way that  $\kappa \leq \min\{\delta, \frac{\mu_1}{4\gamma}, 1\}$ . We then obtain

$$\begin{aligned} &\frac{1}{D_1^2} (2^{(\frac{n}{2}-1)j} \|\Delta_j U(t)\|_{L^2} + 2^{\frac{n}{2}j} \|\Delta_j \sigma\|_{L^2})^2 \\ &\leq 2^{2(\frac{n}{2}-1)j} \left\{ \frac{1}{2} \|\Delta_j U\|_{L^2}^2 + \frac{\kappa \nu}{2\gamma} \|\Lambda \Delta_j \sigma\|_{L^2}^2 - \kappa (\Lambda \Delta_j \sigma, \Delta_j d) \right\} = E_j^2. \end{aligned}$$

For  $j \geq 0$ , by Lemma 2.1, there exists a  $c_0 > 0$  such that

$$2c_0 E_j^2 \leq 2^{2(\frac{n}{2}-1)j} \{ \mu_1 \|\nabla \Delta_j w\|_{L^2}^2 + \mu_1 \|\nabla \cdot \Delta_j w\|_{L^2}^2 + \kappa \|\Lambda \Delta_j \sigma\|_{L^2}^2 - \gamma \kappa \|\Lambda \Delta_j w\|_{L^2}^2 \}.$$

Let us next estimate the right hande side of  $2^{2(\frac{n}{2}-1)j} \times (21)$ . By Hölder's inequality, we obtain

$$\begin{aligned} 2^{2(\frac{n}{2}-1)j} (\Delta_j F_1(U), \Delta_j \sigma) &\leq 2^{(\frac{n}{2}-1)j} \|\Delta_j F_1(U)\|_{L^2} 2^{(\frac{n}{2}-1)j} \|\Delta_j \sigma\|_{L^2}, \\ 2^{2(\frac{n}{2}-1)j} (\Delta_j F_2(U), \Delta_j \sigma) &\leq 2^{(\frac{n}{2}-1)j} \|\Delta_j F_2(U)\|_{L^2} 2^{(\frac{n}{2}-1)j} \|\Delta_j w\|_{L^2}, \\ 2^{2(\frac{n}{2}-1)j} (\Lambda^{-1} \nabla \cdot \Delta_j F_2(U), \Delta_j \sigma) &\leq 2^{(\frac{n}{2}-1)j} \|\Delta_j F_2(U)\|_{L^2} 2^{(\frac{n}{2}-1)j} \|\Delta_j \sigma\|_{L^2}. \end{aligned}$$

By Lemma 4.9 we have

$$\begin{aligned} &2^{2(\frac{n}{2}-1)j} (\Lambda \Delta_j F_1(U), \Lambda \Delta_j \sigma) \\ &= 2^{2(\frac{n}{2}-1)j} (\Lambda \Delta_j (w \cdot \nabla \sigma), \Lambda \Delta_j \sigma) + 2^{2(\frac{n}{2}-1)j} (\Lambda \Delta_j (\sigma \nabla \cdot w), \Lambda \Delta_j \sigma) \\ &\leq C \alpha_j \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} 2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j \sigma\|_{L^2} + 2^{2(\frac{n}{2}-1)j} \|\Lambda \Delta_j (\sigma \nabla \cdot w)\|_{L^2} \|\Lambda \Delta_j \sigma\|_{L^2}, \end{aligned}$$

and

$$\begin{aligned} &2^{2(\frac{n}{2}-1)j} (\Lambda \Delta_j F_1(U), \Delta_j d) \\ &= 2^{2(\frac{n}{2}-1)j} (\Lambda \Delta_j (w \cdot \nabla \sigma), \Delta_j d) + 2^{2(\frac{n}{2}-1)j} (\Lambda \Delta_j (\sigma \nabla \cdot w), \Delta_j d) \\ &\quad + 2^{2(\frac{n}{2}-1)j} (\Delta_j (w \cdot \nabla d), \Lambda \Delta_j \sigma) - 2^{2(\frac{n}{2}-1)j} (\Delta_j (w \cdot \nabla d), \Lambda \Delta_j \sigma) \\ &\leq C \alpha_j \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} (\|d\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} 2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j \sigma\|_{L^2} + \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} 2^{(\frac{n}{2}-1)j} \|\Delta_j d\|_{L^2}) \\ &\quad + 2^{2(\frac{n}{2}-1)j} \|\Delta_j (w \nabla \cdot d)\|_{L^2} \|\Lambda \Delta_j \sigma\|_{L^2} + 2^{2(\frac{n}{2}-1)j} \|\Lambda \Delta_j (\sigma \nabla \cdot w)\|_{L^2} \|\Delta_j d\|_{L^2}, \end{aligned}$$

where  $\sum_{j \in \mathbb{Z}} \alpha_j \leq 1$ . Hence we obtain

$$\begin{aligned} \frac{d}{dt} E_j^2 + 2c_0 E_j^2 &\leq C E_j \{ \alpha_j (1+t)^{-\frac{n}{4}} M(t) f(t) \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j (\sigma \nabla \cdot w)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j (w \cdot \nabla d)\|_{L^2} \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_1(U)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_2(U)\|_{L^2} \}. \end{aligned} \quad (22)$$

Let  $\delta_1 > 0$  be a small parameter (which will tend to 0) and denote  $H_j^2 = E_j^2 + \delta_1^2$ . From (22) and dividing by  $H_j$ , we gather

$$\begin{aligned} \frac{d}{dt} H_j + c_0 H_j &\leq C \{ \alpha_j (1+t)^{-\frac{n}{4}} M(t) f(t) \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j (\sigma \nabla \cdot w)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j (w \cdot \nabla d)\|_{L^2} \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_1(U)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_2(U)\|_{L^2} \} + c_0 \delta_1^2. \end{aligned}$$

Having  $\delta_1$  tend to 0, we get the desired result.  $\square$



### 4.3 Proof of Theorem 1.3.

**Proposition 4.11.** *There exists a constant  $\epsilon_2 > 0$  such that if*

$$\|U_0\|_{B_{2,1}^{\frac{n}{2}-1} \cap \dot{B}_{1,\infty}^0} + \|\sigma_0\|_{B_{2,1}^{\frac{n}{2}}} \leq \epsilon_2,$$

*then there holds*

$$M(t) \leq C \{ \|U_0\|_{B_{2,1}^{\frac{n}{2}-1} \cap \dot{B}_{1,\infty}^0} + \|\sigma_0\|_{B_{2,1}^{\frac{n}{2}}} \}$$

*for  $0 \leq t \leq T$ , where the constant  $C$  does not depend on  $T$ .*

**Proof.** By (20) we have

$$\begin{aligned} E_j(t) &\leq e^{-c_0 t} E_j(0) \\ &\quad + C \int_0^t e^{-c_0(t-\tau)} \{ \alpha_j (1+t)^{-\frac{n}{4}} M(t) f(t) + \alpha_j (1+t)^{-\frac{n}{2}} M^2(t) \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Lambda \Delta_j(\sigma \nabla \cdot w)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j(w \cdot \nabla d)\|_{L^2} \\ &\quad + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_1(U)\|_{L^2} + 2^{(\frac{n}{2}-1)j} \|\Delta_j F_2(U)\|_{L^2} \} d\tau, \end{aligned} \quad (23)$$

where  $\sum_{j=0}^{\infty} \alpha_j \leq 1$  and  $\int_0^{\infty} f(t) dt \leq \int_0^{\infty} \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} dt < C\epsilon_2$ . Hence summing up on  $j \geq 0$ , by the monotone convergence theorem, we obtain

$$\begin{aligned} \sum_{j=0}^{\infty} E_j(t) &\leq e^{-c_0 t} \sum_{j=0}^{\infty} E_j(0) \\ &\quad + C \int_0^t e^{-c_0(t-\tau)} \{ \alpha_j (1+t)^{-\frac{n}{4}} M(t) f(t) + (1+t)^{-\frac{n}{2}} M^2(t) \\ &\quad + \|\sigma \nabla \cdot w\|_{\dot{B}_{2,1}^{\frac{n}{2}}} + \|w \cdot \nabla d\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \|F_1(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \|F_2(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \} d\tau. \end{aligned}$$

We next estimate the right hand side. From Lemma 3.1 and Lemma 3.2, we have

$$\|\sigma \nabla \cdot w\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \leq C \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|\nabla \cdot w\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \leq C(1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau),$$

$$\|w \cdot \nabla d\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \|\nabla d\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \leq C(1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau).$$

Let us next consider  $\|F_1(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}$ ,  $\|F_2(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}$ :

$$\begin{aligned} \|w \cdot \nabla \sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} &\leq C \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|\nabla \sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \\ &\leq C (\|\Delta_{-1} w\|_{\dot{B}_{2,1}^{\frac{n}{2}}} + \|\tilde{S}_0 w\|_{\dot{B}_{2,1}^{\frac{n}{2}}}) \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \\ &\leq C(1+\tau)^{-\frac{n}{2}} M^2(\tau) + C(1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau), \end{aligned}$$

$$\|\sigma \nabla \cdot w\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \|\nabla w\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \leq C(1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau).$$

Hence, we obtain the estimate of  $\|F_1(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}$ . By using Lemma 3.1 and Lemma 3.2,  $\|F_2(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}$  is estimated as

$$\begin{aligned} \left\| \frac{\sigma}{\sigma+1} \Delta w \right\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} &\leq C \left\| \frac{\sigma}{\sigma+1} \right\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|\Delta w\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \\ &\leq C \|\sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}+1}} \leq C(1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau), \end{aligned}$$

$$\left\| \frac{\bar{\rho}\gamma}{\sigma+1} \sigma \nabla \sigma \right\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C \left\| \frac{\bar{\rho}\gamma}{\sigma+1} \sigma \right\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \|\nabla \sigma\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C(1+\tau)^{-\frac{n}{2}} M^2(\tau),$$

$$\|(w \cdot \nabla) w\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C \|w\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \|\nabla w\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} \leq C(1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau).$$

In the same way as we can obtain estimates of other terms on  $\|F_2(U)\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}}$ . Hence, by using Lemma 3.3, the integral of the right hand side of (23) is estimated as

$$\begin{aligned} &\int_0^t e^{-c_0(t-\tau)} \left\{ (1+\tau)^{-\frac{n}{2}} M^2(\tau) + (1+\tau)^{-\frac{n}{4}} M(\tau) f(\tau) \right\} d\tau \\ &\leq M(t) \int_0^t e^{-c_0(t-\tau)} (1+\tau)^{-\frac{n}{4}} f(\tau) d\tau + M^2(t) \int_0^t e^{-c_0(t-\tau)} (1+\tau)^{-\frac{n}{2}} d\tau \\ &\leq C(1+t)^{-\frac{n}{4}} \epsilon_2 M(t) + C(1+t)^{-\frac{n}{2}} M^2(t). \end{aligned}$$

Hence, we obtain

$$M_\infty(t) \leq C \left( \|U_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1}} + \|\sigma_0\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \right) + C\epsilon_2 M(t) + CM^2(t). \quad (24)$$

By Proposition 4.5 and (24), we have

$$M(t) \leq C \left( \|U_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1} \cap \dot{B}_{1,\infty}^0} + \|\sigma_0\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \right) + C\epsilon_2 M(t) + CM^2(t).$$

By taking  $\epsilon_2 > 0$  suitable small, we obtain

$$M(t) \leq C \left( \|U_0\|_{\dot{B}_{2,1}^{\frac{n}{2}-1} \cap \dot{B}_{1,\infty}^0} + \|\sigma_0\|_{\dot{B}_{2,1}^{\frac{n}{2}}} \right)$$

for all  $0 \leq t \leq T$ . □

It follows from Proposition 1.2 and Proposition 4.11 that

$$M(t) \leq C_3 \quad \text{for all } t.$$

Hence we obtain the desired decay estimate in Theorem 1.3.

**Acknowledgment:** The author would like to thank Professor Yoshiyuki Kagei and Professor Takayuki Kobayashi for their valuable advice.

## References

- [1] H. Bahouri, J. Y. Chemin, R. Danchin *Fourier analysis and nonlinear partial differential equation*, Springer, Heidelberg, 2011
- [2] R. Danchin, Global existence in critical spaces for compressible Navier-Stokes equations, *Invent. Math.* **141**, (2000) 579-614
- [3] R. Danchin, On the uniqueness in critical spaces for compressible Navier-Stokes equations, *Nonlinear Differ. Equ. Appl* **12** (2005) 111-128
- [4] B. Haspot, Well-posedness in critical spaces for the system of compressible Navier-Stokes in larger spaces, *J. Differential Equations* **251** (2011) 2262-2295
- [5] Y. Kagei, Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow, *Arch. Rational Mech. Anal.* **205** (2012), 585-650.
- [6] M. Kawashita, On global solution of Cauchy problems for compressible Navier-Stokes equation , *Nonlinear Analysis* **48** (2002), 1087-1105.
- [7] H-Liang. Li, T. Zhang, Large time behavior of isentropic compressible Navier-Stokes system in  $\mathbb{R}^3$ , *Math. Meth Appl. Sci* **34** (2011), 670-682.
- [8] A. Matsumura, T. Nishida, The initial value problem for the equation of motion of compressible viscous and heat-conductive fluids, *Proc. Japan Acad. Ser. A* **55** (1979), 337-342.
- [9] A. Matsumura, T. Nishida, The initial value problems for the equation of motion of compressible viscous and heat-conductive gases, *J. Math. Kyoto Univ.* **20** (1980), 67-104.
- [10] M. Okita, On the convergence rates for the compressible Navier- Stokes equations with potential force, to appear in *Kyushu J. Math.*
- [11] Y. Wang, Z. Tan, Global existence and optimal decay rate for the strong solution in  $H^2$  to the compressible Navier-Stokes equation, *Applied Mathematics Letters* **24** (2011), 1778-1784.

# List of MI Preprint Series, Kyushu University

## The Global COE Program

### Math-for-Industry Education & Research Hub

#### MI

- MI2008-1 Takahiro ITO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Abstract collision systems simulated by cellular automata
- MI2008-2 Eiji ONODERA  
The initial value problem for a third-order dispersive flow into compact almost Hermitian manifolds
- MI2008-3 Hiroaki KIDO  
On isosceles sets in the 4-dimensional Euclidean space
- MI2008-4 Hirofumi NOTSU  
Numerical computations of cavity flow problems by a pressure stabilized characteristic-curve finite element scheme
- MI2008-5 Yoshiyasu OZEKI  
Torsion points of abelian varieties with values in infinite extensions over a p-adic field
- MI2008-6 Yoshiyuki TOMIYAMA  
Lifting Galois representations over arbitrary number fields
- MI2008-7 Takehiro HIROTSU & Setsuo TANIGUCHI  
The random walk model revisited
- MI2008-8 Silvia GANDY, Masaaki KANNO, Hirokazu ANAI & Kazuhiro YOKOYAMA  
Optimizing a particular real root of a polynomial by a special cylindrical algebraic decomposition
- MI2008-9 Kazufumi KIMOTO, Sho MATSUMOTO & Masato WAKAYAMA  
Alpha-determinant cyclic modules and Jacobi polynomials
- MI2008-10 Sangyeol LEE & Hiroki MASUDA  
Jarque-Bera Normality Test for the Driving Lévy Process of a Discretely Observed Univariate SDE
- MI2008-11 Hiroyuki CHIHARA & Eiji ONODERA  
A third order dispersive flow for closed curves into almost Hermitian manifolds
- MI2008-12 Takehiko KINOSHITA, Kouji HASHIMOTO and Mitsuhiro T. NAKAO  
On the  $L^2$  a priori error estimates to the finite element solution of elliptic problems with singular adjoint operator
- MI2008-13 Jacques FARAUT and Masato WAKAYAMA  
Hermitian symmetric spaces of tube type and multivariate Meixner-Pollaczek polynomials

- MI2008-14 Takashi NAKAMURA  
Riemann zeta-values, Euler polynomials and the best constant of Sobolev inequality
- MI2008-15 Takashi NAKAMURA  
Some topics related to Hurwitz-Lerch zeta functions
- MI2009-1 Yasuhide FUKUMOTO  
Global time evolution of viscous vortex rings
- MI2009-2 Hidetoshi MATSUI & Sadanori KONISHI  
Regularized functional regression modeling for functional response and predictors
- MI2009-3 Hidetoshi MATSUI & Sadanori KONISHI  
Variable selection for functional regression model via the  $L_1$  regularization
- MI2009-4 Shuichi KAWANO & Sadanori KONISHI  
Nonlinear logistic discrimination via regularized Gaussian basis expansions
- MI2009-5 Toshiro HIRANOUCI & Yuichiro TAGUCHI  
Flat modules and Groebner bases over truncated discrete valuation rings
- MI2009-6 Kenji KAJIWARA & Yasuhiro OHTA  
Bilinearization and Casorati determinant solutions to non-autonomous 1+1 dimensional discrete soliton equations
- MI2009-7 Yoshiyuki KAGEI  
Asymptotic behavior of solutions of the compressible Navier-Stokes equation around the plane Couette flow
- MI2009-8 Shohei TATEISHI, Hidetoshi MATSUI & Sadanori KONISHI  
Nonlinear regression modeling via the lasso-type regularization
- MI2009-9 Takeshi TAKAISHI & Masato KIMURA  
Phase field model for mode III crack growth in two dimensional elasticity
- MI2009-10 Shingo SAITO  
Generalisation of Mack's formula for claims reserving with arbitrary exponents for the variance assumption
- MI2009-11 Kenji KAJIWARA, Masanobu KANEKO, Atsushi NOBE & Teruhisa TSUDA  
Ultradiscretization of a solvable two-dimensional chaotic map associated with the Hesse cubic curve
- MI2009-12 Tetsu MASUDA  
Hypergeometric  $q$ -functions of the  $q$ -Painlevé system of type  $E_8^{(1)}$
- MI2009-13 Hidenao IWANE, Hitoshi YANAMI, Hirokazu ANAI & Kazuhiro YOKOYAMA  
A Practical Implementation of a Symbolic-Numeric Cylindrical Algebraic Decomposition for Quantifier Elimination
- MI2009-14 Yasunori MAEKAWA  
On Gaussian decay estimates of solutions to some linear elliptic equations and its applications

- MI2009-15 Yuya ISHIHARA & Yoshiyuki KAGEI  
Large time behavior of the semigroup on  $L^p$  spaces associated with the linearized compressible Navier-Stokes equation in a cylindrical domain
- MI2009-16 Chikashi ARITA, Atsuo KUNIBA, Kazumitsu SAKAI & Tsuyoshi SAWABE  
Spectrum in multi-species asymmetric simple exclusion process on a ring
- MI2009-17 Masato WAKAYAMA & Keitaro YAMAMOTO  
Non-linear algebraic differential equations satisfied by certain family of elliptic functions
- MI2009-18 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of an Elliptical Flow Subjected to a Coriolis Force
- MI2009-19 Mitsunori KAYANO & Sadanori KONISHI  
Sparse functional principal component analysis via regularized basis expansions and its application
- MI2009-20 Shuichi KAWANO & Sadanori KONISHI  
Semi-supervised logistic discrimination via regularized Gaussian basis expansions
- MI2009-21 Hiroshi YOSHIDA, Yoshihiro MIWA & Masanobu KANEKO  
Elliptic curves and Fibonacci numbers arising from Lindenmayer system with symbolic computations
- MI2009-22 Eiji ONODERA  
A remark on the global existence of a third order dispersive flow into locally Hermitian symmetric spaces
- MI2009-23 Stjepan LUGOMER & Yasuhide FUKUMOTO  
Generation of ribbons, helicoids and complex scherk surface in laser-matter Interactions
- MI2009-24 Yu KAWAKAMI  
Recent progress in value distribution of the hyperbolic Gauss map
- MI2009-25 Takehiko KINOSHITA & Mitsuhiro T. NAKAO  
On very accurate enclosure of the optimal constant in the a priori error estimates for  $H_0^2$ -projection
- MI2009-26 Manabu YOSHIDA  
Ramification of local fields and Fontaine's property (Pm)
- MI2009-27 Yu KAWAKAMI  
Value distribution of the hyperbolic Gauss maps for flat fronts in hyperbolic three-space
- MI2009-28 Masahisa TABATA  
Numerical simulation of fluid movement in an hourglass by an energy-stable finite element scheme
- MI2009-29 Yoshiyuki KAGEI & Yasunori MAEKAWA  
Asymptotic behaviors of solutions to evolution equations in the presence of translation and scaling invariance

- MI2009-30 Yoshiyuki KAGEI & Yasunori MAEKAWA  
On asymptotic behaviors of solutions to parabolic systems modelling chemotaxis
- MI2009-31 Masato WAKAYAMA & Yoshinori YAMASAKI  
Hecke's zeros and higher depth determinants
- MI2009-32 Olivier PIRONNEAU & Masahisa TABATA  
Stability and convergence of a Galerkin-characteristics finite element scheme of lumped mass type
- MI2009-33 Chikashi ARITA  
Queueing process with excluded-volume effect
- MI2009-34 Kenji KAJIWARA, Nobutaka NAKAZONO & Teruhisa TSUDA  
Projective reduction of the discrete Painlevé system of type  $(A_2 + A_1)^{(1)}$
- MI2009-35 Yosuke MIZUYAMA, Takamasa SHINDE, Masahisa TABATA & Daisuke TAGAMI  
Finite element computation for scattering problems of micro-hologram using DtN map
- MI2009-36 Reiichiro KAWAI & Hiroki MASUDA  
Exact simulation of finite variation tempered stable Ornstein-Uhlenbeck processes
- MI2009-37 Hiroki MASUDA  
On statistical aspects in calibrating a geometric skewed stable asset price model
- MI2010-1 Hiroki MASUDA  
Approximate self-weighted LAD estimation of discretely observed ergodic Ornstein-Uhlenbeck processes
- MI2010-2 Reiichiro KAWAI & Hiroki MASUDA  
Infinite variation tempered stable Ornstein-Uhlenbeck processes with discrete observations
- MI2010-3 Kei HIROSE, Shuichi KAWANO, Daisuke MIIKE & Sadanori KONISHI  
Hyper-parameter selection in Bayesian structural equation models
- MI2010-4 Nobuyuki IKEDA & Setsuo TANIGUCHI  
The Itô-Nisio theorem, quadratic Wiener functionals, and 1-solitons
- MI2010-5 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and detecting change point via the relevance vector machine
- MI2010-6 Shuichi KAWANO, Toshihiro MISUMI & Sadanori KONISHI  
Semi-supervised logistic discrimination via graph-based regularization
- MI2010-7 Teruhisa TSUDA  
UC hierarchy and monodromy preserving deformation
- MI2010-8 Takahiro ITO  
Abstract collision systems on groups

- MI2010-9 Hiroshi YOSHIDA, Kinji KIMURA, Naoki YOSHIDA, Junko TANAKA & Yoshihiro MIWA  
An algebraic approach to underdetermined experiments
- MI2010-10 Kei HIROSE & Sadanori KONISHI  
Variable selection via the grouped weighted lasso for factor analysis models
- MI2010-11 Katsusuke NABESHIMA & Hiroshi YOSHIDA  
Derivation of specific conditions with Comprehensive Groebner Systems
- MI2010-12 Yoshiyuki KAGEI, Yu NAGAFUCHI & Takeshi SUDOU  
Decay estimates on solutions of the linearized compressible Navier-Stokes equation around a Poiseuille type flow
- MI2010-13 Reiichiro KAWAI & Hiroki MASUDA  
On simulation of tempered stable random variates
- MI2010-14 Yoshiyasu OZEKI  
Non-existence of certain Galois representations with a uniform tame inertia weight
- MI2010-15 Me Me NAING & Yasuhide FUKUMOTO  
Local Instability of a Rotating Flow Driven by Precession of Arbitrary Frequency
- MI2010-16 Yu KAWAKAMI & Daisuke NAKAJO  
The value distribution of the Gauss map of improper affine spheres
- MI2010-17 Kazunori YASUTAKE  
On the classification of rank 2 almost Fano bundles on projective space
- MI2010-18 Toshimitsu TAKAESU  
Scaling limits for the system of semi-relativistic particles coupled to a scalar bose field
- MI2010-19 Reiichiro KAWAI & Hiroki MASUDA  
Local asymptotic normality for normal inverse Gaussian Lévy processes with high-frequency sampling
- MI2010-20 Yasuhide FUKUMOTO, Makoto HIROTA & Youichi MIE  
Lagrangian approach to weakly nonlinear stability of an elliptical flow
- MI2010-21 Hiroki MASUDA  
Approximate quadratic estimating function for discretely observed Lévy driven SDEs with application to a noise normality test
- MI2010-22 Toshimitsu TAKAESU  
A Generalized Scaling Limit and its Application to the Semi-Relativistic Particles System Coupled to a Bose Field with Removing Ultraviolet Cutoffs
- MI2010-23 Takahiro ITO, Mitsuhiro FUJIO, Shuichi INOKUCHI & Yoshihiro MIZOGUCHI  
Composition, union and division of cellular automata on groups
- MI2010-24 Toshimitsu TAKAESU  
A Hardy's Uncertainty Principle Lemma in Weak Commutation Relations of Heisenberg-Lie Algebra



- MI2010-25 Toshimitsu TAKAESU  
On the Essential Self-Adjointness of Anti-Commutative Operators
- MI2010-26 Reiichiro KAWAI & Hiroki MASUDA  
On the local asymptotic behavior of the likelihood function for Meixner Lévy processes under high-frequency sampling
- MI2010-27 Chikashi ARITA & Daichi YANAGISAWA  
Exclusive Queueing Process with Discrete Time
- MI2010-28 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA  
Motion and Bäcklund transformations of discrete plane curves
- MI2010-29 Takanori YASUDA, Masaya YASUDA, Takeshi SHIMOYAMA & Jun KOGURE  
On the Number of the Pairing-friendly Curves
- MI2010-30 Chikashi ARITA & Kohei MOTEGI  
Spin-spin correlation functions of the  $q$ -VBS state of an integer spin model
- MI2010-31 Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling and spike detection via Gaussian basis expansions
- MI2010-32 Nobutaka NAKAZONO  
Hypergeometric  $\tau$  functions of the  $q$ -Painlevé systems of type  $(A_2 + A_1)^{(1)}$
- MI2010-33 Yoshiyuki KAGEI  
Global existence of solutions to the compressible Navier-Stokes equation around parallel flows
- MI2010-34 Nobushige KUROKAWA, Masato WAKAYAMA & Yoshinori YAMASAKI  
Milnor-Selberg zeta functions and zeta regularizations
- MI2010-35 Kissani PERERA & Yoshihiro MIZOGUCHI  
Laplacian energy of directed graphs and minimizing maximum outdegree algorithms
- MI2010-36 Takanori YASUDA  
CAP representations of inner forms of  $Sp(4)$  with respect to Klingen parabolic subgroup
- MI2010-37 Chikashi ARITA & Andreas SCHADSCHNEIDER  
Dynamical analysis of the exclusive queueing process
- MI2011-1 Yasuhide FUKUMOTO & Alexander B. SAMOKHIN  
Singular electromagnetic modes in an anisotropic medium
- MI2011-2 Hiroki KONDO, Shingo SAITO & Setsuo TANIGUCHI  
Asymptotic tail dependence of the normal copula
- MI2011-3 Takehiro HIROTSU, Hiroki KONDO, Shingo SAITO, Takuya SATO, Tatsushi TANAKA & Setsuo TANIGUCHI  
Anderson-Darling test and the Malliavin calculus
- MI2011-4 Hiroshi INOUE, Shohei TATEISHI & Sadanori KONISHI  
Nonlinear regression modeling via Compressed Sensing

- MI2011-5 Hiroshi INOUE  
Implications in Compressed Sensing and the Restricted Isometry Property
- MI2011-6 Daeju KIM & Sadanori KONISHI  
Predictive information criterion for nonlinear regression model based on basis expansion methods
- MI2011-7 Shohei TATEISHI, Chiaki KINJYO & Sadanori KONISHI  
Group variable selection via relevance vector machine
- MI2011-8 Jan BREZINA & Yoshiyuki KAGEI  
Decay properties of solutions to the linearized compressible Navier-Stokes equation around time-periodic parallel flow  
Group variable selection via relevance vector machine
- MI2011-9 Chikashi ARITA, Arvind AYYER, Kirone MALLICK & Sylvain PROLHAC  
Recursive structures in the multispecies TASEP
- MI2011-10 Kazunori YASUTAKE  
On projective space bundle with nef normalized tautological line bundle
- MI2011-11 Hisashi ANDO, Mike HAY, Kenji KAJIWARA & Tetsu MASUDA  
An explicit formula for the discrete power function associated with circle patterns of Schramm type
- MI2011-12 Yoshiyuki KAGEI  
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a parallel flow
- MI2011-13 Vladimír CHALUPECKÝ & Adrian MUNTEAN  
Semi-discrete finite difference multiscale scheme for a concrete corrosion model: approximation estimates and convergence
- MI2011-14 Jun-ichi INOGUCHI, Kenji KAJIWARA, Nozomu MATSUURA & Yasuhiro OHTA  
Explicit solutions to the semi-discrete modified KdV equation and motion of discrete plane curves
- MI2011-15 Hiroshi INOUE  
A generalization of restricted isometry property and applications to compressed sensing
- MI2011-16 Yu KAWAKAMI  
A ramification theorem for the ratio of canonical forms of flat surfaces in hyperbolic three-space
- MI2011-17 Naoyuki KAMIYAMA  
Matroid intersection with priority constraints
- MI2012-1 Kazufumi KIMOTO & Masato WAKAYAMA  
Spectrum of non-commutative harmonic oscillators and residual modular forms
- MI2012-2 Hiroki MASUDA  
Mighty convergence of the Gaussian quasi-likelihood random fields for ergodic Levy driven SDE observed at high frequency

- MI2012-3 Hiroshi INOUE  
A Weak RIP of theory of compressed sensing and LASSO
- MI2012-4 Yasuhide FUKUMOTO & Youich MIE  
Hamiltonian bifurcation theory for a rotating flow subject to elliptic straining field
- MI2012-5 Yu KAWAKAMI  
On the maximal number of exceptional values of Gauss maps for various classes of surfaces
- MI2012-6 Marcio GAMEIRO, Yasuaki HIRAOKA, Shunsuke IZUMI, Miroslav KRAMAR, Konstantin MISCHAIKOW & Vidit NANDA  
Topological Measurement of Protein Compressibility via Persistence Diagrams
- MI2012-7 Nobutaka NAKAZONO & Seiji NISHIOKA  
Solutions to a  $q$ -analog of Painlevé III equation of type  $D_7^{(1)}$
- MI2012-8 Naoyuki KAMIYAMA  
A new approach to the Pareto stable matching problem
- MI2012-9 Jan BREZINA & Yoshiyuki KAGEI  
Spectral properties of the linearized compressible Navier-Stokes equation around time-periodic parallel flow
- MI2012-10 Jan BREZINA  
Asymptotic behavior of solutions to the compressible Navier-Stokes equation around a time-periodic parallel flow
- MI2012-11 Daeju KIM, Shuichi KAWANO & Yoshiyuki NINOMIYA  
Adaptive basis expansion via the extended fused lasso
- MI2012-12 Masato WAKAYAMA  
On simplicity of the lowest eigenvalue of non-commutative harmonic oscillators
- MI2012-13 Masatoshi OKITA  
On the convergence rates for the compressible Navier- Stokes equations with potential force
- MI2013-1 Abuduwaili PAERHATI & Yasuhide FUKUMOTO  
A Counter-example to Thomson-Tait-Chetayev's Theorem
- MI2013-2 Yasuhide FUKUMOTO & Hirofumi SAKUMA  
A unified view of topological invariants of barotropic and baroclinic fluids and their application to formal stability analysis of three-dimensional ideal gas flows
- MI2013-3 Hiroki MASUDA  
Asymptotics for functionals of self-normalized residuals of discretely observed stochastic processes
- MI2013-4 Naoyuki KAMIYAMA  
On Counting Output Patterns of Logic Circuits
- MI2013-5 Hiroshi INOUE  
RIPless Theory for Compressed Sensing

- MI2013-6 Hiroshi INOUE  
Improved bounds on Restricted isometry for compressed sensing
- MI2013-7 Hidetoshi MATSUI  
Variable and boundary selection for functional data via multiclass logistic regression modeling
- MI2013-8 Hidetoshi MATSUI  
Variable selection for varying coefficient models with the sparse regularization
- MI2013-9 Naoyuki KAMIYAMA  
Packing Arborescences in Acyclic Temporal Networks
- MI2013-10 Masato WAKAYAMA  
Equivalence between the eigenvalue problem of non-commutative harmonic oscillators and existence of holomorphic solutions of Heun's differential equations, eigenstates degeneration, and Rabi's model
- MI2013-11 Masatoshi OKITA  
Optimal decay rate for strong solutions in critical spaces to the compressible Navier-Stokes equations