Establishment of the Optimal Conditions for Two-Dimensional Gel Electrophoresis of Papaya Seed Proteome

Roan, Su-Feng

Department of Horticulture and Biotechnology, Chinese Culture University

Hsiung, Tung-Chuan

Department of Horticulture and Biotechnology, Chinese Culture University

Yang, Kai-Yun

Department of Horticulture and Landscape Architecture, National Taiwan University

Liu, Wei-Ting

Department of Horticulture and Landscape Architecture, National Taiwan University

他

<https://doi.org/10.5109/27353>

出版情報：九州大学大学院農学研究院紀要. 58 (2), pp.239-246, 2013-09. Faculty of Agriculture, Kyushu University

バージョン：

権利関係：

Establishment of the Optimal Conditions for Two-Dimensional Gel Electrophoresis of Papaya Seed Proteome

Su-Feng ROAN¹, Tung-Chuan HSIUNG¹, Kai-Yun YANG², Wei-Ting LIU²,
Ing-Feng CHANG³, Chun-Ta WU^{2*} and Akira WAKANA

Laboratory of Horticultural Science, Division of Agronomy and Environmental Science,
Department of Agro-environmental Sciences, Faculty of Agriculture,
Kyushu University, Fukuoka 812-8581, Japan

(Received April 12, 2013 and accepted May 16, 2013)

Papaya seeds are considered as a recalcitrant material for protein preparation in two-dimensional gel electrophoresis (2-DE) because of abundance of interfering compounds. We examined two protein extraction methods, the classical trichloroacetic acid (TCA) / acetone precipitation (TCA-A) and phenol ammonium acetate-methanol precipitation (P-AA-M) methods, to evaluate their compatibility and efficiency of extracting papaya seed proteins in SDS-PAGE and two-dimensional (2D) gel electrophoresis analysis. Compared with TCA-A method, P-AA-M method showed better results, i.e., it increased at least five distinguished bands in SDS-PAGE, reduced problems of streaking and smearing in 2D gel, and enhanced resolution of detectable spots from 124 ± 27 to 717 ± 32 . Comparison between traditional Coomassie Brilliant Blue R-250 staining and zinc-reverse staining was also conducted. The result showed that seven times more protein spots could be identified from 2D gel stained with zinc-reverse stain procedure. The resolution of 2D gel analysis for papaya seed proteome was further improved by utilizing pH 4–7 isoelectric focusing gel instead of pH 3–10 one. The protein spots picked from 2D gel through the protocol that we established were compatible with identification assessments of LC-MS/MS. Our data will serve as a platform and reference for future research in proteomics of papaya seeds or other recalcitrant plant materials.

Key words: papaya seed, protein extraction, proteomics, two dimensional gel electrophoresis, zinc-reverse stain

INTRODUCTION

In proteomics research, a crucial step to obtain high resolution of the protein profile is to improve the quality of protein sample preparation (Görg *et al.*, 2000). Usually, plant tissues, unlike bacterial and animal cells, are rich in carbohydrates, organic acids, proteolytic and oxidative enzymes, and secondary metabolites such as pigments, phenolic compounds, and terpenes, which often interfere the resolution of protein separation on two-dimensional (2D) gel electrophoresis. For example, the high levels of secondary metabolites in distinct developmental stages, such as leaf senescence or fruit maturation, have strongly affected protein extraction and, as a consequence, interfered with the results of the 2D gel electrophoresis analysis with horizontal and vertical stripes, viz. streaking and smearing, as well as decrease in the resolution of the protein spots (Saravanan and Rose, 2004).

Typical method used for extracting plant proteins is grinding the plant tissue in liquid nitrogen in extraction buffer, followed by trichloroacetic acid (TCA) / acetone precipitation (Carpentier *et al.*, 2005). For woody plants and other plant tissues that are rich in polyphenol, the extraction by phenol is recommended because

the hydroxyl (OH-based) and protein-hydrophilic functional group of phenol could be used to separate the protein and the interfering substances within the tissue (Mijnsbrugge *et al.*, 2000; Wang *et al.*, 2003). The earliest phenol extraction was done by Hurkman and Tanaka (1986); the ammonium acetate / methanol solution was further used for protein precipitation prior to increase the number of protein spots and resolution on 2D gel electrophoresis.

Papaya (*Carica papaya* L.), an herbaceous perennial fruit crop belonging to the family Caricaceae, is commercially cultivated throughout the tropics and in many subtropical regions of the world. In the past decade, the global production, market demand, and international trade in papaya fruit have shown buoyant growth (Singh 2011). Unlike that most of fruit crops are propagated vegetatively, the propagule of papaya in commercial cultivation almost entirely depends on seeds (Paull and Duarte, 2011). Seed dormancy has long been an obstacle for the reproductive propagation in papaya; therefore, both illustration of the germination biology and establishment of the technologies for dormancy breakage have been extensively explored (Andreoli and Khan, 1993; Bhattacharya and Khuspe, 2001; Chacko and Singh, 1966; Salomao and Mundim, 2000; Wood *et al.*, 2000).

Proteomics has been widely adopted in the research of biology, medicine, and agriculture in recent years largely due to the advances in mass spectrometry and methodology development (Miernyk and Hajdich, 2011). Two-dimensional-gel basis proteomics has also been utilized in seed biology field to provide an overview of

¹ Department of Horticulture and Biotechnology, Chinese Culture University, Taipei, Taiwan

² Department of Horticulture and Landscape Architecture, National Taiwan University, Taipei, Taiwan

³ Institute of Plant Biology, National Taiwan University, Taipei, Taiwan

* Corresponding author (E-mail: wuct@ntu.edu.tw)

protein dynamics during seed development and germination as well as to elucidate the molecular mechanisms of postharvest treatments for the effects of seed vigor and longevity (Gallardo *et al.*, 2001; Miernyk and Hajduch, 2011; Pawlowski, 2010). Along with the information from completed genome sequences of various organisms, further understandings in physiological and molecular mechanisms and gene-related reactions can be established as systems in biology scale. The whole papaya sequence was completed in 2008 (Ming *et al.*, 2008), and can be retrieved from NCBI database (<http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=3649>) to facilitate the genomics and proteomics research. Indeed, Rodrigues *et al.* (2009) has published the procedures for sample preparation and 2D gel electrophoresis to study the proteome of papaya leaves. However, the methodology and proteome analysis of papaya seeds has not yet been reported to date.

The objective of this study was to evaluate the quality of protein samples of papaya seeds prepared by two different extraction methods in terms of protein number and resolution of gel profile. Effects of gel staining protocols and pH gradient ranges of Isoelectric focusing (IEF) electrophoresis on protein separation and detection were also assessed. The results may contribute to facilitating the employment of proteomic approaches for the research of seed biology in papaya.

MATERIALS AND METHODS

Plant material and seed germination condition

The papaya seeds utilized in this research were extracted from ripening 'Tainung No.2' papaya fruit. After water floatation to sort out pulp and immature (white) and empty seeds, the arils of heavy and sound seeds sinking to the bottom were removed via scrubbing and washed thoroughly. The papaya seeds were then sterilized in 10% Chlorex® bleach (0.6% sodium hypochlorite) and 1% Tween 20 solution with sonication for 5 minutes followed by thoroughly rinsing with distilled water. The seeds were dried in an air-circulating oven at 36°C for 48 hrs. The dried seeds (seed water content around $7.93 \pm 0.03\%$) were sealed in plastic bags and stored in a desiccator chamber at 10°C until use.

For germination, the papaya seeds were soaked in deionized water at 26°C in dark for 24 hrs before sowing 25 seeds on two layers of Advantec No. 3 filter paper in a 9-cm-diameter Petri dish moistened with 10 mL of deionized water. The seeds were incubated at 26°C without light for four days prior to protein extraction. Because of seed dormancy, there was no seed germination after the five-day incubation. Each protein sample replicate was prepared from 2 g (about 150 seeds) of papaya seeds. All experiments were independently repeated at least three times.

Protein extraction

Two grams of papaya seeds were grounded in liquid nitrogen with a pre-chilled mortar and pestle and trans-

ferred to a tube containing 10 mL of extraction buffer [20 mM Tris-HCl (pH 7.5), 250 mM sucrose, 10 mM ethylenediaminetetraacetic acid (EDTA), 1 mM dithiothreitol (DTT) and 1% (v/v) Triton X-100 and 2% (w/v) of polyvinylpyrrolidone (PVPP) (Sigma, USA). After thoroughly vortexed, the sample was homogenized by Polytron for one minute, and then centrifuged at 13,500 g for 15 minutes at 4°C. The homogenate was filtrated through a layer of Advantec No. 3 filter paper and the filtrate was collected in another centrifuge tube, with addition of PVPP to 2% (w/v) and mixed thoroughly. The sample was incubated on ice for 10 minutes and then centrifuged at 13,500 g for 15 minutes at 4°C. The supernatant was filtrated through filter papers. The protein samples were then further precipitated by TCA/acetone method or extracted by phenol followed by ammonium acetate/methanol mixture.

TCA/acetone precipitation (TCA-A)

The trichloroacetic acid (TCA)/acetone precipitation method was adapted from Yang *et al.* (2007) used for rice seeds with minor modifications and referred as 'TCA-A' in this research. The filtrated protein solution prepared above was mixed with 3x volume of 10% TCA/acetone solution followed by overnight precipitation at -20°C. After 13,500g centrifugation at 4°C for 15 minutes, the supernatant was removed and the precipitant was washed three times with 80% and 100% acetone, respectively. The pellet was dried at room temperature for three minutes.

Phenol extraction, and precipitation with 0.1 M ammonium acetate/methanol mixture (P-AA-M)

The P-AA-M method was performed as described in Rodrigues *et al.* (2009) with modifications. The filtrated protein solution prepared above was mixed with an equal volume of phenol (pH 8.0) for 10 minutes on a vortex, then centrifuged 13,500 g for 15 minutes at 4°C. The upper aqueous layer was removed by using a micropipette and 5x volume of 0.1 M ammonium acetate/methanol mixture was added and precipitated overnight at -20°C. After 13,500 g centrifugation at 4°C for 15 min, the upper aqueous layer was removed without touching the white precipitants, which were then washed twice with 10 mL of 0.1 M ammonium acetate/methanol mixtures followed by twice with 10 mL of 0.1 M ammonium acetate/80% methanol mixture. Finally, the precipitant was washed three times with 80% and 100% acetone separately to remove excess salts and any interfering substances. The precipitants were dried at room temperature for three minutes.

Protein solubilization and quantitation

Dried protein samples prepared from Section 2.1 and 2.2 was dissolved with 3x volume of rehydration buffer containing 7 M urea (Amersham), 2 M thiourea (Merck), 4% (w/v) CHAPS (Sigma), 60 mM DTT (Sigma) (Görg *et al.* 1997) at room temperature. Protein concentration was determined by the Bio-Rad protein assay (Bio-Rad, Hercules, CA, USA) according to the manu-

facturer's instructions. The standard curve was plotted using bovine serum albumin (BSA) (Bio-Rad) as standard.

Sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE)

SDS-PAGE was performed using a 4% (w/v) acrylamide stacking gel and a 12.5% (w/v) separation gel according to Laemmli (1970) on SE 400 Sturdiel Slab Gel Electrophoresis Units (Hoefler) apparatus. Equal amounts (50 µg) of the soluble protein samples were applied to each lane. The electrophoresis was run with running buffer (25 mM Tris-base, 192 mM glycine, 0.1% SDS) at 110 V of constant voltage until the bromophenol blue tracking dye had reached the bottom of the separating gel.

Two-dimensional (2D) gel electrophoresis

The procedure for 2D gel electrophoresis was performed according the user manual (Bio-Rad or Amersham). An aliquot of 800 µg protein sample was mixed with 5% (v/v) immobilized pH gradient (IPG)-buffer (Bio-Rad) and then rehydration buffer was added to a final volume of 500 µL. The sample was loaded onto an IPG strip holder with a pre-cast IPG gel (pH 3–10 or pH 4–7) strip (Bio-Rad) and covered with 2 mL of mineral oil (Bio-Rad). The holder was placed on the Ettan™ IPGphor™ 3 IEF System (Amersham) to be rehydrated at 50 V for 14 h. Isoelectric focusing (IEF) was carried out with 3 h at 150 V, 1 h at 300 V, 1 h at 600 V, 1 h at 1,200 V, 1 h at 1,200–10,000 V (gradient), and 6 h at 10,000 V. The IPG gel was then rinsed with distilled water to remove the cover oil. The IPG gel is used for SDS-PAGE directly or stored at –80°C until use within two weeks.

Prior to SDS-PAGE, IPG gel was equilibrated in 5 mL of SDS equilibration buffer I [6 M urea, 0.375 M Tris-HCl (pH 8.8), 2% SDS, 20% glycerol, 2% (w/v) DTT] for 15 minutes with orbital shaking and subsequently for 15 minutes with shaking in 5 mL of SDS equilibration buffer II [6 M urea, 0.375 M Tris-HCl (pH 8.8), 2% SDS, 20% glycerol, 2.5% (w/v) iodoacetamide]. The separation in the second dimension was realized on the Mini-Trolean (Bio-Rad) or the EttanDALT System (Amersham) with a lab-cast 1.5 mm SDS polyacrylamide gel (12.5%) at a current of 45 mA · gel⁻¹ for 5 hr.

Gel staining and image analysis

2D gel was stained with either zinc-reverse stain method (Fernandez-Patron *et al.*, 1995, 1998) or Coomassie brilliant blue R-250 (CBB) (Neuhoff *et al.*, 1988). For reverse staining, the acrylamide gel was soaked in an equilibration solution [0.2 M imidazole, 0.1% (w/v) SDS] with shaking for 15 minutes and then transferred into a developer solution of 0.3 M zinc sulfate for 30–40 sec with gentle shaking. The stained gel was either scanned immediately or stored in a storage solution of 0.5% (w/v) sodium carbonate.

The stained gel image was scanned at a 400 dpi resolution and the gel image analysis and comparison were

performed with ImageJ 1.43[®] (National Institutes of Health, USA) software for SDS-PAGE gels and with PDQuest™ 8.01 (Bio-Rad) software for 2 DE gels, respectively.

In-gel digestion and protein identification

In-gel digestion was conducted as described by Hellman *et al.* (1995) with modifications. In brief, protein spots of 2-DE gel spots were excised with cut tips and transferred into eppendorf tubes individually. After adding 200 µL destain solution [50% (w/v) acetonitrile, 25 mM ammonium bicarbonate] and incubated at room temperature for 15 minutes, the tube was centrifuged at 10,000 g for one minute. The destain solution was discarded and the destaining step was repeated until the gel was completely colorless. The gel piece was then covered with 200 µL acetonitrile at room temperature for 5 minutes and subsequently centrifuged at 10,000 g for one minute to remove acetonitrile. The sample was allowed to dry for 10 minutes in SpeedVac (Savant). Dehydrated gel particles were incubated in 40 µL of 25 mM ammonium bicarbonate with 100 ng trypsin (Sigma) at 37°C for at least 16 h. After proteolysis, 50 µL elution solution [50% (w/v) acetonitrile, 5% (v/v) trifluoroacetic acid] was added and incubated for 1 h followed by centrifugation at 10,000 g for one minute. The supernatant containing digested peptides was collected. The gel was extracted again with 50 µL elution solution to collect the remaining peptides. The supernatants from the two extractions were pooled together in a new eppendorf and vacuum-dried in SpeedVac for 1 h. The eluent was determined with liquid chromatography-mass spectrometry (LC-MS/MS). Proteomic mass spectrometry analyses were performed by the Core Facilities for Protein Structural Analysis located at the Institute of Biological Chemistry, Academia Sinica. MS/MS raw data were processed and searched against a target protein sequence database using the Mascot (<http://www.matrixscience.com/>). The following search criteria were used: (1) the database was set to be Swiss-Port or NCBItr; (2) taxonomy was set to be green plants; (3) enzyme setting was trypsin; (4) MS/MS tolerance was set to be 0.25 Da; (5) data format setting was pkl; (6) instrument was set to be ESI-QUAD. Protein hits with Mascot score >50 and P value <0.05 as well as matched molecular weights resolved on the gel were then identified as significant proteins.

RESULTS AND DISCUSSION

Effects of extraction methods on papaya seed proteome analysis

It has been reported that the papaya seeds are rich in phenolic compounds and *p*-hydroxybenzoic acid being the major one (Chow and Lin, 1991; Kothari and Seshadri, 2010). To remove these phenolic compounds interfering the efficiency of protein extraction and resolution of 2D analysis, we have employed both TCA-A and P-AA-M for isolating proteins from papaya seeds, independently. The extracted proteins were analysed

by 1D SDS-PAGE. As shown in Fig. 1, the amounts of proteins extracted by P-AA-M method were significantly higher and resulted in at least five more protein bands on SDS-PAGE than those extracted by TCA-A method. These extra bands were resolved with molecular weights at 75, 37, 30, 25, and 20 kDa.

The same protein samples were also analysed by 2D gel electrophoresis with first dimension in pH 3–10 IPG gel and second dimension in 12.5% SDS-PAGE. As shown in Fig. 2, the proteins isolated by P-AA-M method (Fig. 2B) revealed much better separation between protein blots and higher resolution than those precipitated by TCA-A method (Fig. 2A), in which serious streaking and smearing were evident on the 2D gel. Consequently, P-AA-M method was able to resolve 717 ± 32 protein spots as compared with only 124 ± 27 protein spots were detected from TCA-A method. The result clearly indicated that phenol extraction combined with ammonium acetate/methanol precipitation was a better method than typical TCA/acetone method for papaya seed proteome analysis.

While proteomics plays important roles in research of plant biology and agriculture, not all the organs or cell types were actively investigated at proteome scale.

Fig. 1. SDS-PAGE analysis of papaya seed proteins prepared by 10% trichloroacetic acid (TCA)/acetone precipitation (TCA-A) (lanes 1 and 2) or phenol extraction followed by 0.1M ammonium/methanol precipitation (P-AA-M) (lane 3 and 4). The arrows indicate the different protein bands between the two extraction methods. Marked band A and band B are distinct protein bands prepared by P-AA-M. Fifty microgram of protein was loaded per lane. M: molecular weight marker.

In view of the importance of proteomics research in understanding the molecular mechanisms and pathways underlying the papaya seed germination and responses to various environmental signals, we have now established the methodology for protein extraction and 2D gel electrophoresis of papaya seeds. This provides an important platform for further in-depth seed proteomics analysis of papaya or even other plants. In particular, this platform shall be most useful for proteomics study of woody plants or tissues with high contents of secondary metabolites or phenolic compounds,

Fig. 2. Two-dimensional profiles of papaya seed proteins extracted by using two different extraction methods and visualized by different staining processes. (A) Proteins were extracted by 10% trichloroacetic acid / acetone precipitation (TCA-A) and the gel was stained with zinc reverse-stain. (B) Proteins were extracted by phenol extraction followed by 0.1 M ammonium/methanol precipitation (P-AA-M) and the gel was stained with zinc reverse-stain. (C) Proteins were extracted by P-AA-M method and the gel was stained with Coomassie Brilliant Blue R-250. Molecular weight markers were indicated on the left.

which are the interfering substances that may be not efficiently removed by the typical protein extraction protocol (Wang *et al.*, 2006).

Most popular protein extraction/precipitation method used for plant materials is the utilization of 10% TCA/acetone for protein precipitation prior to 2D gel electrophoresis of plant tissues. This method has been widely used in a variety of plant species and tissues including *Arabidopsis* plantlets (Santoni *et al.*, 1994), rice seeds (Yang *et al.*, 2007), and papaya leaves (Rodrigues *et al.*, 2009). This method has been demonstrated to efficiently remove non-protein macromolecular polymers such as water-soluble cell wall polysaccharides and polyphenols and thereby prevent their interferences on 2D gel electrophoresis (Saravanan and Rose, 2004). However, TCA/acetone precipitation is not sufficient to remove the interferences of papaya seeds and thereby results in significant horizontal and vertical streaking on the 2D gel (Fig. 2A). This poor resolution may be caused by the existence of phenolic compounds that are rich in papaya seeds (Chow and Lin, 1991; Kothari and Seshadri, 2010) but could not be efficiently removed by TCA/acetone precipitation. Even 2% PVPP was added as the phenol compound adsorbent in the protein extraction buffer we used, no further improvements in the number and resolution of protein spots on 2D gel electrophoresis could be detected. This result may be caused by the incapability of TCA-A precipitation step in avoiding the co-precipitation of the interfered compositions of papaya seed from its proteins. In contrast, when the buffer-extracted proteins were extracted again by phenol followed by 0.1 M ammonium acetone/methanol mixtures to perform precipitation, the interferences were significantly suppressed. Moreover, since proteins were partitioned into the phenol phase, phenol extraction step could remove most of the water-soluble contaminants and salts. Wang *et al.* (2006) incorporated phenol extraction process to their extraction protocol and gained protein samples with lower interferences for 2D gel electrophoresis from olive (*Olea europaea* L.) leaves and fruit, as well as apple (*Malus domestica* Borkh.) fruit. Furthermore, Zheng *et al.* (2007) pointed out that protein samples of strawberry (*Fragaria x ananassa* Duchesne) fruit prepared by method including phenol extraction step showed more distinguishable protein spots and better resolution on 2D gel profile than the sample without phenol extraction. The success in the use of phenol for protein extraction from cotton (*Gossypium hirsutum* L.) seed that is rich in polysaccharides, polyphenols, nucleic acids, and secondary metabolites was also demonstrated for reduced background interferences and increased protein spots on 2D gel (Xie *et al.*, 2009). Additionally, the chemical reagents of the P-AA-M method we used did not include any organic acid, thus the pH of sample was less disturbed during extraction and precipitation processes (Wang *et al.*, 2006) and, as a consequence, minimized the frequency of washing steps as well as prevented interference of IEF electrophoresis.

Comparison of zinc-reverse stain and CBB stain for protein separation and detection on 2D profile analysis

Two staining methods, zinc-reverse stain and CBB stains, were used to reveal the protein profiles of papaya seeds. We observed seven folds or more of protein spots on zinc-reverse stained gel (Fig. 2B) than CBB-stained gel (Fig. 2C). This result is consistent with higher sensitivity of zinc reverse stains for detecting 1 ng of BSA, as compared with CBB at sensitivity at 100 ng band⁻¹ (Patton, 2002). Importantly, zinc reverse stains also revealed linear range of sensitivity from 1 to 100 ng of BSA, suggesting that this is a sensitive and quantifiable staining method suitable for 2D gel analysis.

CBB with minimal sensitivity 100 ng band⁻¹ (Patton, 2002) is routinely used in protein gel staining because of its high reproducibility, simple operation, and wide compatibility for most of proteins. However, proteins accumulated in a low abundant level cannot be visualized via CBB stain, thus silver stain or fluorescent dye SYBRO Ruby with a higher sensitivity are used as a substitute technique (Miller *et al.*, 2006). Despite its high sensitivity (1 ng band⁻¹) to protein staining, the reproducibility, narrow reliable linear range, and spot intensity highly influenced by the timing of staining procedure are problematic and therefore easily render in an incorrect estimation for silver stained 2D gel when the profile is analyzed (Miller *et al.*, 2006). On the other hand, the sensitivity of SYBRO Ruby is comparable to silver staining. The former displays better reproducibility and reliable linear range than the later. However, the fluorescent dye is expensive and images have to be analysed by the specialized fluorescent machine such as Typhoon Imager (Amersham), thus caused many restrictions. Zinc-reverse stain used in the present study showed comparable in both the time required for staining process and detection sensitivity as compared with other two staining methods. The intensity of protein spots on reverse-stained gel was almost independent from the incubation time of staining. In addition, reverse-stained gel was less likely contaminated by chemicals during the procedure and was feasible for destaining and restaining, which is an advantage that CBB stain and silver stain lack. According to the results we obtained, the profile of reverse-stained 2D gel showed more distinguishable protein spots than those of CBB-stained one (Fig. 2B and C). The fact that proteins invisible on CBB-stained gels were detected on the zinc-reverse stained gels proves that zinc-reverse staining enhanced sensitivity of protein detection. Moreover, if gel staining was overdone and was not in an optimal situation, zinc-reverse stained gels could be destained and restained again to optimize the staining intensity or restained with CBB without any obvious interference. In considering its comparable sensitivity and several advantages, zinc-reverse stain may be considered as the most convenient, speedy, and affordable staining method for proteomics study.

Effect of the pH gradient range of IPG gel strip on the resolution of 2D profile

The pH 3–10 IPG gel was initially used for 2D gel electrophoresis (Fig. 2B), which exhibited that the majority of proteins are resolved in the pH range between 4 and 8. To further improve the resolution of 2D gel, the protein samples extracted by P-AA-M method were separated in the pH 4–7 IPG gel prior to SDS-PAGE. The protein profile significantly increased the resolution of protein spots in this pH range (Fig. 3) when compared to those in pH 3–10 gel (Fig. 2B).

High quality of protein samples is absolutely essential for good result of IEF. The presence of interfering substances, such as nucleic acids and salts, would cause proteins focusing incompletely during electrophoresis, resulting in blurry and distorted 2D gel images (Saravanan and Rose, 2004). Protein samples extracted by P-AA-M method enhanced the resolution of the 2D gel profile of papaya seeds protein when compared to those by TCA-A protocol (Fig. 2A and B). The protein spots of papaya seeds on 2D gel were found mainly in the region between pH 6 and pH 7.5, when a pH 3–10 IPG gel strip was used for IEF electrophoresis (Fig. 2B). We, therefore, further successfully improved the number and resolution of protein spots on 2D profile by separation in IPG gel at pH 4–7 (Fig. 3). Our data also suggested that the pI values of the water-soluble proteins of papaya seeds were barely resolved in extreme pH ranges (Fig. 2 B). The lower representatives of acidic and alkaline proteins could be due to the technical issue as proteins with extreme acidic or basic pI or hydrophobic characteristics, such as the membrane protein and ribosomal protein are poorly extracted or resolved in IEF. Alternatively, the proteins from papaya seeds may contain less highly acidic or basic proteins. Future

work to use IPG gel with narrow range in acidic or basic pH may reveal more acidic and basic proteins.

The efficacy of protein separation by IEF was also affected by the amount of proteins loaded. When proteins was over loaded, it spends more time for each protein to move to its isoelectric point; smearing or aggregation of protein samples at a certain region on the 2D gel was observed especially in case of electrophoresis process was not long enough. We found that the optimal concentration of protein sample for IEF electrophoresis was $1.6 \mu\text{g } \mu\text{L}^{-1}$ (i.e. $800 \mu\text{g}$ proteins in $500 \mu\text{L}$ IPG buffer solution).

Protein identification by LC-MS/MS

To further compare the differences of extraction efficacy between TCA/A and P-AA-M methods and to examine the compatibility of the P-AA-M method with LC-MS/MS analysis, two outstanding 30- (Band A) and 20-kDa (Band B) protein bands on the SDS-PAGE analysis (Fig. 1) were chosen for protein identifications by LC-MS/MS. A total of eight and five of proteins were identified from 30-kDa (Band A) and 20-kDa (Band B) bands, respectively, as significant hits (Table 1). For the 2D gel analysis, five protein spots were selected for protein identification (Table 1).

Although 15 of proteins isolated from papaya seeds were identified from selected protein bands and spots, only two proteins, namely small GTP-binding protein (Table 1) and 17.5 kDa class I heat shock protein (Table 1), matched the papaya protein database whereas other proteins were identified by matching with the orthologous proteins encoded in other green plants (Table 1). Similar observation was also reported in the proteomics analysis of papaya leaves, in which only a few proteins were identified from papaya protein database (Rodrigues *et al.* 2009). These results indicated that papaya genome has been completed, but the protein database is barely annotated for protein identifications by MS/MS ions.

In conclusion, we have clearly demonstrated that protein samples of papaya seeds, a high phenolics-contained plant material, prepared by the P-AA-M method included more protein spots and less contaminants than traditional TCA/A protocol via SDS-PAGE and 2D gel analysis. Smearing and streaking problems occurred on the 2D gel profile of TCA/A-extracted samples were markedly suppressed when P-AA-M-prepared samples utilized. The protein number and resolution on 2D gel profile were further improved by incorporation of pH 4–10 IPG gel strip for IEF electrophoresis and zinc-reverse stain technique. The protein samples of papaya seeds isolated from the protocol we established were compatible with LC-MS/MS analysis. The high-resolution 2D map with identifications of several proteins from papaya seeds shall serve as a platform and reference for future research in seed biology of papaya or other plants.

Fig. 3. Two-dimensional gel profile of papaya seed proteins prepared by phenol extraction followed by 0.1 M ammonium/methanol precipitation (P-AA-M) after first-dimension separation using a pH 4–7 immobilized pH gradient (IPG) gel strip and visualized by zinc reverse-staining. Spot numbers corresponding with the numbers in Table 2 are labeled with white arrows pointing to each protein spot.

REFERENCES

- Andreoli, C. and A. A. Khan 1993 Improving papaya seedling emergence by matricconditioning and gibberellins treatment.

Table 1. Identifications of the distinct protein bands (bands A and B seen in Fig. 1) prepared by phenol extraction followed by 0.1 M ammonium/methanol precipitation (P-AA-M), those prepared by 10% trichloroacetic acid / acetone precipitation (TCA-A) in SDS-PAGE analysis and protein spots from 2D gel analysis. The proteins were identified by LC-MS/MS

Putative protein	Accession No.	Organism	Calculated MW (kDa)/pI	Peptide sequence hit	Sequence coverage (%)	Protein scores
SDS-PAGE						
<i>Band A</i>						
Malate dehydrogenase	P19446	<i>Citrullus lanatus</i>	37.6/8.67	RGFLGQQQLEAALTGMDLIIVPAGVPRK; RAIVNLISNPVNSTVPAAAEVFKK; RAIVNLISNPVNSTVPAAAEVFKKA; KRLLGVTMLDVVRA; RLLGVTMLDVVRA	17	199
Hypothetical protein	XP_002273346	<i>Vitis vinifera</i>	33.1/5.50	KTVLVDNEDFLKE; KITRQPGPIPLNTKI; KGNAYAQVAISTDDVYKS	14	140
Rubber elongation factor protein	P15252	<i>Heveabra siliensis</i>	14.7/5.04	KSGPLQPGVDIIEGPVK; KNVAVPLYNRF; KDASIQVVSIRA	26	96
Putative uncharacterized protein	ACJ85383	<i>Medicago truncatula</i>	31.5/7.06	KVPGGLAVVIVGNRQ; RSNIVGLPVSLLLKA; KVAGWITPVPGGVGPMTVTMLLN	16	66
Glucose and ribitol dehydrogenase homolog 1	AAG51119	<i>Arabidopsis thaliana</i>	31.2/5.92	RVVDEVVNAFGRI; KRVVDEVVNAFGRI; KRVVDEVVNAFGRI	4	59
Cytosolic cysteine synthase	BAB20861	<i>Solanum tuberosum</i>	34.3/5.93	RAFGAELVLTDPAGK; KLIITMPSSMSLERR	8	54
Putative uncharacterized protein	XP_002894087	<i>Arabidopsis lyrata</i>	34.3/8.77	MAITKERR; RTPEEILRI	4	47
Putative NADPH dependent mannose 6-phosphate reductase	NP_179721	<i>Arabidopsis thaliana</i>	35.0/6.16	KTVAQVVLRW; RDLILNAIKI	5	41
<i>Band B</i>						
RAB1C	CAA98160	<i>Lotus japonicus</i>	22.6/5.30	KLLIGDSGVGKS; KNATNVEQAFMAMAAEIKN; KNATNVEQAFMAMAAEIKN; RFADDSYLDYSISTIGVDFKI	23	152
Ras-related protein RIC2	BAA02904	<i>Oryza sativa</i>	24.0/5.37	RAITSAYYRG; KSTIGVEFATRS; RGAVGALLVYDVTRH	14	68
Predicted protein	XP_002299359	<i>Populus trichocarpa</i>	18.7/5.07	RNFNIIQHQLGKL; RNFNIIQHQLGKL	6	63
Small GTP-binding protein	BAA88497	<i>Carica papaya</i>	23.3/8.43	KTVAQVVLRW; RDLILNAIKI	17	51
2D-gel						
<i>Spot 4</i>						
17.5 kDa class I heat shock protein	AAG51119	<i>Carica papaya</i>	17.47/5.31	DFPFPPSSISR; VELEDDRVLQISGER; ASMEN GVLTVTVPK	25	191
<i>Spot 5</i>						
Late embryogenesis abundant protein	AAB01563	<i>Picea glauca</i>	22.47/6.22	AELSQVKR	4	59

Hortscience, **28**: 708–709

Bhattacharya, J. and S. S. Khuspe 2001 *In vitro* and *in vivo* germination of papaya (*Carica papaya* L.) seeds. *Sci. Hort.*, **91**: 39–49

Bradford, M. M. 1976 A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, **72**: 248–254

Carpentier, S. C., E. Witters, K. Laukens, P. Deckers, R. Swennen and B. Panis 2005 Preparation of protein extracts from recalcitrant plant tissues: An evaluation of different methods for two-dimensional gel electrophoresis analysis. *Proteomics*, **5**:

2497–2507

Chacko, E. K. and R. N. Singh 1966 The effect of gibberellic acid on the germination of papaya seeds and subsequent seedling growth. *Trop. Agri. Trin.*, **43**: 341–346

Chow, Y. J. and C. H. Lin 1991 p-Hydroxybenzoic acid as the major phenolic germination inhibitor of papaya seed. *Seed Sci. & Technol.*, **19**: 167–174

Fernandez-Patron, C., M. Calero, P. R. Collazo, J. R. Garcia, J. Madrazo, A. Musacchio, F. Soriano, R. Estrada, R. Frank, L. R. Castellanoserra and E. Mendez 1995 Protein reverse staining: high-efficiency microanalysis of unmodified proteins

- detected on electrophoresis gels. *Anal. Biochem.*, **224**:203–211
- Görg, A. 2000 Advances in 2D gel techniques. *Trends in Biotechnol.*, **18**: supplement 13–16
- Görg, A., C. Obermaier, G. Boguth, A. Csordas, J. J. Diaz and J. J. Madjar 1997 Very alkaline immobilized pH gradients for two-dimensional electrophoresis of ribosomal and nuclear proteins. *Electrophoresis*, **18**: 328–337
- Hellman, U., C. Wernstedt, J. Genezand and C. H. Heldin 1995 Improvement of an “In-Gel” digestion procedure for the micropreparation of internal protein fragments for amino acid sequencing. *Anal. Biochem.*, **224**: 451–455
- Hurkman, W. J. and C. K. Tanaka 1986 Solubilization of plant membrane proteins for analysis by two-dimensional gel electrophoresis. *Plant Physiol.*, **81**: 802–806
- Kothari, V. and S. Seshadri 2010 Antioxidant activity of seed extracts of *Annona squamosa* and *Carica papaya*. *Nutr. & Food Sci.*, **40**: 403–408
- Laemmli, U. K. 1970 Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**: 680–685
- Miernyk, J. A. and M. Hajdich 2011 Seed proteomics. *J. Proteomics*, **74**: 389–400
- Mijnsbrugge, K. V., H. Meyermans, M. Van Montagu, G. Bauw and W. Boerjan 2000 Wood formation in poplar: identification, characterization, and seasonal variation of xylem proteins. *Planta*, **210**: 589–598
- Ming, R., S. Hou, Y. Feng, Q. Yu, A. Dionne-Laporte, J. H. Saw, P. Senin, W. Wang, B. V. Ly, K. L. Lewis, S. L. Salzberg, L. Feng, M. R. Jones, R. L. Skelton, J. E. Murray, C. Chen, W. Qian, J. Shen, P. Du, M. Eustice, E. Tong, H. Tang, E. Lyons, R. E. Paull, T. P. Michael, K. Wall, D. W. Rice, H. Albert, M. L. Wang, Y. J. Zhu, M. Schatz, N. Nagarajan, R. A. Acob, P. Guan, A. Blas, C. M. Wai, C. M. Ackerman, Y. Ren, C. Liu, J. Wang, J. Wang, J. K. Na, E. V. Shakirov, B. Haas, J. Thimmapuram, D. Nelson, X. Wang, J. E. Bowers, A. R. Gschwend, A. L. Delcher, R. Singh, J. Y. Suzuki, S. Tripathi, K. Neupane, H. Wei, B. Irikura, M. Paidi, N. Jiang, W. Zhang, G. Presting, A. Windsor, R. Navajas-Perez, M. J. Torres, F. A. Feltus, B. Porter, Y. Li, A. M. Burroughs, M. C. Luo, L. Liu, D. A. Christopher, S. M. Mount, P. H. Moore, T. Sugimura, J. Jiang, M. A. Schuler, V. Friedman, T. Mitchell-Olds, D. E. Shippen, C. W. dePamphilis, J. D. Palmer, M. Freeling, A. H. Paterson, D. Gonsalves, L. Wang and M. Alam 2008 The draft genome of the transgenic tropical fruit tree papaya (*Carica papaya* Linnaeus). *Nature*, **452**: 991–997
- Neuhoff, V., N. Arold, D. Taube and W. Ehrhardt 1988 Improved staining of proteins in polyacrylamide gels including isoelectric focusing gels with clear background at nanogram sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis*, **9**: 255–262
- Patton, W. F. 2002 Detection technologies in proteome analysis. *J. Chromatogr. B Analyt. Technol. Biomed. Life Sci.*, **771**: 3–31
- Paull, R. E. and O. Duarte 2011 Papaya. In “Tropical Fruits”, Vol.1 (2nd ed.), eds. by R. E. Paull, and O. Duarte, CABI, Oxfordshire, pp. 291–326
- Rodrigues, S. P., J. A. Ventura, R. B. Zingali and P. M. B. Fernandes 2009 Evaluation of sample preparation methods for the analysis of papaya leaf proteins through two-dimensional gel electrophoresis. *Phytochem. Anal.*, **20**: 456–464
- Salomao, A. N. and R. C. Mundim 2000 Germination of papaya seed in response to desiccation, exposure to subzero temperatures, and gibberellic acid. *HortScience*, **35**: 904–906
- Saravanan, R. S. and J. K. C. Rose 2004 A critical evaluation of sample extraction techniques for enhanced proteomic analysis of recalcitrant plant tissues. *Proteomics*, **4**: 2522–2532
- Singh, S. P. and D. V. Sudhakar 2011 Papaya (*Carica papaya* L.). In “Postharvest Biology and Technology of Tropical and Subtropical Fruits”, Vol. 4: Mangosteen to White Sapote, ed. by E. M. Yahia, Woodhead Publishing, Oxford, pp. 86–124
- Wang, W., M. Scali, R. Vignani, A. Spadafora, E. Sensi, S. Mazzuca and M. Cresti 2003 Protein extraction for two-dimensional electrophoresis from olive leaf, a plant tissue containing high levels of interfering compounds. *Electrophoresis*, **24**: 2369–2375
- Wang, W., R. Vignani, M. Scali and M. Cresti 2006 A universal and rapid protocol for protein extraction from recalcitrant plant tissues for proteomic analysis. *Electrophoresis*, **27**: 2782–2786
- Wood, C. B., H. W. Pritchard and D. Amritphale 2000 Desiccation-induced dormancy in papaya (*Carica papaya* L.) seeds is alleviated by heat shock. *Seed Sci. Res.*, **10**: 135–145
- Xie, C., D. Wang and X. Yang 2009 Protein extraction methods compatible with proteomic analysis for cotton seedling. *Crop Sci.*, **49**: 395–402
- Yang, P., X. Li, X. Wang, H. Chen, F. Chen and S. Shen 2007 Proteomics analysis of rice (*Oryza sativa*) seeds during germination. *Proteomics*, **7**: 3358–3368
- Zheng, Q., J. Song, K. Doncaster, E. Rowland and D. M. Byers 2007 Qualitative and quantitative evaluation of protein extraction protocols for apple and strawberry fruit suitable for two-dimensional electrophoresis and mass spectrometry analysis. *J. Agric. Food Chem.*, **55**: 1663–1673