The Labyrinth model and products of two Cantor sets

Takahashi, Yuki Advanced Institute for Materials Research, Tohoku University

https://doi.org/10.15017/2924850

出版情報:応用力学研究所研究集会報告. 2019A0-S2, pp.37-38, 2020-03. Research Institute for Applied Mechanics, Kyushu University バージョン: 権利関係: 応用力学研究所研究集会報告 No.2019AO-S2 「非線形波動研究の多様性」(研究代表者 永井 敦)

Reports of RIAM Symposium No.2019AO-S2

Diversity in the research of nonlinear waves

Proceedings of a symposium held at Chikushi Campus, Kyushu University, Kasuga, Fukuoka, Japan, October 31 - November 2, 2019

Article No. 07 (pp. 37 - 38)

The Labyrinth model and products of two Cantor sets

高橋 悠樹(Takahashi Yuki)



Research Institute for Applied Mechanics Kyushu University March, 2020

The Labyrinth model and products of two Cantor sets

YUKI TAKAHASHI

Advanced Institute for Materials Research, Tohoku University

ABSTRACT. We consider the Labyrinth model, which is a separable two-dimensional quasicrystal model. The spectrum of the Labyrinth model is given by products of two Cantor sets. We will show that the spectrum of the Labyrinth model is an interval for coupling constants sufficiently close to 1.

1. Products of two Cantor sets

For any gap U of a Cantor set, we denote the right (resp. left) endpoint of U by U^R (resp. U^L). If gaps U_1, U_2 satisfy $U_1^R < U_2^L$, we denote $U_1 < U_2$.

DEFINITION 1.1. Let K be a Cantor set. We define the *thickness* of K by

$$\inf_{U_1 < U_2} \max \left\{ \frac{U_2^L - U_1^R}{|U_1|}, \frac{U_2^L - U_1^R}{|U_2|} \right\},\,$$

where the infimum is taken for all pairs of gaps of K, with at least one of them being a finite gap. We denote this value by $\tau(K)$.

In [1], the author considered products of two Cantor sets and obtained the optimal estimate in terms of thickness that $K \cdot L$ is an interval. We call K a 0-Cantor set if $K_+, K_- \neq \phi$, inf $K_+ = 0$ and inf $K_- = 0$.

THEOREM 1.1 (Theorem 1.4 in [1]). Let K, L be 0-Cantor sets. Then, if

$$2(\tau(K) + 1)(\tau(L) + 1) \leq (\tau(K)\tau(L) - 1)^{2},$$

 $K \cdot L$ is an interval. In particular, if

$$\tau(K) = \tau(L) \ge 1 + \sqrt{2},$$

then $K \cdot L$ is an interval. Furthermore, for any M, N > 0 with

$$2(M+1)(N+1) > (MN-1)^2,$$

there exist 0-Cantor sets K, L such that $\tau(K) = M$, $\tau(L) = N$, and $K \cdot L$ is a disjoint union of two intervals.

2. Spectrum of the Labyrinth model

Let $\lambda>1$ and define

$$\omega_{\lambda}(n) = \begin{cases} \lambda & \text{if } n\alpha \mod 1 \in [1-\alpha, 1) \\ 1 & \text{if } n\alpha \mod 1 \in [0, 1-\alpha), \end{cases}$$

where $\alpha = \frac{-1+\sqrt{5}}{2}$ is the inverse of the golden mean. The *Labyrinth model* is given by the following self-adjoint operator that acts on $\ell^2(\mathbb{Z}^2)$:

$$(H_{\lambda_{1},\lambda_{2}}\psi)(m,n) = \omega_{\lambda_{1}}(m+1)\omega_{\lambda_{2}}(n+1)\psi(m+1,n+1) + \omega_{\lambda_{1}}(m+1)\omega_{\lambda_{2}}(n)\psi(m+1,n-1) + \omega_{\lambda_{1}}(m)\omega_{\lambda_{2}}(n+1)\psi(m-1,n+1) + \omega_{\lambda_{1}}(m)\omega_{\lambda_{2}}(n)\psi(m-1,n-1).$$

In [2], the author proved the following:

THEOREM 2.1 (Theorem 1.1 in [2]). The spectrum $\sigma(H_{\lambda_1,\lambda_2})$ is given by products of two Cantor sets, and is an interval for sufficiently large $\lambda_1, \lambda_2 > 1$.

References

- [1] Y. Takahashi, Products of two Cantor sets, Nonlinearity 30 (2017), 2114–2137.
- Y. Takahashi, Quantum and spectral properties of the Labyrinth model, J. Math. Phys. 57, 2016.