

Type D personality and periodontal disease in university students: A prospective cohort study

Mizutani, Shinsuke

Section of Geriatric Dentistry, Fukuoka Dental College

Ekuni, Daisuke

Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University

Takeuchi, Mayu

Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University

Azuma, Tetsuji

Graduate School of Medicine, Dentistry and Pharmaceutical Sciences, Okayama University

他

<https://hdl.handle.net/2324/3053979>

出版情報 : Journal of Health Psychology. 23 (5), pp.754-762, 2018-04-01. Sage Publications
バージョン :
権利関係 : © 2018 by the author(s).

**Type D personality and periodontal disease in university students: A prospective
cohort study**

ABSTRACT

The purpose of this prospective cohort study was to investigate whether Type D personality was related to periodontal disease in Japanese university students. Among students (n = 600) who were interested in receiving oral health examinations, logistic regression analysis showed that the risk of developing periodontal disease during a 3-year period was associated with BMI \geq 25 (OR: 2.543; 95% CI: 1.297-4.989; p = .007) and Type D personality (OR: 1.473; 95% CI: 1.027-2.111; p = .035). In this short-term prospective cohort study, a significant association between periodontal disease and Type D personality was observed in Japanese university students.

Keywords: Health behavior; Cohort; Periodontal disease; Type D personality; University students.

INTRODUCTION

Periodontal disease is one of the most widespread chronic diseases, and is predominantly a bacterial infection involving the dental biofilm or dental plaque (Genco and Borgnakke, 2013). Numerous studies have investigated the risk of periodontal diseases, and have provided evidence supporting the important role of risk factors of periodontal disease. The prevalent risk factors are being of male sex (Eke et al., 2012), smoking (Tonetti, 1998), diabetes (Simpson et al., 2010), obesity (Suvan et al., 2011), osteoporosis (Martinez-Maestre et al., 2010), psychological stress/psychiatric symptoms (Peruzzo et al., 2007), and genetic factors (Meng et al., 2011). In addition, behavior such as coping stress (Wimmer et al., 2005) and oral health behavior (usage of dental floss or regular dental check-ups) (Mizutani et al., 2012) are related to periodontal disease. However, few studies have investigated the relationship between personality and periodontal disease.

Type D personality was established to describe individuals who exhibit both negative affectivity (NA) and social inhibition (SI) (Denollet, 2005). Individuals with high NA tend to experience undesired emotions such as anger, distress, hostility and depression. Individuals with high SI tend to experience social difficulties because they hide their emotions in social interactions due to fear of rejection or disapproval from

others. In recent years, Type D personality has been linked to a series of illnesses. Type D personality is associated with negative outcomes in various diseases such as type 2 diabetes, irritable bowel syndrome, heart diseases and coronary heart disease (Ginting et al., 2016; Nefs et al., 2012; Schiffer et al., 2010; Yildirim et al., 2013). The relationship between Type D personality and various psychological problems, including depression, anxiety, and posttraumatic stress disorder, has been investigated (Howard and Hughes, 2012; Kupper et al., 2007). Moreover, Type D personality is associated with work-related problems such as job stress and burnout (Mommersteeg et al., 2012; Oginska-Bulik, 2006), and subjective health (Williams et al., 2015).

There are some studies about Type D personality and health behavior in healthy young adults. Individuals with Type D personality exhibited an approximately four times increased risk of having sleep disturbances, and type D personality was associated with fewer sleeping hours (Condén et al., 2013). Type D individuals engaged in fewer health behaviors like eating a balanced diet and getting regular exercise, and were less likely to get regular medical check-ups than non-Type D individuals (Williams et al., 2008). Type D was also associated with smoking, lack of physical activity, low levels of self-efficacy, maladaptive coping, and less healthy food intake (Booth and Williams, 2015; Gilmour and Williams, 2012; Wiencierz and Williams, 2016). Thus, Type D

personality affects health behavior in even the young generation. Although periodontal health is critically dependent upon the oral health behaviors of the patient (Newton et al., 2015), no studies have examined the relationship between Type D personality and oral health behaviors.

It is important to evaluate the relationship between Type D personality and periodontal disease in young adults, as progression of periodontal disease, which is one of the major causes of tooth loss, was observed in university students (Mizutani et al., 2015). We hypothesized that Type D personality may predict onset of periodontal disease in young adults. The aim of the present prospective cohort study was to explain the relationship between Type D personality and periodontal disease in university students.

METHODS

Study population

Of the first-year students ($n = 2,297$) who underwent both general and oral health examinations (pre-university) at the Health Service Center of Okayama University in April 2012, 692 students volunteered to receive a 3-year follow-up examination before graduation in April 2015 (follow-up rate; 30.1%). After excluding participants with incomplete data in their questionnaires ($n = 44$) and with periodontal disease at baseline ($n = 48$), data from 600 systemically healthy students (282 males, 318 females), aged 21.2 ± 0.6 years (range, 21-28 years) at follow-up were analyzed (Figure 1). The study was approved by the Ethics Committee of Okayama University. Verbal consent was obtained from all participants.

Type D personality

Type D personality was assessed using the 14-item Type D Personality Scale (DS14) (Denollet, 2005). The DS14 comprises two subscales: negative affectivity (NA; 7 items; range 0–28) and social inhibition (SI; 7 items; range 0–28). Participants were asked to rate their responses on a 5-point Likert-type scale (from 0=false to 4=true). We defined participants with Type D personality as those with scores of greater than 10 on both

subscales. (i.e., NA & SI) (Denollet et al., 1996; Emons et al., 2007). The Cronbach's alpha was 0.88 for the NA and 0.86 for the SI, respectively and test-retest reliability was 0.72 and 0.82 for the NA and SI, respectively in the original DS14 (Denollet, 2005). In the current study, the Cronbach's alpha observed was 0.891 for the NA subscale, and 0.874 for the SI subscale. Moreover, the test-retest reliability was 0.827 for the NA subscale, and 0.824 for the SI subscale in the preliminary study targeted at other university students (n = 38).

Oral examination

Three dentists (D.E., S.M., and T.A.) examined the oral health status of the participants. The number of teeth present was recorded. Periodontal condition was assessed using the Community Periodontal Index (CPI) (Furuta et al., 2011). Ten teeth were selected for periodontal examination: two molars in each posterior sextant, and the upper right and lower left central incisors. Measurements were made using a CPI probe (YDM, Tokyo, Japan) at six sites (mesio-buccal, mid-buccal, disto-buccal, disto-lingual, mid-lingual and mesio-lingual) per tooth. The percentage of teeth exhibiting bleeding on probing at least one site (%BOP: number of BOP-positive teeth per total number of teeth) was calculated. Since BOP has been shown to be an earlier and more sensitive indicator of

inflammation than visual signs of inflammation (redness and swelling) (Greenstein, 1984). Thus, we also assessed %BOP as an indicator of periodontal disease in this study (Furuta et al., 2011). Periodontal disease was defined using following criteria: presence of probing pocket depth (PPD) \geq 4 mm (CPI score = 3 or 4) and BOP \geq 30% (active periodontal pocket) (Kawabata et al., 2008). Good intra- and inter-examiner agreement was achieved for repeated PPD measurements in the 10 teeth used for CPI with Kappa statistic $>$ 0.8.

Body Mass Index (BMI)

In the general health examination, the height and body weight of participants were measured by the University's public health nurses using the Tanita body fat analyzer (Model No. BF-220; Tanita, Tokyo, Japan). Obesity was assessed on the basis of BMI (kg/m^2) at 4 levels: $<$ 22.0, 22.0 to 24.9, 25.0 to 29.9, and \geq 30.0 (Ekuni et al., 2014; Morita et al., 2011).

Questionnaire

At baseline, participants filled out a self-administered questionnaire. In addition to gender, age (continuous), general health condition and smoking status (current/never),

the questionnaire also asked about the following items: (i) daily frequency of tooth brushing (\leq One time/ \geq Two times); (ii) use of dental floss (yes/no); and (iii) visits to dental clinics for regular check-up (yes/no). The questionnaire also evaluated the change of oral health behaviors at the 3-year follow-up (Mizutani et al., 2015).

Statistical analysis

The Cronbach's alpha coefficient was used to examine the consistency of scales. Differences between the baseline and follow-up were evaluated with paired t-test or McNemar test, as appropriate. When the participants have periodontal disease after 3 years, they were defined as the subjects with onset of periodontal disease. First, we examined the association between Type D personality and oral health behaviors at baseline and after 3 years using chi-square test. Second, unpaired t-test and chi-square test were used to determine whether there were any significant differences between the participants with and without the onset of periodontal disease at re-examination. We used Fisher's exact test when expected cells were found to be less than 5. Multivariate logistic regression was used to calculate ORs and 95% confidence intervals (95% CI) after controlling simultaneously for potential confounders. The onset of periodontal disease was used as the dependent variable. Independent variables were selected when

the P value was $< .20$ for the chi-square test or unpaired t-test in each variable. Since there were only two students who were $\text{BMI} \geq 30$, we combined two categories ($\text{BMI} \geq 30$ and 25.0 to 29.9) into one ($\text{BMI} \geq 25.0$). Results of logistic regression analyses were considered statistically significant when $p < .05$. We evaluated the predictive performance of the model by assessing its discrimination (ability to classify correctly), and assessed the model fit using the Hosmer–Lemeshow goodness-of-fit test for logistic regression. We conducted all analyses using SPSS version 22.0 (IBM, Tokyo, Japan).

RESULTS

The characteristics of the 600 participants at baseline and after 3 years are shown in Table 1. The number of participants with onset of periodontal disease was 186 (31.0%) during 3 years. Although the percentage of participants who use dental floss increased during 3 years ($p < .001$), regular dental check-up and smoking state turned worse ($p = .042$ and $p < .001$, respectively). “INSERT HERE (Table 1)”

Table 2 shows the association between Type D personality and oral health behaviors at baseline and after 3 years. A significant difference was not observed between Type D and non-Type D personality at both baseline and follow-up. “INSERT HERE (Table 2)”

The comparisons of covariates between the participants with and without progression of periodontal disease are shown in Table 3. The percentage of males with onset of periodontal disease after 3 years was 54.8%, while that without the onset of periodontal disease was 43.5%. The onset of periodontal disease was more prevalent in males than in females ($p = .010$). In the participants with Type D personality, the percentage of the onset of periodontal disease was higher than those in Non-type D personality ($p = .019$). In addition, there was a significant difference between BMI at baseline and the onset of periodontal disease ($p = .003$), however no significant

differences between oral health behaviors at baseline and the onset of periodontal disease were observed (Table 3). “INSERT HERE (Table 3)”

In the logistic regression analyses, the risk of periodontal disease after 3 years was significantly associated with “BMI \geq 25” (OR: 2.543; 95% CI: 1.297-4.989; p = .007) and “Type D personality” (OR: 1.473; 95% CI: 1.027-2.111; p = .035) (Table 4). The accuracy of discrimination was 69.5%, and the Hosmer–Lemeshow test found the model fit to be acceptable, with chi-square statistics of 7.704 (p = .359). “INSERT HERE (Table 4)”

DISCUSSION

To the best of our knowledge, this is the first study which adopted Type D personality in the field of dentistry. In addition, we are the first to report the prevalence of Type D personality in Japanese young population. Our findings suggest that Type D personality is associated with the onset of periodontal disease during a 3-year-period in young adults (Table 4). Although there are many risk factors, including diabetes (Simpson et al., 2010) and smoking (Tonetti, 1998), for periodontal disease, Type D personality may be evaluated as a new risk factor. In addition, the evaluation of Type D personality might be effective in the field of public health activity, because measuring Type D personality is easy (Denollet, 2005), inexpensive and can predict the occurrence of other diseases or health behaviors (Howard and Hughes, 2012; Kupper et al., 2007; Nefs et al., 2012; Schiffer et al., 2010; Yildirim et al., 2013).

Cortisol and stress are the common factors in the association between Type D personality and periodontal disease. Type D individuals tend to experience negative emotions (Denollet, 2000; Habra et al., 2003). Stressful situations involving fear, anxiety, helplessness, and loss of control result in release of cortisol (Sher, 2005). On the other hand, stress is one of the important risk factors for periodontal disease, and cortisol levels are associated with the extent and severity of periodontitis (Hilgert et al.,

2006). Exposure to stress can induce the release of noradrenaline through the sympathetic nervous system by activation of the adrenal medulla, which can have immunosuppressive effects. Such immunosuppressive effect can enhance periodontal tissue destruction (Breivik and Thrane, 2011). Since the stress or cortisol levels of the participants in this study were not measured, we could not make certain whether Type D personality preceded the stress related to the onset of periodontal disease. Clarifying the mechanisms to explain the association among periodontal disease, stress and Type D personality is the next critical step.

Type D individuals engaged in unhealthier behaviors including smoking, poor diet and lack of physical activity and less likely to get regular medical check-ups than non-Type D individuals (Williams et al., 2008). However, there was no relationship between Type D personality and oral health behaviors in our results (Table 2). The discrepancy between our study and the previous study may depend on field (Japan vs. United Kingdom and Ireland). However, further studies may be needed to examine the relationship.

In this study, the risk of the onset of periodontal disease during 3 years was significantly associated with BMI \geq 25 (overweight /obesity) at baseline. Morita et al. (2011) showed a direct dose–response relationship between BMI and the subsequent

development of periodontal disease in a population of 3,590 Japanese individuals followed over 5 years. Our data are consistent with the previous study. Since obese individuals develop infections of various types more often than those with normal weight (Falagas and Kompoti, 2006), improving overweight during undergraduate days might be effective to prevent the onset of periodontal disease.

In our study, the prevalence of students with Type D personality was found to be 39.2% (Table 1). A previous study in UK reported that the prevalence of Type D was 38.5% in 1,012 healthy young adults (mean age 20.5 years), which supports our result (Williams et al., 2008). On the other hand, a survey in Japan showed the prevalence of elderly Japanese people (≥ 65) with Type D personality was 46.2%. The discrepancy of prevalence may be mainly due to differences in age groups (Kasai et al., 2013). Although many studies explained that Type D personality predicts the occurrence of various diseases (Nefs et al., 2012; Schiffer et al., 2010; Yildirim et al., 2013), it is unknown whether a disease affects the personality yet. Further studies may need to examine the relationship among Type D personality and age, race, or diseases.

The presence of periodontal disease was defined as presence of both PPD ≥ 4 mm and BOP $\geq 30\%$. BOP is an earlier and sensitive indicator of inflammation (Greenstein, 1984), and deepened PPD with BOP (+) increases the risk of progression of periodontal

disease (Oliveira Costa et al., 2011). Dentists usually diagnose periodontal disease using the results of probing pocket depth, clinical attachment level, bleeding on probing and destruction of the alveolar bone. Thus, full-mouth periodontal examination is recommended (Dye et al., 2014), however it is rather invasive and unsuitable for screening. We examined only 10 teeth, which can be underestimated compared with full-mouth examination.

The present study had other limitations. First, all participants were recruited from among students at Okayama University. Thus, extrapolation of our findings to a general population of young-aged people may be limited. Second, the follow-up rate in our current study was low (30.1%), as applicants received the 3-year follow-up examinations only just prior to graduation. However, there were no significant differences in oral health behavior, periodontal states and body composition at baseline between participants who were followed up and those who were not (data not shown). Thus, the effect of sampling bias on the results may be small. Finally, we used BMI as measure of body composition. Although BMI is often used to assess obesity in young people (Ekuni et al., 2014), we did not know whether it is muscle or obesity.

In conclusion, a significant association between Type D personality and the onset of periodontal disease was observed in Japanese university students.

Conflicts of interest

The authors have no conflicts of interest to declare.

Funding

This study was self-supported.

Figure legend

Figure 1. Flow chart of participants

REFERENCES

- Booth L, Williams L (2015) Type D personality and dietary intake: The mediating effects of coping style. *Journal of Health Psychology* 20: 921-927.
- Breivik T and Thrane OS (2011) Psychoneuroimmune interactions in periodontal disease. In: Ader R, Felten DL, Cohen N (eds) *Psychoneuroimmunology*. vol 63. New York: Academic Press, pp. 627-644.
- Condén E, Ekselius L, Aslund C (2013) Type D personality is associated with sleep problems in adolescents. Results from a population-based cohort study of Swedish adolescents. *Journal of Psychosomatic Research* 74: 290-295.
- Denollet J (2000) Type D personality. A potential risk factor defined. *Journal of Psychosomatic Research* 49: 255-266.
- Denollet J (2005) DS14: standard assessment of negative affectivity, social inhibition, and Type D personality. *Psychosomatic Medicine* 67: 89-97.
- Denollet J, Sys S, Stoobant N, et al. (1996) Personality as independent predictor of long-term mortality in patients with coronary heart disease. *Lancet* 347: 417-421.
- Dye BA, Li X, Lewis BG et al. (2014) Overview and quality assurance for the oral health component of the National Health and Nutrition Examination Survey (NHANES), 2009-2010. *Journal of Public Health Dentistry* 74: 248-256.

Eke PI, Page RC, Wei L, et al. (2012) Update of the case definitions for population-based surveillance of periodontitis. *Journal of Periodontology* 83: 1449-1454.

Ekuni D, Mizutani S, Kojima A, et al. (2014) Relationship between increases in BMI and changes in periodontal status: a prospective cohort study. *Journal of Clinical Periodontology* 41: 772-778.

Emons WH, Meijer RR, Denollet J (2007) Negative affectivity and social inhibition in cardiovascular disease: evaluating type-D personality and its assessment using item response theory *Journal of Psychosomatic Research* 63: 27-39.

Falagas ME and Kompoti M (2006) Obesity and infection. *The Lancet. Infectious Diseases* 6: 438-446.

Furuta M, Ekuni D, Irie K, et al. (2011) Sex differences in gingivitis relate to interaction of oral health behaviors in young people. *Journal of Periodontology* 82: 558-565.

Genco RJ and Borgnakke WS (2013) Risk factors for periodontal disease. *Periodontology 2000* 62: 59-94.

Gilmour J, Williams L (2012) Type D personality is associated with maladaptive health-related behaviours. *Journal of Health Psychology* 17: 471-478.

Ginting H, van de Ven M, Becker ES, et al. (2016) Type D personality is associated with

health behaviors and perceived social support in individuals with coronary heart disease.

Journal of Health Psychology 21: 727-737.

Greenstein G (1984) The role of bleeding upon probing in the diagnosis of periodontal disease. A literature review. *Journal of Periodontology* 55: 684-688.

Habra ME, Linden W, Anderson JC, et al. (2003) Type D personality is related to cardiovascular and neuroendocrine reactivity to acute stress. *Journal of Psychosomatic Research* 55: 235-245.

Hilgert JB, Hugo FN, Bandeira DR et al. (2006) Stress, cortisol, and periodontitis in a population aged 50 years and over. *Journal of Dental Research* 85: 324-328.

Howard S and Hughes BM (2012) Construct, concurrent and discriminant validity of Type D personality in the general population: associations with anxiety, depression, stress and cardiac output. *Psychology & Health* 27: 242-258.

Kasai Y, Suzuki E, Iwase T, et al. (2013) Type D personality is associated with psychological distress and poor self-rated health among the elderly: a population-based study in Japan. *PLoS One* 8: e77918.

Kawabata Y, Ekuni D, Miyai H, et al. (2016) Relationship Between Prehypertension/Hypertension and Periodontal Disease: A Prospective Cohort Study. *American Journal of Hypertension* 29: 388-396.

- Kupper N, Denollet J, de Geus EJ, et al. (2007) Heritability of type-D personality. *Psychosomatic Medicine* 69: 675-681.
- Martinez-Maestre MA, Gonzalez-Cejudo C, Machuca G, et al. (2010) Periodontitis and osteoporosis: a systematic review. *Climacteric* 13: 523-529.
- Meng H, Ren X, Tian Y, et al. (2011) Genetic study of families affected with aggressive periodontitis. *Periodontology 2000* 56: 87-101.
- Mizutani S, Ekuni D, Furuta M, et al. (2012) Effects of self-efficacy on oral health behaviors and gingival health in university students aged 18- or 19-years-old. *Journal of Clinical Periodontology* 39: 844-849.
- Mizutani S, Ekuni D, Tomofuji T, et al. (2015) Self-efficacy and progression of periodontal disease: A prospective cohort study. *Journal of Clinical Periodontology* 42: 1083-1089.
- Mommersteeg PM, Denollet J, Martens EJ (2012) Type D personality, depressive symptoms and work-related health outcomes. *Scandinavian Journal of Public Health* 40: 35-42.
- Morita I, Okamoto Y, Yoshii S, et al. (2011) Five-year incidence of periodontal disease is related to body mass index. *Journal of Dental Research* 90: 199-202.
- Nefs G, Pouwer F, Pop V, et al. (2012) Type D (distressed) personality in primary care

patients with type 2 diabetes: validation and clinical correlates of the DS14 assessment.

Journal of Psychosomatic Research 72: 251-257.

Newton JT and Asimakopoulou K (2015) Managing oral hygiene as a risk factor for periodontal disease: a systematic review of psychological approaches to behaviour change for improved plaque control in periodontal management. *Journal of Clinical Periodontology* 42: S36-46.

Oginska-Bulik N (2006) Occupational stress and its consequences in healthcare professionals: the role of type D personality. *International Journal of Occupational Medicine and Environmental Health* 19: 113-122.

Oliveira Costa F, Miranda Cota LO, Pereira Lages EJ, et al (2011) Progression of periodontitis in a sample of regular and irregular compliers under maintenance therapy: a 3-year follow-up study. *Journal of Periodontology* 82: 1279-1287.

Peruzzo DC, Benatti BB, Ambrosano GM, et al. (2007) A systematic review of stress and psychological factors as possible risk factors for periodontal disease. *Journal of Periodontology* 78: 1491-1504.

Schiffer AA, Smith OR, Pedersen SS, et al. (2010) Type D personality and cardiac mortality in patients with chronic heart failure. *International Journal of Cardiology* 142: 230-235.

Sher L (2005) Type D personality: the heart, stress, and cortisol. *QJM : monthly journal of the Association of Physicians* 98: 323-329.

Simpson TC, Needleman I, Wild SH, et al. (2010) Treatment of periodontal disease for glycaemic control in people with diabetes. *The Cochrane Database of Systematic Reviews* 12: CD004714.

Suvan J, D'Aiuto F, Moles DR, et al. (2011) Association between overweight/obesity and periodontitis in adults. A systematic review. *Obesity reviews* 12: e381-e404.

Tonetti MS (1998) Cigarette smoking and periodontal diseases: etiology and management of disease. *Annals of Periodontology* 3: 88-101.

Wiencierz S, Williams L (2016) Type D personality and physical inactivity: The mediating effects of low self-efficacy. *Journal of Health Psychology* [Epub ahead of print].

Williams L, Abbott C, Kerr R (2015) Health behaviour mediates the relationship between Type D personality and subjective health in the general population. *Journal of Health Psychology* [Epub ahead of print].

Williams L, O'Connor RC, Howard S et al. (2008) Type-D personality mechanisms of effect: the role of health-related behavior and social support. *Journal of Psychosomatic Research* 64: 63-69.

Wimmer G, Kohldorfer G, Mischak I, et al. (2005) Coping with stress: its influence on periodontal therapy. *Journal of Periodontology* 76: 90-98.

Yildirim O, Alçelik A, Canan F, et al. (2013) Impaired subjective sleep quality in irritable bowel syndrome patients with a Type D personality. *Sleep and Biological Rhythms* 11: 135-138.

Table 1. Participant characteristics ($N = 600$)

Covariate		Baseline	After 3 years
Age		-	21.22 ± 0.57
Gender	Male	282 (47.0)	-
	Female	318 (53.0)	-
Type D personality	Non-Type D	-	365 (60.8)
	Type D	-	235 (39.2)
Presence of periodontal disease	Yes	0 (0.0)	186 (31.0)
	No	600 (100.0)	414 (69.0)
Tooth brushing (daily frequency)	≤ One time	102 (17.0)	95 (15.8)
	≥ Two times	498 (83.0)	505 (84.2)
Usage of dental floss	Yes	35 (5.8)	91 (15.2)
	No	565 (94.2)	509 (84.8)
Regular dental check-up	Yes	100 (16.7)	77 (12.8)
	No	500 (83.3)	523 (87.2)
Smoking state	Current	2 (0.3)	27 (4.5)
	Never	598 (99.7)	573 (95.5)

Note. Periodontal disease was defined as having probing pocket depth ≥ 4 mm and BOP $\geq 30\%$; Values are reported as mean \pm standard deviation or number (percentage).

Table 2. Association between Type D personality and oral health behaviors at baseline and after 3 years in participants ($N = 600$)

Oral health behaviors		Baseline			After 3 years		
		Type D $N = 235$	Non-Type D $N = 365$	p value	Type D $N = 235$	Non-Type D $N = 365$	p value
Age		-	-	-	21.26 ± 0.74	21.20 ± 0.44	.053
Gender	Male	119 (50.6)	163 (44.7)	.155	-	-	-
	Female	116 (49.4)	202 (55.3)		-	-	-
Body Mass Index	< 22.0	168 (71.5)	281 (77.0)	.389	-	-	-
	22.0 to 24.9	47 (20.0)	64 (17.5)		-	-	-
	25.0 to 29.9	19 (8.1)	19 (5.2)		-	-	-
	≥ 30.0	1 (0.4)	1 (0.3)		-	-	-
Tooth brushing (daily frequency)	\leq One time	46 (19.6)	56 (15.3)	.178	43 (18.3)	52 (14.2)	.185
	\geq Two times	189 (80.4)	309 (84.7)		192 (81.7)	313 (85.8)	
Usage of dental floss	Yes	12 (5.1)	23 (6.3)	.542	29 (12.3)	62 (17.0)	.121
	No	223 (94.9)	342 (93.7)		206 (87.7)	303 (83.0)	
Regular dental check-up	Yes	46 (19.6)	54 (14.8)	.125	29 (12.9)	48 (13.2)	.772
	No	189 (80.4)	311 (85.2)		206 (87.7)	317 (86.8)	
Smoking state	Yes	1 (0.4)	1 (0.3)	1.000	12 (5.1)	15 (4.1)	.565
	No	234 (99.6)	364 (99.7)		223 (94.9)	350 (95.9)	

Note. Values are reported as number (percentage); P value obtained with chi-square test or Fisher's exact test (Type D vs. Non-type D).

Table 3. Association between onset of periodontal disease and related factors ($N = 600$)

Parameters at baseline		Onset of periodontal disease		p value
		No ($N = 414$)	Yes ($N = 186$)	
Age		18.2 \pm 0.72	18.2 \pm 0.46	.497
Gender	Male	180 (43.5)	102 (54.8)	.010
	Female	234 (56.5)	84 (45.2)	
Type D personality	Non-Type D	265 (64.0)	100 (53.8)	.019
	Type D	149 (36.0)	86 (46.2)	
Body Mass Index	< 22.0	325 (78.5)	124 (66.7)	.003
	22.0 to 24.9	69 (16.7)	42 (22.6)	
	25.0 to 29.9	18 (4.8)	20 (10.8)	
	\geq 30.0	2 (0.5)	0 (0.0)	
Tooth brushing (daily frequency)	\leq One time	76 (18.4)	26 (14.0)	.187
	\geq Two times	338 (81.6)	160 (86.0)	
Usage of dental floss	Yes	26 (6.3)	9 (4.8)	.486
	No	388 (93.7)	177 (95.2)	
Regular dental check-up	Yes	71 (17.1)	29 (15.6)	.636
	No	343 (82.9)	157 (84.4)	
Smoking state	Smoker	1 (0.2)	1 (0.5)	.561
	Non-smoker	413 (99.8)	185 (99.5)	

Note. Values are reported as mean \pm standard deviation or number (percentage); P value obtained with Unpaired t-test, chi-square test or Fisher's exact test, as appropriate.

Table 4. Adjusted odds ratios and 95% CIs for onset of periodontal disease.

		Odds ratio	95% CI	p value
Age		1.363	0.988-1.881	.059
Gender	Female	1 (Reference)		
	Male	1.414	0.983-2.033	.062
Body Mass Index	< 22.0	1 (Reference)		
	22.0 to 24.9	1.537	0.984-2.399	.059
	≥ 25	2.543	1.297-4.989	.007
Type D personality	Non-Type D	1 (Reference)		
	Type D	1.473	1.027-2.111	.035
Tooth brushing (daily frequency)	≥ Two times	1 (Reference)		
	≤ One time	0.632	0.382-1.046	.074
Regular dental check-up	Yes	1 (Reference)		
	No	1.150	0.708-1.868	.573

Note. Adjusted for age, gender, Body Mass Index, Type D personality, tooth brushing (daily frequency) and regular dental check-up.