

## A methodology for numerical simulations to a singular limit

**Ikota, Ryo**  
Faculty of Mathematics, Kyushu University

**Mimura, Masayasu**  
Department of Mathematical and Life Sciences, Hiroshima University

**Nakaki, Tatsuyuki**  
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/3347>

---

出版情報 : MHF Preprint Series. 2004-6, 2004-03-08. 九州大学大学院数理学研究院  
バージョン :  
権利関係 :

# MHF Preprint Series

Kyushu University  
21st Century COE Program  
Development of Dynamic Mathematics with  
High Functionality

## A methodology for numerical simulations to a singular limit

R. Ikota, M. Mimura  
T. Nakaki

MHF 2004-6

( Received March 8, 2004 )

Faculty of Mathematics  
Kyushu University  
Fukuoka, JAPAN

# A methodology for numerical simulations to a singular limit

Ryo Ikota

Faculty of Mathematics,  
Kyushu University, Fukuoka, Japan

Masayasu Mimura

Department of Mathematical and Life Sciences,  
Hiroshima University, Hiroshima, Japan

Tatsuyuki Nakaki

Faculty of Mathematics,  
Kyushu University, Fukuoka, Japan

## Abstract

A numerical methodology is proposed for a singular limit of certain reaction-diffusion systems. The limit problem arises in competition-diffusion systems and chemical reaction equations. Convergence of the semi-discrete-in-time solutions obtained by the methodology is proved. In a particular case, a convergence rate is also shown.

*Keywords:* reaction-diffusion systems, singular limit, operator splitting method

## 1 Introduction

In ecological and chemical problems, we encounter nonlinear equations of the form

$$w_t = \nabla \cdot (d(w)\nabla w) + h(w) \quad x \in \Omega, \quad t > 0, \quad (1.1)$$

$$\frac{\partial w}{\partial \nu} = 0 \quad x \in \partial\Omega, \quad t > 0, \quad (1.2)$$

$$w(0, x) = w_0(x) \in L^\infty(\Omega) \quad x \in \Omega, \quad (1.3)$$

where  $d(s)$  is a step function:

$$d(s) = \begin{cases} d_1 & (s \geq 0), \\ d_2 & (s < 0). \end{cases} \quad (1.4)$$

The function  $w = w(t, x)$  is real-valued,  $d_1$  and  $d_2$  are positive constants,  $\Omega$  is a bounded region in  $\mathbb{R}^N$  with a smooth boundary  $\partial\Omega$ , and  $\nu$  is the unit outer normal to  $\partial\Omega$ . The aim of this paper is to propose a numerical methodology for (1.1)–(1.3).

One origin of (1.1)–(1.3) is in theoretical ecology. A reaction-diffusion system, specifically called a competition-diffusion system, has been studied as a model of spatially distributed competing species (see Cantrell (1996), Cosner & Lazer (1984), Dancer, Hilhorst, Mimura & Peletier (1999), Ei, Ikota & Mimura (1999), Iida, Muramatsu, Ninomiya & Yanagida (1998), Mimura & Fife (1986) for examples). Let  $u_i(t, x)$  be the population density of an  $i$ th competing

species  $U_i$  ( $i = 1, 2, \dots, n$ ) at the time  $t > 0$  and the position  $x \in \Omega$ . Then the competition-diffusion system is written as

$$\frac{\partial u_i}{\partial t} = d_i \Delta u_i + (r_i - \sum_{j=1}^n a_{ij} u_j) u_i \quad (i = 1, 2, \dots, n), \quad x \in \Omega, \quad t > 0, \quad (1.5)$$

where  $d_i$  is the diffusion rate,  $r_i$  the intrinsic growth rate,  $a_{ii}$  the intraspecific competition rate, and  $a_{ij}$  ( $i \neq j$ ) the interspecific competition rate between  $U_i$  and  $U_j$ . These parameters are all positive constants. We impose the homogeneous Neumann boundary condition and suppose the initial functions are non-negative.

One of our interests in (1.5) is spatially segregating patterns of the solutions, which appear when the interspecific competition rates  $a_{ij}$  ( $i \neq j$ ) are large. Let  $k$  be a positive parameter and put  $a_{ij} = kb_{ij}$  ( $i \neq j$ ). Then from (1.5) we have

$$\frac{\partial u_i}{\partial t} = d_i \Delta u_i + (r_i - a_{ii} u_i) u_i - k \sum_{j \neq i} b_{ij} u_i u_j \quad (i = 1, 2, \dots, n) \quad x \in \Omega, \quad t > 0. \quad (1.6)$$

What we have to observe is behavior of the solutions as  $k$  is large.

On the asymptotic behavior, an analytical result is known in the case  $n = 2$ . By putting  $u = u_1$  and  $v = b_{12} u_2$  the equations (1.6) lead to the following form:

$$u_t = d_1 \Delta u + f(u)u - kuv \quad x \in \Omega, \quad t > 0, \quad (1.7)$$

$$v_t = d_2 \Delta v + g(v)v - \alpha kuv \quad x \in \Omega, \quad t > 0, \quad (1.8)$$

where  $\alpha = b_{21}$ . The boundary and initial conditions are

$$\frac{\partial u}{\partial \nu} = 0, \quad \frac{\partial v}{\partial \nu} = 0 \quad x \in \partial \Omega, \quad t > 0, \quad (1.9)$$

$$u(0, x) = u_0(x), \quad v(0, x) = v_0(x) \quad x \in \Omega. \quad (1.10)$$

The functions  $f$  and  $g$  are written as  $f(s) = a_u(R_u - s)$  and  $g(s) = a_v(R_v - s)$ , where  $a_u = a_{11}$ ,  $R_u = r_1/a_{11}$ ,  $a_v = a_{22}/b_{12}$  and  $R_v = r_2 b_{12}/a_{22}$ . Thus they satisfy

$$f(s) > 0 \quad (0 < s < R_u), \quad f(s) < 0 \quad (s > R_u), \quad (1.11)$$

$$g(s) > 0 \quad (0 < s < R_v), \quad g(s) < 0 \quad (s > R_v). \quad (1.12)$$

Let  $(u^{(k)}, v^{(k)})$  be a solution to (1.7)–(1.10). Dancer et al. (1999) have shown that as  $k \rightarrow \infty$  the function  $w^{(k)} = u^{(k)} - v^{(k)}/\alpha$  converges to a weak solution  $w$  to (1.1)–(1.3) with a function  $h$  such that

$$h(s) = \begin{cases} f(s)s & (s \geq 0), \\ g(-\alpha s)s & (s < 0). \end{cases} \quad (1.13)$$

Furthermore they proved that  $u^{(k)} \rightarrow [w]^+$  and  $v^{(k)} \rightarrow \alpha[w]^-$  respectively, where  $[a]^\pm$  is  $\max\{\pm a, 0\}$ . See Proposition 2.1 in Dancer et al. (1999) for the detail. Thus (1.1)–(1.3) describe the asymptotic behavior of (1.7)–(1.10).

Another origin of (1.1)–(1.3) is chemical reaction equations, which we obtain by dropping the nonlinear terms  $f$  and  $g$  from (1.7)–(1.10). The singular limits of the chemical reaction equations have been studied by Evans and Tonegawa earlier on. Evans (1980) gave convergence proof for restrictive initial data. Tonegawa (1998) proved regularity properties of solutions to the limiting problem.

To solve (1.1)–(1.3) numerically, for example, we can apply finite volume methods (FVM) (Eymard, Gallouët, Hilhorst & Slimane 1998). FVM is capable of dealing with wide range of nonlinear diffusion equations.

Although the target of our numerical methodology, named *Threshold Competition Dynamics* (TCD), is rather restricted, it has flexibility in implementations; TCD includes a process to solve reaction-diffusion equations, to which we can apply finite difference methods, finite element methods or others. Hence, even if the shape of  $\Omega$  is complicated we can use TCD without numerical difficulties. In addition, TCD has possibility to solve the singular limits of (1.6) in the case  $n \geq 3$ . In fact we have applied TCD to a three-component competition-diffusion system (Ikota, Mimura & Nakaki 2001).

We should note that TCD is similar to the so-called diffusion-generated approach for mean curvature flow (see Merriman, Bence & Osher (1994) and Ruuth (1998)). Its convergence has been proved by Barles & Georgelin (1995), Evans (1993), and for more general geometric motions of hypersurfaces by Ishii, Pires & Souganidis (1999).

Throughout the rest of this paper we assume  $h$  satisfies (1.13) with (1.11) and (1.12). It should be noticed that all the statements are still valid for the case  $h \equiv 0$ .

This paper is organized as follows. In the next section we present TCD and state our results. The results are composed of two theorems. One refers to the convergence of TCD in a general situation. The other gives a convergence rate in a restricted situation. The former theorem is proved in section 3 and the latter in section 4. In order to demonstrate practical usefulness of TCD, we perform numerical experiments in section 5. In the appendix we see notation index.

## 2 Results

A scheme is shown and results on it are stated in this section.

Put

$$F(s) := f(s)s, \quad G(s) := g(s)s. \quad (2.1)$$

Then the scheme that we propose is written as follows.

### Threshold Competition Dynamics (TCD)

Let  $M$  be a positive integer. The approximate solution  $(u_M(t, x), v_M(t, x))$  by TCD to the limiting problem of (1.7)–(1.10) as  $k \rightarrow \infty$  is defined by

$$u_M(0, x) = u_0(x), \quad v_M(0, x) = v_0(x) \quad \text{for } x \in \Omega, \quad (2.2)$$

$$u_M(t, x) = \bar{u}_M^j(t, x), \quad v_M(t, x) = \bar{v}_M^j(t, x), \quad \text{for } t \in (t_j, t_{j+1}], \quad x \in \Omega, \quad (2.3)$$

where

$$\tau := T/M, \quad t_j := j\tau \quad (j = 0, 1, \dots, M). \quad (2.4)$$

The functions  $\bar{u}_M^j(t, x)$  and  $\bar{v}_M^j(t, x)$  are constructed by the following steps:

Step 1. Put  $u_M^0(x) = u_0(x)$ ,  $v_M^0(x) = v_0(x)$  ( $x \in \Omega$ ).

Step 2. For given  $u_M^j(x)$  and  $v_M^j(x)$ ,

(i) Find  $\bar{u}_M^j(t, x)$  and  $\bar{v}_M^j(t, x)$  such that

$$\begin{cases} \frac{\partial \bar{u}_M^j}{\partial t} = d_1 \Delta \bar{u}_M^j + F(\bar{u}_M^j) & x \in \Omega, \quad t_j < t < t_{j+1}, \\ \frac{\partial \bar{u}_M^j}{\partial \nu} = 0 & x \in \partial\Omega, \quad t_j < t < t_{j+1}, \\ \bar{u}_M^j(t_j, x) = u_M^j(x) & x \in \Omega, \end{cases} \quad (2.5)$$

$$\begin{cases} \frac{\partial \bar{v}_M^j}{\partial t} = d_2 \Delta \bar{v}_M^j + G(\bar{v}_M^j) & x \in \Omega, \quad t_j < t < t_{j+1}, \\ \frac{\partial \bar{v}_M^j}{\partial \nu} = 0 & x \in \partial\Omega, \quad t_j < t < t_{j+1}, \\ \bar{v}_M^j(t_j, x) = v_M^j(x) & x \in \Omega. \end{cases} \quad (2.6)$$

(ii) Define  $u_M^{j+1}(x)$  and  $v_M^{j+1}(x)$  by

$$u_M^{j+1}(x) = \lim_{\theta \rightarrow \infty} \hat{u}_M^j(\theta; x), \quad v_M^{j+1}(x) = \lim_{\theta \rightarrow \infty} \hat{v}_M^j(\theta; x), \quad (2.7)$$

where  $\hat{u}_M^j$  and  $\hat{v}_M^j$  solve

$$\begin{cases} \frac{d\hat{u}_M^j}{d\theta} = -\hat{u}_M^j \hat{v}_M^j & x \in \Omega, \quad 0 < \theta < k\tau, \\ \frac{d\hat{v}_M^j}{d\theta} = -\alpha \hat{u}_M^j \hat{v}_M^j & x \in \Omega, \quad 0 < \theta < k\tau, \\ \hat{u}_M^j(0; x) = \bar{u}_M^j(x, t_{j+1}), \quad \hat{v}_M^j(0; x) = \bar{v}_M^j(x, t_{j+1}), & x \in \Omega. \end{cases} \quad (2.8)$$

We note that an operator-splitting method is used in Step 2, that is, (1.7) and (1.8) are splitted into

$$u_t = d_1 \Delta u + F(u), \quad v_t = d_2 \Delta v + G(v), \quad (2.9)$$

and

$$\frac{du}{dt} = -kuv, \quad \frac{dv}{dt} = -k\alpha uv. \quad (2.10)$$

The main idea of TCD is Step 2 (ii). Let  $\theta = kt$ ; then (2.10) are rewritten to (2.8). Instead of passing to the limit  $k \rightarrow \infty$  in (2.10), we use the asymptotic limit  $\theta \rightarrow \infty$  in a solution to (2.8). The limit is easily obtained. In fact, by using the fact that  $d(u - v/\alpha)/d\theta = 0$ , it follows that

$$\lim_{\theta \rightarrow \infty} \hat{u}_M^j(\theta; x) = \left[ \bar{u}_M^j(t_{j+1}, x) - \bar{v}_M^j(t_{j+1}, x)/\alpha \right]^+, \quad (2.11)$$

$$\lim_{\theta \rightarrow \infty} \hat{v}_M^j(\theta; x) = \alpha \left[ \bar{u}_M^j(t_{j+1}, x) - \bar{v}_M^j(t_{j+1}, x)/\alpha \right]^-. \quad (2.12)$$

We now define a weak solution to (1.1)–(1.3).

**Definition 2.1.** *We call  $w$  a weak solution if it satisfies:*

$$w \in L^\infty(\Omega \times (0, T)) \cap L^2(0, T; H^1(\Omega)) \cap C([0, T]; L^2(\Omega)), \quad (2.13)$$

$$\int_\Omega w(T)\phi(T) - \iint_{Q_T} \{w\phi_t - d(w)\nabla w \nabla \phi + h(w)\phi\} = \int_\Omega w_0\phi(0), \quad (2.14)$$

for all  $\phi \in C^1(\bar{Q}_T)$ , where  $Q_T = \Omega \times (0, T)$ .

**Remark 2.1.** *There exists a unique solution to (2.13)–(2.14) if  $w_0 \in L^\infty(\Omega)$  (Dancer et al. 1999).*

We are ready to state our results.

**Theorem 2.1.** *Suppose  $w_0 \in L^\infty(\Omega)$ . Set  $u_0 = [w_0]^+$  and  $v_0 = \alpha[w_0]^-$ . Let  $w$  be a weak solution for the initial data  $w_0$  and  $(u_M, v_M)$  an approximate solution by Threshold Competition Dynamics for the initial data  $(u_0, v_0)$ . Then  $u_M$ ,  $v_M$  and  $w_M = u_M - v_M/\alpha$  converge to  $[w]^+$ ,  $\alpha[w]^-$  and  $w$  in  $L^2(0, T; L^2(\Omega))$  respectively as  $M$  tends to  $\infty$ .*

Moreover, if  $d_1 = d_2$  we have information about the convergence rate.

**Theorem 2.2.** *Functions  $w_0$ ,  $u_0$ ,  $v_0$ ,  $w_M$  and  $w$  are the same as those in Theorem 2.1. Assume that*

$$u_0, v_0 \in H^2(\Omega) \cap L^\infty(\Omega), \quad (2.15)$$

and

$$\left. \begin{aligned} \frac{\partial u_0}{\partial \nu} = \frac{\partial v_0}{\partial \nu} = 0 \quad \text{on } \partial\Omega, \\ \int_{\Omega} u_0 v_0 = 0. \end{aligned} \right\} \quad (2.16)$$

In addition if  $d_1 = d_2$ , then

$$\|(u_M(T) - v_M(T)/\alpha) - w(T)\|_{L^2(\Omega)} \leq C(1/M)^{1/2}, \quad (2.17)$$

$$\|(u_M(T) - v_M(T)/\alpha) - w(T)\|_{L^1(\Omega)} \leq C'(1/M), \quad (2.18)$$

where  $C$  and  $C'$  are positive constants independent of  $M$ .

### 3 Proof of Theorem 2.1

We use the evolution triple  $H^1(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow (H^1(\Omega))^*$  (see chapter 23 in Zeidler (1990) or chapter 3 in Temam (1984)) and prove the following four lemmas.

**Lemma 3.1.** *Functions  $u_M$  and  $v_M$  are uniformly bounded with respect to  $M$  in  $L^\infty(\Omega)$ . More precisely, for any  $t$  ( $0 \leq t \leq T$ )  $u_M$  and  $v_M$  satisfy*

$$0 \leq u_M(t, x) \leq \max\{R_u, \|u_0\|_{L^\infty(\Omega)}\} \quad \text{a.e. in } \Omega, \quad (3.1)$$

$$0 \leq v_M(t, x) \leq \max\{R_v, \|v_0\|_{L^\infty(\Omega)}\} \quad \text{a.e. in } \Omega. \quad (3.2)$$

*Proof.* We establish only (3.1) because the same argument yields (3.2).

Recall that

$$u_M^j(x) = \left[ \bar{u}_M^{j-1}(t_j, x) - \bar{v}_M^{j-1}(t_j, x)/\alpha \right]^+ \leq \bar{u}_M^{j-1}(t_j, x).$$

Thus it suffices to show that

$$0 \leq \bar{u}_M^j(t, x) \leq \max\{R_u, \|u_M^j\|_{L^\infty(\Omega)}\} \quad \text{a.e. in } \Omega \quad (t_j \leq t \leq t_{j+1}). \quad (3.3)$$

By regularizing the function  $u_M^j(x)$ , we have only to show (3.3) for smooth solutions to (2.5). For smooth solutions, (3.3) is easily deducible from the comparison theorem.  $\square$

**Lemma 3.2.** *Functions  $u_M$ ,  $v_M$  and  $w_M$  are uniformly bounded in  $L^2(0, T; H^1(\Omega))$  with respect to  $M$ .*

*Proof.* We see in the scalar distribution sense on  $(t_j, t_{j+1})$  that

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} (\bar{u}_M^j)^2 = -d_1 \int_{\Omega} |\nabla \bar{u}_M^j|^2 + \int_{\Omega} F(\bar{u}_M^j) \bar{u}_M^j.$$

Thus we have

$$d_1 \int_{t_j}^{t_{j+1}} \int_{\Omega} |\nabla \bar{u}_M^j|^2 = \frac{1}{2} \left[ \lim_{t \downarrow t_j} \int_{\Omega} (u_M)^2 - \lim_{t \uparrow t_{j+1}} \int_{\Omega} (u_M)^2 \right] + \int_{t_j}^{t_{j+1}} \int_{\Omega} F(u_M) u_M. \quad (3.4)$$

Therefore we obtain

$$\begin{aligned} d_1 \iint_{Q_T} |\nabla u_M|^2 &= \frac{1}{2} \sum_{j=0}^{M-1} \left[ \lim_{t \downarrow t_j} \int_{\Omega} (u_M)^2 - \lim_{t \uparrow t_{j+1}} \int_{\Omega} (u_M)^2 \right] + \iint_{Q_T} F(u_M) u_M \\ &= \frac{1}{2} \int_{\Omega} u_0^2 + \frac{1}{2} \sum_{j=0}^{M-2} \left[ - \lim_{t \uparrow t_{j+1}} \int_{\Omega} (u_M)^2 + \lim_{t \downarrow t_{j+1}} \int_{\Omega} (u_M)^2 \right] \\ &\quad - \frac{1}{2} \lim_{t \uparrow T} \int_{\Omega} (u_M)^2 + \iint_{Q_T} F(u_M) u_M \\ &\leq \frac{1}{2} \int_{\Omega} u_0^2 + \iint_{Q_T} F(u_M) u_M. \end{aligned} \quad (3.5)$$

Combining this and (3.1) we get the stated result for  $u_M$ . The boundedness for  $v_M$  is obtained in the same way. Since  $w_M = u_M - v_M/\alpha$ , it is also uniformly bounded.  $\square$

**Lemma 3.3.**

$$\iint_{Q_T} u_M v_M \rightarrow 0 \quad \text{as } M \rightarrow \infty. \quad (3.6)$$

*Proof.* From the equation (25) in section 23.6 (Zeidler 1990), in each interval  $[t_j, t_{j+1}]$  ( $j = 0, 1, \dots, M-1$ ) we have

$$\begin{aligned} \int_{\Omega} \bar{u}_M^j(t, \cdot) \bar{v}_M^j(t, \cdot) &= \int_{\Omega} \bar{u}_M^j(t, \cdot) \bar{v}_M^j(t, \cdot) - \int_{\Omega} u_M^j(\cdot) v_M^j(\cdot) \\ &= -(d_1 + d_2) \int_{t_j}^t ds \int_{\Omega} \nabla u_M(s, \cdot) \cdot \nabla v_M(s, \cdot) \\ &\quad + \int_{t_j}^t ds \int_{\Omega} \{F(u_M(s, \cdot)) v_M(s, \cdot) + G(v_M(s, \cdot)) u_M(s, \cdot)\} \\ &\leq \frac{d_1 + d_2}{2} \int_{t_j}^t ds \int_{\Omega} (|\nabla u_M(s, \cdot)|^2 + |\nabla v_M(s, \cdot)|^2) + C_1(t - t_j), \end{aligned} \quad (3.7)$$

where

$$\begin{aligned} C_1 &= |\Omega| \sup\{F(p)q + G(q)p \mid \\ &\quad 0 \leq p \leq \max\{R_u, \|u_0\|_{L^\infty(\Omega)}\}, 0 \leq q \leq \max\{R_v, \|v_0\|_{L^\infty(\Omega)}\}\}. \end{aligned}$$



Here notice that  $u_M^j v_M^j = 0$  in  $\Omega$ . Now we observe

$$\begin{aligned} & \int_{t_j}^{t_{j+1}} dt \int_{t_j}^t ds \int_{\Omega} (|\nabla u_M(s, \cdot)|^2 + |\nabla v_M(s, \cdot)|^2) \\ & \leq \int_{t_j}^{t_{j+1}} dt \int_{t_j}^{t_{j+1}} ds \int_{\Omega} (|\nabla u_M(s, \cdot)|^2 + |\nabla v_M(s, \cdot)|^2) \\ & \leq \tau \int_{t_j}^{t_{j+1}} ds \int_{\Omega} (|\nabla u_M(s, \cdot)|^2 + |\nabla v_M(s, \cdot)|^2). \end{aligned}$$

Thus we have

$$\iint_{Q_T} u_M v_M \leq \frac{d_1 + d_2}{2} \tau \int_0^T \int_{\Omega} (|\nabla u_M(s, \cdot)|^2 + |\nabla v_M(s, \cdot)|^2) + \frac{C_1}{2} \tau T \rightarrow 0$$

as  $M \rightarrow \infty$ . □

**Lemma 3.4.** *The generalized derivative  $\partial_t w_M$  exists and it is written as*

$$\partial_t w_M = d_1 \Delta u_M - (d_2/\alpha) \Delta v_M + F(u_M) - G(v_M)/\alpha. \quad (3.8)$$

*Proof.* Although  $u_M$  and  $v_M$  are generally discontinuous as functions from  $[0, T]$  to  $L^2(\Omega)$ ,  $w_M$  belongs to  $C([0, T]; L^2(\Omega))$ . In addition  $\partial_t u_M$  and  $\partial_t v_M$  exist in each interval  $[t_j, t_{j+1}]$ . Hence, for  $z \in H^1(\Omega)$  and  $\psi \in C_0^\infty(0, T)$ , we have

$$\begin{aligned} \int_0^T \int_{\Omega} w_M z \psi_t &= \sum_{j=0}^{M-1} \left[ \int_{\Omega} w_M(t_{j+1}, \cdot) z(\cdot) \psi(t_{j+1}) - \int_{\Omega} w_M(t_j, \cdot) z(\cdot) \psi(t_j) \right. \\ & \quad \left. - \int_{t_j}^{t_{j+1}} \int_{\Omega} (\partial_t u_M - \partial_t v_M / \alpha) z \psi \right] \\ &= - \int_0^T \int_{\Omega} \{d_1 \Delta u_M + F(u_M) - (d_2 \Delta v_M + G(v_M)) / \alpha\} z \psi. \end{aligned}$$

Thus the statement follows. □

Now we are in a position to prove Theorem 2.1.

*Proof of Theorem 2.1.* From Lemmas 3.2 and 3.4, we observe that  $\|w_M\|_{L^2(0, T; H^1(\Omega))}$  and  $\|\partial_t w_M\|_{L^2(0, T; H^1(\Omega)^*)}$  are uniformly bounded with respect to  $M$ . Hence thanks to the compactness property (Theorem 2.1, chapter 3 in Temam (1984)) we obtain a subsequence from  $\{w_M\}$ , which is denoted by  $\{w_M\}$  again, converging in  $L^2(0, T; L^2(\Omega))$ . We write the limit as  $w_\infty$ :

$$w_M \rightarrow w_\infty \quad \text{in } L^2(0, T; L^2(\Omega)) \quad \text{as } M \rightarrow \infty. \quad (3.9)$$

Note that

$$|u - [u - v/\alpha]^+|^2 \leq \frac{uv}{\alpha}, \quad (3.10)$$

$$|v/\alpha - [u - v/\alpha]^-|^2 \leq \frac{uv}{\alpha}. \quad (3.11)$$

Thus  $u_M - [w_M]^+$  and  $v_M/\alpha - [w_M]^-$  converge to 0 in  $L^2(0, T; L^2(\Omega))$  from Lemma 3.3. On the other hand we see that

$$[w_M]^\pm \rightarrow [w_\infty]^\pm \quad \text{in } L^2(0, T; L^2(\Omega)), \quad (3.12)$$

because  $|[w_M]^\pm - [w_\infty]^\pm| \leq |w_M - w_\infty|$ . Therefore we have

$$u_M \rightarrow [w_\infty]^+ \quad \text{in } L^2(0, T; L^2(\Omega)), \quad (3.13)$$

$$v_M/\alpha \rightarrow [w_\infty]^- \quad \text{in } L^2(0, T; L^2(\Omega)). \quad (3.14)$$

Extracting a subsequence again if necessary we see  $u_M$  and  $v_M/\alpha$  converge weakly in  $L^2(0, T; H^1(\Omega))$  and  $\partial_t w_M$  does in  $L^2(0, T; H^1(\Omega)^*)$  from Lemmas 3.2 and 3.4. The limits are identical with  $\partial_t w_M$ ,  $[w_\infty]^+$  and  $[w_\infty]^-$  respectively owing to (3.13), (3.14) and Lemma 3.4:

$$\begin{aligned} \partial_t w_M &\rightarrow \partial_t w_\infty && \text{weakly in } L^2(0, T; H^1(\Omega)^*), \\ u_M &\rightarrow [w_\infty]^+ && \text{weakly in } L^2(0, T; H^1(\Omega)), \\ v_M/\alpha &\rightarrow [w_\infty]^- && \text{weakly in } L^2(0, T; H^1(\Omega)). \end{aligned}$$

In addition, from the Lebesgue's bounded integral lemma we have

$$\begin{aligned} \int_{Q_T} F(u_M)\varphi &\rightarrow \int_{Q_T} F([w_\infty]^+)\varphi, \\ \int_{Q_T} \frac{G(v_M)}{\alpha}\varphi &\rightarrow \int_{Q_T} \frac{G(\alpha[w_\infty]^-)}{\alpha}\varphi. \end{aligned}$$

Now we observe for  $\varphi \in C^1(\overline{Q_T})$  that

$$\begin{aligned} \int_0^T \int_\Omega (\partial_t w_M)\varphi &= - \int_0^T \int_\Omega d_1 \nabla u_M \nabla \varphi + \int_0^T \int_\Omega d_2 \frac{\nabla v_M}{\alpha} \nabla \varphi \\ &\quad + \int_0^T \int_\Omega F(u_M)\varphi - \int_0^T \int_\Omega \frac{G(v_M)}{\alpha}\varphi. \end{aligned}$$

Thus we get

$$\int_0^T \int_\Omega (\partial_t w_\infty)\varphi = - \int_0^T \int_\Omega d(w_\infty) \nabla w_\infty \nabla \varphi + h(w_\infty)\varphi. \quad (3.15)$$

On the other hand, for  $\tilde{\varphi} \in C^1(\overline{Q_T})$  satisfying  $\tilde{\varphi}(T, \cdot) = 0$  we see

$$\int_0^T \int_\Omega (\partial_t w_M)\tilde{\varphi} = - \int_\Omega w_0(\cdot)\tilde{\varphi}(0, \cdot) - \iint_{Q_T} w_M \tilde{\varphi}_t. \quad (3.16)$$

Passing to the limit along a subsequence, we have

$$\int_0^T \int_\Omega (\partial_t w_\infty)\tilde{\varphi} = - \int_\Omega w_0(\cdot)\tilde{\varphi}(0, \cdot) - \iint_{Q_T} w_\infty \tilde{\varphi}_t. \quad (3.17)$$

This implies  $w_\infty(0, \cdot) = w_0(\cdot)$ . Consequently  $w_\infty$  is a weak solution to (1.1)–(1.3). Since the limit is unique, the original whole sequence  $\{w_M\}$  converges to  $w_\infty$ .  $\square$

## 4 Proof of Theorem 2.2

Throughout this section we assume that the conditions for Theorem 2.2 are satisfied. Set

$$e_j^{(p)} := \|w_M(t_j, \cdot) - w(t_j, \cdot)\|_{L^p(\Omega)} \quad (p = 1, 2). \quad (4.1)$$

Our strategy is to deduce a recursive inequality for  $e_j^{(p)}$ .

In this section we choose a positive constant  $R_1$  so that

$$R_1 > \max\{\|u_0\|_{L^\infty(\Omega)}, \|v_0\|_{L^\infty(\Omega)}, R_u, R_v\}. \quad (4.2)$$

In addition we set

$$R_2 = \max\left\{\sup_{0 \leq p \leq R_1} |F(p)|, \sup_{0 \leq p \leq R_1} |G(p)|\right\}, \quad (4.3)$$

$$R_3 = \max\left\{\sup_{0 \leq p \leq R_1} |F'(p)|, \sup_{0 \leq p \leq R_1} |G'(p)|\right\}. \quad (4.4)$$

Here we state the convergence theorem obtained by Dancer et al. (1999).

**Proposition 4.1 (Dancer et al. (1999)).** *Suppose*

$$u_0^{(k)}, v_0^{(k)} \in C(\overline{\Omega}), \quad (4.5)$$

$$0 \leq u_0^{(k)} \leq R_1, \quad 0 \leq v_0^{(k)} \leq R_1, \quad (4.6)$$

$$u_0^{(k)} \rightarrow u_0, \quad v_0^{(k)} \rightarrow v_0, \quad \text{weakly in } L^2(\Omega) \text{ as } k \rightarrow \infty. \quad (4.7)$$

Denote solutions to (1.7)–(1.9) with initial conditions  $u(0, \cdot) = u_0^{(k)}$  and  $v(0, \cdot) = v_0^{(k)}$  by  $u^{(k)}$  and  $v^{(k)}$ . Set  $w^{(k)} = u^{(k)} - v^{(k)}/\alpha$ . Then there exists a weak solution  $w$  to (2.13)–(2.14) such that it satisfies

$$u^{(k)} \rightarrow [w]^+, \quad v^{(k)} \rightarrow \alpha[w]^- \quad \text{weakly in } L^2(0, T; H^1(\Omega)), \quad (4.8)$$

$$u^{(k)} \rightarrow [w]^+, \quad v^{(k)} \rightarrow \alpha[w]^- \quad \text{in } L^1(Q_T), \quad (4.9)$$

$$u^{(k)}v^{(k)} \rightarrow 0 \quad \text{in } L^1(Q_T), \quad (4.10)$$

$$w^{(k)} \rightarrow w \quad \text{in } L^2(Q_T). \quad (4.11)$$

In order to prove Theorem 2.2 we work within the framework of the following spaces:

$$\hat{C}^2(\overline{\Omega}) \stackrel{\text{def}}{=} \left\{v \in C^2(\overline{\Omega}); \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega\right\}, \quad (4.12)$$

$$\hat{C}^3(\overline{\Omega}) \stackrel{\text{def}}{=} \left\{u \in C^3(\overline{\Omega}); \frac{\partial u}{\partial \nu} = 0, \frac{\partial \Delta u}{\partial \nu} = 0 \text{ on } \partial\Omega\right\}. \quad (4.13)$$

The next proposition is obtained by putting  $N = 2$ ,  $N' = 0$ ,  $m = 1$ ,  $q = 1$ ,  $r = 0$ ,  $k = 2$  and  $l = 0$  in Theorem 4.1 of Mora (1983).

**Proposition 4.2.** *The equations (1.7)–(1.9) determines a semiflow of class  $C^0$  on the space  $\hat{C}^3(\overline{\Omega})$ .*

Here we prove several lemmas.

**Lemma 4.1.** *Suppose  $u_0^{(k)} \in \hat{C}^3(\overline{\Omega})$  and  $v_0^{(k)} \in \hat{C}^3(\overline{\Omega})$ . Then solutions  $u^{(k)}$  and  $v^{(k)}$  to (1.7)–(1.9), with initial conditions  $u^{(k)}(0, \cdot) = u_0^{(k)}$  and  $v^{(k)}(0, \cdot) = v_0^{(k)}$ , satisfy*

$$\begin{aligned} & \sup_{0 \leq t \leq T} \left( \int_{\Omega} |u_t^{(k)}(t, \cdot)| + \int_{\Omega} |v_t^{(k)}(t, \cdot)/\alpha| \right) \\ & \leq e^{R_3 T} \int_{\Omega} \left( |d_1 \Delta u_0^{(k)}| + |d_2 \Delta v_0^{(k)}/\alpha| + |F(u_0^{(k)})| + |G(v_0^{(k)})/\alpha| + 2k u_0^{(k)} v_0^{(k)} \right). \end{aligned} \quad (4.14)$$

*Proof.* An argument similar to that in Evans (1980) provides the following inequality:

$$\begin{aligned} \int_{\Omega} (|u_t(T, \cdot)| + |v_t(T, \cdot)/\alpha|) \\ \leq \int_{\Omega} (|u_t(0, \cdot)| + |v_t(0, \cdot)/\alpha|) + R_3 \iint_{Q_T} (|u_t| + |v_t/\alpha|). \end{aligned}$$

Whence we get (4.14) from the Gronwall's inequality.  $\square$

**Lemma 4.2.** *If the nonnegative functions  $u_0$  and  $v_0$  satisfy (2.15) and (2.16), then there exist  $\{k_i\}_{i=1}^{\infty} \subset \mathbb{R}_+$ ,  $\{u_0^{(i)}\}_{i=1}^{\infty} \subset \hat{C}^3(\bar{\Omega})$  and  $\{v_0^{(i)}\}_{i=1}^{\infty} \subset \hat{C}^3(\bar{\Omega})$  such that*

$$k_i \rightarrow \infty \quad \text{as } i \rightarrow \infty, \quad (4.15)$$

$$0 \leq u_0^{(i)} \leq R_1, \quad 0 \leq v_0^{(i)} \leq R_1 \quad (i = 1, 2, \dots), \quad (4.16)$$

$$u_0^{(i)} \rightarrow u_0, \quad v_0^{(i)} \rightarrow v_0 \quad \text{in } H^1(\Omega) \quad \text{as } i \rightarrow \infty, \quad (4.17)$$

$$\int_{\Omega} |d_1 \Delta u_0^{(i)}| + \int_{\Omega} |d_2 \Delta v_0^{(i)}/\alpha| \leq C_2 \quad (i = 1, 2, \dots), \quad (4.18)$$

$$k_i \int_{\Omega} u_0^{(i)} v_0^{(i)} \leq C_3 \quad (i = 1, 2, \dots), \quad (4.19)$$

where  $C_2, C_3$  are independent of  $i$ .

*Proof.* We use the heat equation with the homogeneous Neumann boundary condition as a mollifier. From the standard regularity argument (see Brezis (1983) for example), we observe that

$$\begin{aligned} e^{t\Delta} u_0, e^{t\Delta} v_0 &\in \hat{C}^3(\bar{\Omega}) \quad \text{for } t > 0, \\ e^{t\Delta} u_0 &\rightarrow u_0, \quad e^{t\Delta} v_0 \rightarrow v_0 \quad \text{in } H^2(\Omega) \quad \text{as } t \downarrow 0. \end{aligned}$$

Moreover we have  $0 \leq e^{t\Delta} u_0 \leq \|u_0\|_{L^\infty(\Omega)}$ ,  $0 \leq e^{t\Delta} v_0 \leq \|v_0\|_{L^\infty(\Omega)}$ . Then we set

$$u_0^{(i)} = e^{t\Delta} u_0 \Big|_{t=1/i}, \quad v_0^{(i)} = e^{t\Delta} v_0 \Big|_{t=1/i}.$$

Since

$$\int_{\Omega} u_0^{(i)} v_0^{(i)} \rightarrow \int_{\Omega} u_0 v_0 = 0,$$

we have

$$k_i \stackrel{\text{def}}{=} \frac{1}{1/i + \int_{\Omega} u_0^{(i)} v_0^{(i)}} \rightarrow \infty \quad \text{as } i \rightarrow \infty.$$

Hence we prove the statement.  $\square$

**Lemma 4.3.** *If the conditions (4.15)–(4.19) hold, then solutions  $u^{(k_i)}$  and  $v^{(k_i)}$  to (1.7)–(1.9) for  $k = k_i$ ,  $u^{(k_i)}(0, \cdot) = u_0^{(i)}$  and  $v^{(k_i)}(0, \cdot) = v_0^{(i)}$  satisfy*

$$\sup_{0 \leq t \leq T} \int_{\Omega} (|u_t^{(k_i)}| + |v_t^{(k_i)}/\alpha|) \leq C_4, \quad (4.20)$$

$$\sup_{0 \leq t \leq T} k_i \int_{\Omega} u^{(k_i)} v^{(k_i)} \leq C_5, \quad (4.21)$$

$$\sup_{0 \leq t \leq T} \left( d_1 \int_{\Omega} |\nabla u^{(k_i)}|^2 + d_2 \int_{\Omega} |\nabla v^{(k_i)}/\alpha|^2 \right) \leq C_6, \quad (4.22)$$

$$\sup_{0 \leq t \leq T} \left( d_1 \int_{\Omega} |\Delta u^{(k_i)}| + d_2 \int_{\Omega} |\Delta v^{(k_i)}/\alpha| \right) \leq C_7, \quad (4.23)$$

where  $C_5, C_6$  and  $C_7$  are positive constants independent of  $k_i$ .

*Proof.* From Lemmas 4.1 and 4.2 we obtain (4.20) by putting  $C_4 = e^{R_3 T}(C_2 + (1 + \alpha^{-1})R_2 + 2C_3)$ . Thus we have

$$k_i \int_{\Omega} u^{(k_i)} v^{(k_i)} = - \int_{\Omega} u_t^{(k_i)} + \int_{\Omega} F(u^{(k_i)}) \leq C_5,$$

where  $C_5 = C_4 + R_2|\Omega|$ .

Next, multiplying the both hand sides of (1.7) by  $u^{(k_i)}$  and integrating them over  $\Omega$ , we observe

$$\begin{aligned} d_1 \int_{\Omega} |\nabla u^{(k_i)}|^2 &= - \int_{\Omega} u^{(k_i)} u_t^{(k_i)} + \int_{\Omega} F(u^{(k_i)}) u^{(k_i)} - k_i \int_{\Omega} (u^{(k_i)})^2 v^{(k_i)} \\ &\leq R_1 C_4 + R_1 R_2 |\Omega|. \end{aligned}$$

Note that  $\|u^{(k_i)}\|_{L^\infty(\Omega)} \leq R_1$  from the comparison theorem. A similar estimate holds for  $d_2 \int_{\Omega} |\nabla v^{(k_i)}|^2$ . Setting  $C_6 = (1 + \alpha^{-1})R_1 C_4 + (1 + \alpha^{-2})R_1 R_2 |\Omega|$  provides (4.22).

Finally we see

$$\begin{aligned} d_1 \int_{\Omega} |\Delta u^{(k_i)}| &= \int_{\Omega} |u_t^{(k_i)} - F(u^{(k_i)}) + k_i u^{(k_i)} v^{(k_i)}| \\ &\leq C_4 + R_2 |\Omega| + C_5. \end{aligned}$$

We obtain the estimate for  $d_2 \int_{\Omega} |\Delta v^{(k_i)}|$  likewise. Putting  $C_7 = 2C_4 + (1 + \alpha^{-1})R_2 |\Omega| + 2C_5$  we prove (4.23).  $\square$

**Proposition 4.3.** *Suppose the same conditions as those in Lemma 4.3 hold. Let  $w$  be the weak solution to (1.1)–(1.3). Then  $w$  satisfies the followings:*

$$u^{(k_i)} \rightarrow [w]^+ \quad \text{in } C([0, T]; L^2(\Omega)), \quad (4.24)$$

$$v^{(k_i)}/\alpha \rightarrow [w]^- \quad \text{in } C([0, T]; L^2(\Omega)), \quad (4.25)$$

$$w^{(k_i)} = u^{(k_i)} - v^{(k_i)}/\alpha \rightarrow w \quad \text{in } C([0, T]; L^2(\Omega)). \quad (4.26)$$

*Proof.* We see for  $t \geq s$

$$\begin{aligned} &\int_{\Omega} \left| u^{(k_i)}(t, \cdot) - u^{(k_i)}(s, \cdot) \right|^2 \\ &= \int_{\Omega} \left\{ (u^{(k_i)}(t, x) - u^{(k_i)}(s, x)) \int_s^t u_t^{(k_i)}(\sigma, x) d\sigma \right\} dx \\ &\leq 2R_1 \int_s^t \left( \int_{\Omega} |u_t^{(k_i)}(\sigma, x)| dx \right) d\sigma \\ &\leq 2R_1 C_4 (t - s). \end{aligned} \quad (4.27)$$

Hence  $u^{(k_i)}$  is uniformly bounded in  $C^{1/2}([0, T]; L^2(\Omega))$  and so is  $v^{(k_i)}$ . Therefore  $u^{(k_i)}$  and  $v^{(k_i)}$  are equicontinuous in  $C([0, T]; L^2(\Omega))$ . Moreover from (4.22)  $u^{(k_i)}(t, \cdot)$  and  $v^{(k_i)}(t, \cdot)$  are contained in a compact subset of  $L^2(\Omega)$ . By Ascoli-Arzelà's theorem we obtain a subsequence converging in  $C([0, T]; L^2(\Omega))$ . In view of Proposition 4.1 the limit of  $w^{(k_i)}$  along the subsequence is the weak solution. Uniqueness of the limit assures the convergence of the original sequence.  $\square$

From this proposition we immediately have the next lemma.

**Lemma 4.4.** For any positive  $\varepsilon$ , there exist  $k = k(\varepsilon) > 0$  and  $u_0^{(\varepsilon)}, v_0^{(\varepsilon)} \in \hat{C}^3(\bar{\Omega})$  such that solutions  $u^{(k(\varepsilon))}, v^{(k(\varepsilon))}$  to (1.7)–(1.9) with the initial conditions  $u^{(k(\varepsilon))}(0, \cdot) = u_0^{(\varepsilon)}$  and  $v^{(k(\varepsilon))} = v_0^{(\varepsilon)}$  satisfy the following:

$$\|w^{(k(\varepsilon))} - w\|_{C([0,T];L^p(\Omega))} \leq \varepsilon, \quad (4.28)$$

$$\|u^{(k(\varepsilon))} - [w]^+\|_{C([0,T];L^p(\Omega))} \leq \varepsilon, \quad (4.29)$$

$$\|v^{(k(\varepsilon))}/\alpha - [w]^-\|_{C([0,T];L^p(\Omega))} \leq \varepsilon, \quad (4.30)$$

where  $w^{(k(\varepsilon))} = u^{(k(\varepsilon))} - v^{(k(\varepsilon))}/\alpha$  and  $p = 1, 2$ . Moreover  $k(\varepsilon)$ ,  $u^{(k(\varepsilon))}$  and  $v^{(k(\varepsilon))}$  satisfy the inequalities (4.20)–(4.23).

**Lemma 4.5.** Consider the following equations in each interval  $[t_j, t_{j+1}]$ :

$$\frac{\partial \bar{u}_M^{j,\varepsilon}}{\partial t} = d_1 \Delta \bar{u}_M^{j,\varepsilon} + F(\bar{u}_M^{j,\varepsilon}), \quad t_j < t \leq t_{j+1}, \quad x \in \Omega, \quad (4.31)$$

$$\frac{\partial \bar{v}_M^{j,\varepsilon}}{\partial t} = d_2 \Delta \bar{v}_M^{j,\varepsilon} + G(\bar{v}_M^{j,\varepsilon}), \quad t_j < t \leq t_{j+1}, \quad x \in \Omega, \quad (4.32)$$

$$\frac{\partial \bar{u}_M^{j,\varepsilon}}{\partial \nu} = \frac{\partial \bar{v}_M^{j,\varepsilon}}{\partial \nu} = 0, \quad t_j < t \leq t_{j+1}, \quad x \in \partial\Omega, \quad (4.33)$$

$$\bar{u}_M^{j,\varepsilon}(t_j, x) = u^{(k(\varepsilon))}(t_j, x), \quad (4.34)$$

$$\bar{v}_M^{j,\varepsilon}(t_j, x) = v^{(k(\varepsilon))}(t_j, x). \quad (4.35)$$

Set

$$\bar{w}_M^{j,\varepsilon} = \bar{u}_M^{j,\varepsilon} - \bar{v}_M^{j,\varepsilon}/\alpha. \quad (4.36)$$

If  $d_1 = d_2$ , the following inequality holds:

$$\|w_M(t_{j+1}, \cdot) - \bar{w}_M^{j,\varepsilon}(t_{j+1}, \cdot)\|_{L^p(\Omega)} \leq (e_j^{(p)} + \varepsilon)(1 + E\tau), \quad (p = 1, 2), \quad (4.37)$$

where  $E$  is independent of  $M$ ,  $j$  and  $\varepsilon$ .

*Proof.* Throughout this proof we assume  $p = 1, 2$ . Using the Duhamel formula we have

$$\begin{aligned} & \|w_M(t_{j+1}, \cdot) - \bar{w}_M^{j,\varepsilon}(t_{j+1}, \cdot)\|_{L^p(\Omega)} \\ & \leq \|e^{(t_{j+1}-t_j)d_1\Delta}(w_M(t_j, \cdot) - \bar{w}_M^{j,\varepsilon}(t_j, \cdot))\|_{L^p(\Omega)} \\ & \quad + \left\| \int_{t_j}^{t_{j+1}} e^{(t_{j+1}-s)d_1\Delta} (F(\bar{w}_M^j(s, \cdot)) - F(\bar{u}_M^{j,\varepsilon}(s, \cdot))) ds \right\|_{L^p(\Omega)} \\ & \quad + \frac{1}{\alpha} \left\| \int_{t_j}^{t_{j+1}} e^{(t_{j+1}-s)d_1\Delta} (G(\bar{v}_M^j(s, \cdot)) - G(\bar{v}_M^{j,\varepsilon}(s, \cdot))) ds \right\|_{L^p(\Omega)} \\ & = I + II + III. \end{aligned}$$

Let  $w$  be the weak solution to (1.1)–(1.3). Recall that  $\|e^{td_1\Delta}z\|_{L^p(\Omega)} \leq \|z\|_{L^p(\Omega)}$  for  $z \in L^p(\Omega)$  ( $p = 1, 2$ ). Then we observe

$$\begin{aligned} I & \leq \|w_M(t_j, \cdot) - \bar{w}_M^{j,\varepsilon}(t_j, \cdot)\|_{L^p(\Omega)} \\ & \leq \|w_M(t_j, \cdot) - w(t_j, \cdot)\|_{L^p(\Omega)} + \|w(t_j, \cdot) - \bar{w}_M^{j,\varepsilon}(t_j, \cdot)\|_{L^p(\Omega)} \\ & \leq e_j^{(p)} + \varepsilon. \end{aligned}$$

Next we estimate *II*. For  $t_j < t \leq t_{j+1}$  by means of the Duhamel formula we obtain

$$\begin{aligned}
& \|\bar{w}_M^j(t, \cdot) - \bar{u}_M^{j,\varepsilon}(t, \cdot)\|_{L^p(\Omega)} \\
& \leq \|\bar{w}_M^j(t_j, \cdot) - \bar{u}_M^{j,\varepsilon}(t_j, \cdot)\|_{L^p(\Omega)} + \int_{t_j}^t \|F(\bar{w}_M^j(s, \cdot)) - F(\bar{u}_M^{j,\varepsilon}(s, \cdot))\|_{L^p(\Omega)} ds \\
& \leq \|u_M^j(\cdot) - [w(t_j, \cdot)]^+\|_{L^p(\Omega)} + \|[w(t_j, \cdot)]^+ - \bar{u}_M^{j,\varepsilon}(t_j, \cdot)\|_{L^p(\Omega)} \\
& \quad + \int_{t_j}^t \|F(\bar{w}_M^j(s, \cdot)) - F(\bar{u}_M^{j,\varepsilon}(s, \cdot))\|_{L^p(\Omega)} ds \\
& \leq e_j^{(p)} + \varepsilon + R_3 \int_{t_j}^t \|\bar{w}_M^j(s, \cdot) - \bar{u}_M^{j,\varepsilon}(s, \cdot)\|_{L^p(\Omega)} ds.
\end{aligned}$$

Recall that  $\bar{u}_M^j$  and  $\bar{v}_M^j$  are defined by (2.5)–(2.6). Hence we obtain by Gronwall's inequality

$$\|\bar{w}_M^j(t, \cdot) - \bar{u}_M^{j,\varepsilon}(t, \cdot)\|_{L^p(\Omega)} \leq (e_j^{(p)} + \varepsilon)e^{R_3(t-t_j)}.$$

Therefore

$$II \leq R_3 \int_{t_j}^{t_{j+1}} \|\bar{w}_M^j(s, \cdot) - \bar{u}_M^{j,\varepsilon}(s, \cdot)\|_{L^p(\Omega)} ds \leq R_3(e_j^{(p)} + \varepsilon)e^{R_3T} \tau.$$

In a similar fashion we have

$$III \leq (R_3/\alpha)(e_j^{(p)} + \varepsilon)e^{R_3T} \tau.$$

Setting  $E = (1 + 1/\alpha)R_3e^{R_3T}$  we complete the proof.  $\square$

**Lemma 4.6.** *Suppose  $\bar{w}_M^{j,\varepsilon}$  is given by (4.36). If  $d_1 = d_2$ ,  $\bar{w}_M^{j,\varepsilon}$  satisfies*

$$\|\bar{w}_M^{j,\varepsilon}(t_{j+1}, \cdot) - w^{(k(\varepsilon))}(t_{j+1}, \cdot)\|_{L^2(\Omega)} \leq C_8 \tau^{3/2}, \quad (4.38)$$

$$\|\bar{w}_M^{j,\varepsilon}(t_{j+1}, \cdot) - w^{(k(\varepsilon))}(t_{j+1}, \cdot)\|_{L^1(\Omega)} \leq C_9 \tau^2, \quad (4.39)$$

where  $C_8$  and  $C_9$  are independent of  $M$ ,  $j$  and  $\varepsilon$ .

*Proof.* Recall the inequalities (4.21)–(4.23). Then from an argument similar to that in Lemma 4.1 we obtain

$$\sup_{t_j \leq t \leq t_{j+1}} \left( \int_{\Omega} |\partial_t \bar{w}_M^{j,\varepsilon}(t, \cdot)| + \int_{\Omega} |\partial_t \bar{v}_M^{j,\varepsilon}(t, \cdot)/\alpha| \right) \leq C_{10},$$

where  $C_{10}$  is independent of  $\varepsilon$ ,  $j$ ,  $M$ . Noting that  $u^{(k(\varepsilon))}(t_j, \cdot) = \bar{u}_M^{j,\varepsilon}(t_j, \cdot)$  we have

$$\begin{aligned}
& \int_{\Omega} |u^{(k(\varepsilon))}(s, \cdot) - \bar{u}_M^{j,\varepsilon}(s, \cdot)|^2 \\
& = \int_{\Omega} \left\{ (u^{(k(\varepsilon))}(s, x) - \bar{u}_M^{j,\varepsilon}(s, x)) \int_{t_j}^s (\partial_t u^{(k(\varepsilon))}(\sigma, x) - \partial_t \bar{u}_M^{j,\varepsilon}(\sigma, x)) d\sigma \right\} dx \\
& \leq 2R_1 \int_{t_j}^s \int_{\Omega} (|\partial_t u^{(k(\varepsilon))}(\sigma, x)| + |\partial_t \bar{u}_M^{j,\varepsilon}(\sigma, x)|) dx d\sigma \\
& \leq 2R_1(C_4 + C_{10})(s - t_j).
\end{aligned}$$

Thus we get

$$\|(u^{(k(\varepsilon))}(s, \cdot) - \bar{u}_M^{j,\varepsilon}(s, \cdot))\|_{L^2(\Omega)} \leq \sqrt{2R_1(C_4 + C_{10})}(s - t_j)^{1/2}.$$

Likewise we observe

$$\begin{aligned} \|u^{(k(\varepsilon))}(s, \cdot) - \bar{u}_M^{j, \varepsilon}(s, \cdot)\|_{L^1(\Omega)} &\leq (C_4 + C_{10})(s - t_j), \\ \|(v^{(k(\varepsilon))}(s, \cdot) - \bar{v}_M^{j, \varepsilon}(s, \cdot))/\alpha\|_{L^2(\Omega)} &\leq \sqrt{2(R_1/\alpha)(C_4 + C_{10})(s - t_j)^{1/2}}, \\ \|(v^{(k(\varepsilon))}(s, \cdot) - \bar{v}_M^{j, \varepsilon}(s, \cdot))/\alpha\|_{L^1(\Omega)} &\leq (C_4 + C_{10})(s - t_j). \end{aligned}$$

Here we use the Duhamel formula again. The condition  $d_1 = d_2$  makes the term  $kuv$  vanish so that we have

$$\begin{aligned} &\|w^{(k(\varepsilon))}(t_{j+1}, \cdot) - \bar{w}_M^{j, \varepsilon}(t_{j+1}, \cdot)\|_{L^2(\Omega)} \\ &\leq \int_{t_j}^{t_{j+1}} \|F(u^{(k(\varepsilon))}(s, \cdot)) - F(\bar{u}_M^{j, \varepsilon}(s, \cdot))\|_{L^2(\Omega)} ds \\ &\quad + \frac{1}{\alpha} \int_{t_j}^{t_{j+1}} \|G(v^{(k(\varepsilon))}(s, \cdot)) - G(\bar{v}_M^{j, \varepsilon}(s, \cdot))\|_{L^2(\Omega)} ds \\ &\leq R_3 \int_{t_j}^{t_{j+1}} \|u^{(k(\varepsilon))}(s, \cdot) - \bar{u}_M^{j, \varepsilon}(s, \cdot)\|_{L^2(\Omega)} ds \\ &\quad + \frac{R_3}{\alpha} \int_{t_j}^{t_{j+1}} \|v^{(k(\varepsilon))}(s, \cdot) - \bar{v}_M^{j, \varepsilon}(s, \cdot)\|_{L^2(\Omega)} ds \\ &\leq \frac{2}{3}(1 + \alpha^{-1/2})R_3(\sqrt{2R_1(C_4 + C_{10})})\tau^{3/2}. \end{aligned}$$

Analogously we get

$$\|w^{(k(\varepsilon))}(t_{j+1}, \cdot) - \bar{w}_M^{j, \varepsilon}(t_{j+1}, \cdot)\|_{L^1(\Omega)} \leq R_3(C_4 + C_{10})\tau^2,$$

which completes the proof.  $\square$

Now we are ready to prove Theorem 2.2.

*Proof of Theorem 2.2.* From (4.28), (4.37) and (4.38) we observe

$$\begin{aligned} e_{j+1}^{(2)} &= \|w_M(t_{j+1}, \cdot) - w(t_{j+1}, \cdot)\|_{L^2(\Omega)} \\ &\leq \|w_M(t_{j+1}, \cdot) - \bar{w}_M^{j, \varepsilon}(t_{j+1}, \cdot)\|_{L^2(\Omega)} \\ &\quad + \|\bar{w}_M^{j, \varepsilon}(t_{j+1}, \cdot) - w^{(k(\varepsilon))}\|_{L^2(\Omega)} \\ &\quad + \|w^{(k(\varepsilon))}(t_{j+1}, \cdot) - w(t_{j+1}, \cdot)\|_{L^2(\Omega)} \\ &\leq (e_j^{(2)} + \varepsilon)(1 + E\tau) + C_8\tau^{3/2} + \varepsilon. \end{aligned}$$

Recall that we can choose arbitrary small  $\varepsilon > 0$  for the above inequality. Whence we have

$$e_{j+1}^{(2)} \leq (1 + E\tau)e_j^{(2)} + C_8\tau^{3/2}.$$

Consequently we are led to

$$e_M^{(2)} \leq \frac{C_8\tau^{1/2}}{E}((1 + E\tau)^M - 1) \tag{4.40}$$

$$\leq \frac{C_8}{E}(T/M)^{1/2}(e^{ET} - 1) \equiv C(1/M)^{1/2}. \tag{4.41}$$

In a similar way we arrive at

$$e_M^{(1)} \leq C'(1/M),$$

thereby completing the proof.  $\square$



## 5 Numerical Experiments

In this section we perform numerical experiments to observe how TCD converges. This section is composed of two parts. First, we pose a test problem we used in this section. After describing the numerical scheme to TCD, we report numerical results.

### 5.1 A Test Problem

We consider the singular limit of the following system

$$\begin{cases} u_t = d_1 \Delta u + (r_1 - a_1 u)u - kb_1 uv, \\ v_t = d_2 \Delta v + (r_2 - a_2 v)v - kb_2 uv, \end{cases} \quad (x, y) \in \Omega = [0, X] \times [0, Y], 0 < t < T. \quad (5.1)$$

We impose the Neumann boundary condition on  $u$  and  $v$ . The initial functions  $u|_{t=0}$  and  $v|_{t=0}$  are shown in Fig. 1. In this test problem, the parameters in (5.1) are fixed as




Figure 1: Initial functions  $u|_{t=0}$  (left) and  $v|_{t=0}$  (right) used in our computations.

$$\begin{aligned} d_1 &= 0.7, & r_1 &= 2.5, & a_1 &= 1, & b_1 &= 1.2 \\ d_2 &= 1.5, & r_2 &= 1, & a_2 &= 1, & b_2 &= 3.5, \\ k &= 10^{10}, & X &= 2, & Y &= 2, & T &= 0.1. \end{aligned}$$

Since no exact solutions of (5.1) are available, the numerical solution on the uniform  $256 \times 256$  mesh and the time increment  $2 \times 10^{-7}$  will be referred as an “exact” solution. By our careful numerical computations we conclude that the value  $k = 10^{10}$  is large enough to regard the numerical result as the singular limit solution of (5.1). The “exact” solution, which is shown in Fig. 2, is computed by the standard finite difference scheme. The zero-level sets of  $w = u - v/\alpha$




Figure 2: “Exact” solutions  $u|_{t=0.1}$  (left) and  $v|_{t=0.1}$  (right).

is displayed in Fig. 3.




Figure 3: The zero-level sets of  $w = u - v/\alpha$  at  $t = 0$  (left) and  $t = 0.1$  (right).

## 5.2 Numerical Scheme and Results

Let  $M_X$ ,  $M_Y$  and  $M_T$  be positive integers. We denote the spatial mesh sizes by  $\delta x = X/M_X$  and  $\delta y = Y/M_Y$ , the time increment by  $\delta t = T/M_T$ . Let  $u_{i,j}^n$  ( $0 \leq i \leq M_X$ ,  $0 \leq j \leq M_Y$  and  $0 \leq n \leq M_T$ ) be an approximation to  $u(x, y, t)$  at the location  $(x, y) = (i\delta x, j\delta y)$  and the time  $t = \delta t$ . We define operators  $L$  by

$$Lu_{i,j}^n = \frac{u_{i+1,j}^n - 2u_{i,j}^n + u_{i-1,j}^n}{\delta x^2} + \frac{u_{i,j+1}^n - 2u_{i,j}^n + u_{i,j-1}^n}{\delta y^2}. \quad (5.2)$$

We introduce an implementation of TCD. For Step 1 of TCD, we use the following semi-implicit scheme:

$$\begin{cases} \frac{u_{i,j}^{n+1/2} - u_{i,j}^n}{\delta t} = d_1 Lu_{i,j}^{n+1/2} + (r_1 - a_1 u_{i,j}^{n+1/2}) u_{i,j}^n, \\ \frac{v_{i,j}^{n+1/2} - v_{i,j}^n}{\delta t} = d_2 Lv_{i,j}^{n+1/2} + (r_2 - a_2 v_{i,j}^{n+1/2}) v_{i,j}^n \end{cases} \quad (5.3)$$

for  $0 \leq i \leq M_X$ ,  $0 \leq j \leq M_Y$ . Here we define

$$\begin{aligned} u_{-1,j}^{n+1/2} &= u_{1,j}^{n+1/2}, & u_{M_X+1,j}^{n+1/2} &= u_{M_X-1,j}^{n+1/2}, & u_{i,-1}^{n+1/2} &= u_{i,1}^{n+1/2}, \\ u_{i,M_Y+1}^{n+1/2} &= u_{i,M_Y-1}^{n+1/2}, & v_{-1,j}^{n+1/2} &= v_{1,j}^{n+1/2}, & v_{M_X+1,j}^{n+1/2} &= v_{M_X-1,j}^{n+1/2}, \\ v_{i,-1}^{n+1/2} &= v_{i,1}^{n+1/2}, & v_{i,M_Y+1}^{n+1/2} &= v_{i,M_Y-1}^{n+1/2}, \end{aligned} \quad (5.4)$$

which are led by the Neumann boundary condition.

Step 2 of TCD can be written by

$$\begin{cases} u_{i,j}^{n+1} = [u_{i,j}^{n+1/2} - v_{i,j}^{n+1/2}/\alpha]^+, \\ v_{i,j}^{n+1} = \alpha [u_{i,j}^{n+1/2} - v_{i,j}^{n+1/2}/\alpha]^-, \end{cases} \quad (5.5)$$

where  $a = b_2/b_1$ .

Now we report the results of numerical simulations by the above implementation of TCD. We choose  $M_X = M_Y = 128$  and  $M_T = 1600, 3200, 6400, \dots, 51200$ . Fig. 4 shows the relative errors  $E_u$  and  $E_v$  at  $t = 0.1$  defined by

$$\begin{aligned} E_u &= \max_{(i,j) \in \Lambda} |u_{i,j}^{M_T} - u(i\delta x, j\delta y, 0.1)| / \max_{(i,j) \in \Lambda} |u(i\delta x, j\delta y, 0.1)|, \\ E_v &= \max_{(i,j) \in \Lambda} |v_{i,j}^{M_T} - v(i\delta x, j\delta y, 0.1)| / \max_{(i,j) \in \Lambda} |v(i\delta x, j\delta y, 0.1)|, \end{aligned} \quad (5.6)$$

where  $(u(x, y, t), v(x, y, t))$  is the “exact” solution and  $\Lambda = \{(i, j) \in \mathbb{Z}^2 \mid 0 \leq i \leq M_X, 0 \leq j \leq M_Y\}$ . One can observe that  $E_u$  and  $E_v$  tends to zero as  $\delta t \rightarrow 0$ .




Figure 4: The relative errors versus the time increment.

## Acknowledgments

The authors thank Professors Masato Iida of Iwate University, Hideo Ikeda of Toyama University and Masato Kimura of Kyushu University for stimulus discussions and Professor Danielle Hilhorst for introducing the finite volume method to them. They also express their gratitude to Professor Hisashi Inaba for his encouragement.

The first author was supported by Research Fellowships of the Japan Society for the Promotion of Science for Young Scientists

## A Notation Index

$T$  : a time interval.

$M$  : a time step.

$\tau$  : a time mesh size;  $\tau = T/M$ .

$\bar{u}_M^j, \bar{v}_M^j$  : auxiliary functions associated with Step 1 of TCD; defined by (2.5), (2.6).

$\hat{u}_M^j, \hat{v}_M^j$  : auxiliary functions associated with Step 2 of TCD; see (2.8).

$u_M, v_M$  : approximate functions to the singular limit via TCD; see (2.3).

$w_M$  : an approximate function to (1.1)–(1.3) via TCD;  $w_M = u_M - v_M/\alpha$ .

$u_M^j, v_M^j$  : approximate functions to the singular limit via TCD at the time  $t = t_j$ ; see (2.7), (2.11) and (2.12).

$\bar{\bar{u}}_M^{j,\varepsilon}, \bar{\bar{v}}_M^{j,\varepsilon}$  : auxiliary functions; see (4.31)–(4.35)

## References

- Barles, G. & Georgelin, C. (1995), ‘A simple proof of convergence for an approximation scheme for computing motions by mean curvature’, *SIAM J. Numer. Anal.* **32**(2), 484–500.
- Brezis, H. (1983), *Analyse fonctionnelle*, Masson.
- Cantrell, R. S. (1996), ‘Antibifurcation and the  $n$ -species lotka-volterra competition model with diffusion’, *Differential Integral Equations* **9**(2), 305–322.
- Cosner, C. & Lazer, A. C. (1984), ‘Stable coexistence states in the volterra-lotka competition model with diffusion’, *SIAM J. Appl. Math.* **44**(6), 1112–1132.
- Dancer, E. N., Hilhorst, D., Mimura, M. & Peletier, L. A. (1999), ‘Spatial segregation limit of a competition-diffusion system’, *European J. Appl. Math.* **10**(2).
- Ei, S.-I., Ikota, R. & Mimura, M. (1999), ‘Segregating partition problem in competition-diffusion systems’, *Interfaces and Free Boundaries* **1**(1), 57–80.
- Evans, L. C. (1980), ‘A convergence theorem for a chemical diffusion-reaction system’, *Houston J. Math.* **6**(2), 259–267.
- Evans, L. C. (1993), ‘Convergence of an algorithm for mean curvature motion’, *Indiana Univ. Math. J.* **42**(2), 533–557.
- Eymard, R., Gallouët, T., Hilhorst, D. & Slimane, Y. N. (1998), ‘Finite volumes and nonlinear diffusion equations’, *RAIRO Modél. Math. Anal. Numér.* **32**(6), 747–761.
- Iida, M., Muramatsu, T., Ninomiya, H. & Yanagida, E. (1998), ‘Diffusion-induced extinction of a superior species in a competition system’, *Japan J. Indust. Appl. Math.* **15**(2), 233–252.
- Ikota, R., Mimura, M. & Nakaki, T. (2001), Numerical computation for some competition-diffusion systems on a parallel computer, in T. Chan, T. Kako, H. Kawarada & O. Pironneau, eds, ‘Proceedings of 12th international conference on domain decomposition methods’, DDM.org, pp. 373–379.
- Ishii, H., Pires, G. E. & Souganidis, P. E. (1999), ‘Threshold dynamics type approximation schemes for propagating fronts’, *J. Math. Soc. Japan* **51**(2), 267–308.
- Merriman, B., Bence, J. K. & Osher, S. J. (1994), ‘Motion of multiple junctions: a level set approach’, *J. Comput. Phys.* **112**, 334–363.
- Mimura, M. & Fife, P. C. (1986), ‘A 3-component system of competition and diffusion’, *Hiroshima Math. J.* **16**, 189–207.
- Mora, X. (1983), ‘Semilinear parabolic problems define semiflows on  $c^k$  spaces’, *Trans. Amer. Math. Soc.* **278**(1), 21–55.
- Ruuth, S. J. (1998), ‘A diffusion-generated approach to multiphase motion’, *J. Comput. Phys.* **145**, 166–192.
- Temam, R. (1984), *Navier-Stokes Equations*, third edition edn, North-Holland.
- Tonegawa, Y. (1998), ‘On the regularity of a chemical reaction interface’, *Comm. Partial Differential Equations* **23**(7–8), 1181–1207.
- Zeidler, E. (1990), *Nonlinear Functional Analysis and its Applications*, Vol. II/A, Springer-Verlag.

# List of MHF Preprint Series, Kyushu University

## 21st Century COE Program

### Development of Dynamic Mathematics with High Functionality

#### MHF

- 2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE  
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- 2003-2 Masahisa TABATA & Daisuke TAGAMI  
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- 2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO  
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- 2003-4 Kazuhiro YOKOYAMA  
On systems of algebraic equations with parametric exponents
- 2003-5 Masao ISHIKAWA & Masato WAKAYAMA  
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- 2003-6 Atsushi SUZUKI & Masahisa TABATA  
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- 2003-7 Setsuo TANIGUCHI  
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- 2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA  
Computable sequences in the Sobolev spaces
- 2003-9 Toru FUJII & Takashi YANAGAWA  
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- 2003-10 Atsushi YOSHIKAWA  
Maple and wave-front tracking — an experiment
- 2003-11 Masanobu KANEKO  
On the local factor of the zeta function of quadratic orders
- 2003-12 Hidefumi KAWASAKI  
Conjugate-set game for a nonlinear programming problem

- 2004-1 Koji YONEMOTO & Takashi YANAGAWA  
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- 2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI  
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- 2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA  
Cubic pencils and Painlevé Hamiltonians
- 2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA  
Estimating the correlation dimension from a chaotic system with dynamic noise
- 2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO  
Detection of auroral breakups using the correlation dimension
- 2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI  
A methodology for numerical simulations to a singular limit