

Finite Element Schemes for Two-Fluids Flow Problems

Tabata, Masahisa
Faculty of Mathematics, Kyushu University

Kaizu, Satoshi
Faculty of Education, Ibaraki University

<https://hdl.handle.net/2324/3371>

出版情報 : Proceedings of the Seventh China-Japan Seminar on Numerical Mathematics, pp.139-148, 2006

バージョン :

権利関係 :

MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

Finite element schemes for two-fluids flow problems

M. Tabata & S. Kaizu

MHF 2005-23

(Received June 17, 2005)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Finite Element Schemes for Two-Fluids Flow Problems

Masahisa TABATA ^a and Satoshi KAIZU ^b

^a *Faculty of Mathematics, Kyushu University*

Hakozaki, Fukuoka, 812-8581 JAPAN

E-mail: tabata@math.kyushu-u.ac.jp

^b *Faculty of Education, Ibaraki University*

Bunkyo, Mito, 310-8512 Japan

E-mail: kaizu@mx.ibaraki.ac.jp

A weak formulation suitable for two-fluids flow problems is shown. Stability of the velocity is proved for a wide class of finite element spaces under a mild assumption on the density. Full stability is proved for a mass-conservative upwind scheme without any condition.

Keywords: two-fluids flow, density-dependent Navier-Stokes equations, finite element method, mass-conservative upwind scheme

AMS Subject classification: Primary 65M12, 65M60; Secondary 76M10

1. Introduction

Two-fluids flow problems describe many interesting phenomena such as Rayleigh-Taylor instability and sloshing of fluid, and various computations have been done by many authors, e.g., [5], [4]. The problems include unknown interfaces, that is, the domains where the problems are solved are also unknown. To the best of our knowledge there are no numerical schemes whose solutions are proved to converge to the exact one. Recently we have developed a finite element scheme for density-dependent Navier-Stokes equations and proved the convergence of the finite element solutions [3]. In this paper we consider two-fluids flow problems in the framework of density-dependent Navier-Stokes problems. We present a weak formulation suitable for free-interface problems. In this formulation a wide class of finite element methods is stable with respect to the velocity under the non-negativity assumption on the density. The assumption is satisfied by a mass-conservative upwind finite element approximation developed by Baba-Tabata [1]. The finite element scheme is proved unconditionally stable for all variables. The study to show the convergence of the finite element solutions is ongoing.

2. Two-layers flow problems

Let Ω be a bounded domain in \mathbf{R}^d , $d = 2, 3$, with piecewise smooth boundary Γ , and T be a positive number. At the initial time $t = 0$ the domain Ω is occupied by two immiscible incompressible viscous fluids; each domain is denoted by Ω_k^0 , $k = 1, 2$, whose interface $\partial\Omega_1^0 \cap \partial\Omega_2^0$ is denoted by Γ_{12}^0 . At $t \in (0, T)$ the two fluids occupy domains $\Omega_k(t)$, $k = 1, 2$, and the interface $\partial\Omega_1(t) \cap \partial\Omega_2(t)$ is denoted by $\Gamma_{12}(t)$.

Let ρ_k and μ_k , $k = 1, 2$, be the densities and the viscosities of the two fluids. Let

$$u : \Omega \times (0, T) \rightarrow \mathbf{R}^d, \quad p : \Omega \times (0, T) \rightarrow \mathbf{R}$$

be the velocity and the pressure to be found. The Navier-Stokes equations are satisfied in each domain $\Omega_k(t)$, $k = 1, 2$, $t \in (0, T)$,

$$\rho_k \left\{ \frac{\partial u}{\partial t} + (u \cdot \nabla)u \right\} - \nabla (2\mu_k D(u)) + \nabla p = \rho_k f \quad (1a)$$

$$\nabla \cdot u = 0 \quad (1b)$$

where $f : \Omega \times (0, T) \rightarrow \mathbf{R}^d$ is a given function, $D(u)$ is the strain tensor defined by

$$D_{ij}(u) = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right)$$

and

$$[\nabla (2\mu_k D(u))]_i = \sum_{j=1}^d \frac{\partial}{\partial x_j} (2\mu_k D_{ij}(u)).$$

On the boundary Γ , $t \in (0, T)$, the non-slip conditions

$$u = 0 \quad (2)$$

are imposed. On the interface $\Gamma_{12}(t)$, $t \in (0, T)$, the velocity and the stress vector should be continuous,

$$[u] = 0 \quad (3a)$$

$$[\sigma(\mu, u, p)n_{12}] = 0 \quad (3b)$$

where $[\cdot]$ means the difference of the values approaching from the domain Ω_2 and the domain Ω_1 , n_{12} is the unit outer normal to $\Gamma_{12}(t)$ from $\Omega_1(t)$ to $\Omega_2(t)$, and σ is the stress tensor defined by

$$\sigma(\mu, u, p) = -pI + 2\mu D(u).$$




Figure 1. Domains $\Omega_k(t)$, $k = 1, 2$, and the interface $\Gamma_{12}(t)$

Let $\rho(x, t)$ be the solution of the convection equation

$$\frac{\partial \rho}{\partial t} + u \cdot \nabla \rho = 0, \quad (x, t) \in \Omega \times (0, T) \quad (4a)$$

$$\rho(x, 0) = \rho^0(x), \quad x \in \Omega \quad (4b)$$

where u is the unknown velocity and

$$\rho^0(x) = \begin{cases} \rho_1 & (x \in \Omega_1^0) \\ \rho_2 & (x \in \Omega_2^0) \end{cases} \quad (5)$$

Note that no boundary condition is required as the velocity u vanishes on the boundary. Then, $\Omega_k(t)$ can be written as

$$\Omega_k(t) = \{x \in \Omega; \rho(x, t) = \rho_k\}, \quad k = 1, 2.$$

Using $\rho(x, t)$, we can express the viscosity $\mu(x, t)$ as the function of ρ ,

$$\mu(\rho) = \mu_1 \frac{\rho_2 - \rho}{\rho_2 - \rho_1} + \mu_2 \frac{\rho - \rho_1}{\rho_2 - \rho_1}. \quad (6)$$

3. A weak formulation

The function ρ^0 is discontinuous and does not belong to $H^1(\Omega)$. We, however, assume that $\rho(\cdot, t) \in H^1(\Omega)$, $t \in (0, T)$, for the time being. Deriving a weak formulation under the assumption, we devise a finite element scheme for the original problem. Apart from (5) and (6) we set a more general setting on $\mu(\rho)$,

$$\mu : \mathbf{R} \rightarrow [\mu_1, \mu_2] \quad (7)$$

where μ_i , $i = 1, 2$, are positive constants and μ is continuously differentiable.

Let $X = H^1(\Omega)$, $Y = X^d$, and V and Q be function spaces defined by

$$V = H_0^1(\Omega)^d, \quad Q = \{q \in L^2(\Omega); \int_{\Omega} q \, dx = 0\}.$$

We consider the variational problem to find the functions

$$\rho : (0, T) \rightarrow X, \quad u : (0, T) \rightarrow V, \quad p : (0, T) \rightarrow Q \quad (8)$$

satisfying

$$\left(\frac{\partial \rho}{\partial t}, \phi \right) + c_1(u, \rho, \phi) = 0, \quad \forall \phi \in X \quad (9a)$$

$$\left(\rho \frac{\partial u}{\partial t} + \frac{1}{2} u \frac{\partial \rho}{\partial t}, v \right) + a_1(\rho, u, u, v) + a_0(\rho, u, v) + b(v, p) = (\rho f, v), \quad \forall v \in V \quad (9b)$$

$$b(u, q) = 0, \quad \forall q \in Q \quad (9c)$$

subject to the initial conditions

$$\rho(0) = \rho^0, \quad u(0) = u^0 \quad (10)$$

where

$$\rho^0 : \Omega \rightarrow \mathbf{R}, \quad u^0 : \Omega \rightarrow \mathbf{R}^d$$

are given functions, (\cdot, \cdot) shows the inner product in $L^2(\Omega)$ or $L^2(\Omega)^d$, and

$$\begin{aligned} c_1(u, \rho, \phi) &= \int_{\Omega} (\nabla \cdot (u\rho)) \phi \, dx \\ a_1(\rho, w, u, v) &= \int_{\Omega} \left(\frac{1}{2}(w \cdot \nabla \rho)u + \frac{1}{2}\rho(\nabla \cdot w)u + \rho(w \cdot \nabla)u \right) \cdot v \, dx \\ a_0(\rho, u, v) &= \int_{\Omega} 2\mu(\rho)D(u) : D(v) \, dx \\ b(v, q) &= - \int_{\Omega} (\nabla \cdot v)q \, dx. \end{aligned}$$

The Sobolev theorem implies the following.

Proposition 1. (i) c_1 is a continuous tri-linear form on $Y \times X \times X$.

(ii) a_1 is a continuous tetra-linear form on $X \times Y \times Y \times Y$.

(iii) a_0 is a continuous tri-linear form on $L^\infty(\Omega) \times Y \times Y$.

(iv) b is a continuous bi-linear form on $Y \times X$.

We now assume that the space-time domain $Q(T) \equiv \Omega \times (0, T)$ is divided into two domains $Q_1(T)$ and $Q_2(T)$ such that the common surface $\Sigma \equiv \bar{Q}_1(T) \cap \bar{Q}_2(T)$ is a smooth surface in \mathbf{R}^{d+1} , and that $\Sigma \cap \{(x, t); x \in \Omega, t = s\}$ is not empty for all $s \in [0, T]$.

Theorem 2. Suppose that (ρ, u, p) is smooth in $Q_1(T)$ and $Q_2(T)$, and continuous in $Q(T)$. If (ρ, u, p) is the solution of the variational problem (8)-(10), then (ρ, u, p) satisfies the equation

$$\rho \left\{ \frac{\partial u}{\partial t} + (u \cdot \nabla)u \right\} - \nabla (2\mu(\rho)D(u)) - \nabla p = \rho f$$

and (1b) in each domain $Q_k(T)$, $k = 1, 2$, and the interface conditions (3) on Σ .

We omit the proof.

Remark 3. In Theorem 2 we assumed that ρ is continuous in $Q(T)$. In (9a), however, the required differentiability of ρ is concerned only with the material differential

$$\frac{D}{Dt} \equiv \frac{\partial}{\partial t} + u \cdot \nabla.$$

Hence, the continuity to the direction $(u, 1)$ in \mathbf{R}^{d+1} is required, but the continuity to the transverse direction is not necessary. If the normal to Σ is orthogonal to $(u, 1)$, ρ can be discontinuous at Σ . It is the case of the two-fluids flow problems.

4. Finite element schemes

Now we consider the finite element approximation to the variational problem (8)-(10). Let $\Phi_h \subset X$, $V_h \subset V$, and $Q_h \subset Q$ be finite element spaces for the density, the velocity, and the pressure. These

spaces are equipped with the norms $L^2(\Omega)$, $H^1(\Omega)^d$, and $L^2(\Omega)$, respectively. L^2 -norm and H^1 -norm are denoted simply by $\|\cdot\|_0$ and $\|\cdot\|_1$. We assume the inf-sup condition on V_h and Q_h

$$\inf_{q_h \in Q_h} \sup_{v_h \in V_h} \frac{b(v_h, q_h)}{\|q_h\|_0 \|v_h\|_1} \geq \beta \quad (11)$$

where β is a positive constant independent of h , the representative mesh size. As the problem (8)-(10) includes the Navier-Stokes equations, the inf-sup condition is necessary when we discuss the convergence of the finite element solutions, though we do not do it in this paper. Let Δt be a time increment and $N_T = \lfloor T/\Delta t \rfloor$. We denote by (ρ_h^n, u_h^n, p_h^n) the value at time $n\Delta t$, and by $\bar{D}_{\Delta t}$ the backward difference operator, e.g.,

$$\bar{D}_{\Delta t} u_h^n = \frac{u_h^n - u_h^{n-1}}{\Delta t}.$$

We find $\{(\rho_h^n, u_h^n, p_h^n) \in \Phi_h \times V_h \times Q_h; n = 1, \dots, N_T\}$ satisfying

$$(\bar{D}_{\Delta t} \rho_h^n, \phi_h) + c_1(u_h^{n-1}, \rho_h^n, \phi_h) = 0, \quad \forall \phi_h \in \Phi_h \quad (12a)$$

$$\left(\rho_h^{n-1} \bar{D}_{\Delta t} u_h^n + \frac{1}{2} u_h^n \bar{D}_{\Delta t} \rho_h^n, v_h \right) + a_1(\rho_h^n, u_h^{n-1}, u_h^n, v_h) + a_0(\rho_h^n, u_h^n, v_h) + b(v_h, p_h^n) = (\rho_h^n \Pi_h f^n, v_h), \quad \forall v_h \in V_h \quad (12b)$$

$$b(u_h^n, q_h) = 0, \quad \forall q_h \in Q_h \quad (12c)$$

subject to the initial conditions

$$\rho_h^0 = \Pi_h \rho^0, \quad u_h^0 = \Pi_h u^0 \quad (13)$$

where Π_h is the interpolation operator to the corresponding finite element spaces.

For a series of functions $\phi_h = \{\phi_h^n\}_{n=0}^{N_T}$ in a Banach space W we prepare the norms defined by

$$\|\phi_h\|_{\ell^\infty(W)} \equiv \max\{\|\phi_h^n\|_W; 0 \leq n \leq N_T\},$$

$$\|\phi_h\|_{\ell^2(W)} \equiv \left\{ \Delta t \sum_{n=0}^{N_T} \|\phi_h^n\|_W^2 \right\}^{1/2}.$$

Lemma 4. Suppose that $(\rho_h^n, u_h^n, p_h^n) \in \Phi_h \times V_h \times Q_h$, $n = 0, \dots, N_T$, satisfy (12b) and (12c) and that $\rho_h \geq 0$. Then we have

$$\|\sqrt{\rho_h} u_h\|_{\ell^\infty(L^2)}, \|\sqrt{\mu_1} u_h\|_{\ell^2(H^1)} \leq c \{ \|\sqrt{\rho_h^0} u_h^0\|_0 + \|\sqrt{\rho_h} \Pi_h f\|_{\ell^2(L^2)} \} \quad (14)$$

where c is a positive constant independent of h and Δt .

Proof. We substitute $v_h = u_h^n$ in (12b). The first term is equal to

$$\begin{aligned} & \left(\rho_h^{n-1} \bar{D}_{\Delta t} u_h^n + \frac{1}{2} u_h^n \bar{D}_{\Delta t} \rho_h^n, u_h^n \right) \\ &= \bar{D}_{\Delta t} \left(\frac{1}{2} \|\sqrt{\rho_h^n} u_h^n\|_0^2 \right) + \frac{1}{2} \|\sqrt{\Delta t} \sqrt{\rho_h^{n-1}} \bar{D}_{\Delta t} u_h^n\|_0^2 \end{aligned}$$

By using the Gauss-Green theorem and the boundary condition, the second term becomes

$$a_1(\rho_h^n, u_h^{n-1}, u_h^n, u_h^n) = \int_{\Gamma} \rho_h^n (u_h^{n-1} \cdot n) |u_h^n|^2 ds = 0.$$

The third term is estimated as

$$a_0(\rho_h^n, u_h^n, u_h^n) \geq \|\sqrt{2\mu_1}D(u_h^n)\|_0^2 \geq c_0\|\sqrt{2\mu_1}u_h^n\|_1^2$$

where c_0 is a positive constant in the Korn inequality. The fourth term vanishes from (11c). The right-hand side is evaluated as

$$|(\rho_h^n \Pi_h f^n, u_h^n)| \leq \epsilon \|\sqrt{\rho_h^n} u_h^n\|_0^2 + \frac{1}{4\epsilon} \|\sqrt{\rho_h^n} \Pi_h f^n\|_0^2$$

where ϵ is any positive constant. Combining these estimates and applying the discrete Gronwall inequality, we get (14). \square

Remark 5. If we replace the tri-linear form $c_1(u, \rho, \phi)$ by a skew symmetric form

$$\hat{c}_1(u, \rho, \phi) = \frac{1}{2} \int_{\Omega} ((u \cdot \nabla \rho) \phi - (u \cdot \nabla \phi) \rho) dx,$$

we can easily get the estimate

$$\|\rho_h\|_{\ell^\infty(L^2)} \leq \|\rho_h^0\|_0$$

by substituting $\phi_h = \rho_h^n$ in (12a). When ρ is continuous, $\hat{c}_1(u, \rho, \phi) = c_1(u, \rho, \phi)$. If not so, \hat{c}_1 includes another condition $[\rho u \cdot n_{12}] = 0$ on the interface.

5. A mass-conservative upwind finite element scheme

We now replace the tri-linear form $c_1(\rho, u, p)$ by a mass-conservative upwind approximation $c_{1h}(u_h, \rho_h, \phi_h)$ [1]. Using the lumping technique, we can show the stability of ρ_h , u_h , and p_h without any assumption. We describe the scheme only in the case $d = 2$. At first we make the dual decomposition




Figure 2. A node P and the barycentric domain D_P

$\{D_P\}$, where D_P is the barycentric domain of the node P , see Fig. 2. The D_P is made by connecting the midpoints of the sides and the centroids of the triangles around $P \in \Omega$. When the node P is on Γ , some parts of the (approximate) boundary are used. Let Φ_h be the P1-finite element space. We define the lumping operator $\bar{\cdot}$ from Φ_h to $L^2(\Omega)$ by

$$\phi_h \mapsto \bar{\phi}_h(x) \equiv \sum_P \phi_h(P) \chi_P(x)$$

where χ_P is the characteristic function of D_P . We denote by Λ_P the set of nodes adjacent to P . Let $u_h \in V_h$ and $\rho_h, \phi_h \in \Phi_h$. The mass-conservative upwind approximation c_{1h} is defined by

$$c_{1h}(u_h, \rho_h, \phi_h) = \sum_P \phi_h(P) \sum_{Q \in \Lambda_P} \left\{ \beta_{PQ}^+(u_h) \rho_h(P) - \beta_{PQ}^-(u_h) \rho_h(Q) \right\}$$

where

$$\beta_{PQ}(u_h) = \int_{\Gamma_{PQ}} u_h \cdot n_{PQ} \, ds, \quad \Gamma_{PQ} = \partial D_P \cap \partial D_Q,$$

n_{PQ} is the unit outer normal to Γ_{PQ} from D_P to D_Q ,

$$\beta^+ = \max(\beta, 0), \quad \beta^- = \max(-\beta, 0).$$

We replace (12a) by

$$(\bar{D}_{\Delta t} \bar{\rho}_h^n, \bar{\phi}_h) + c_{1h}(u_h^{n-1}, \rho_h^n, \phi_h) = 0, \quad \forall \phi_h \in \Phi_h. \quad (15)$$

Our variational problem is to find $\{(\rho_h^n, u_h^n, p_h^n) \in \Phi_h \times V_h \times Q_h; n = 1, \dots, N_T\}$ satisfying (15), (12b), (12c), and (13).

Theorem 6. Suppose that Φ_h is the P1-finite element space and $V_h \subset V$ and $Q_h \subset Q$ satisfy the inf-sup condition (11).

- (i) For any functions $\rho^0 \geq 0$, u^0 , and f there exists the unique solution (ρ_h, u_h, p_h) of (15), (12b), (12c), and (13).
- (ii) The solution ρ_h is non-negative and it satisfies

$$\|\bar{\rho}_h^n\|_{L^1(\Omega)} = \|\bar{\rho}_h^0\|_{L^1(\Omega)}, \quad \forall n = 0, \dots, N_T.$$

- (iii) The solution satisfies the estimate (14).

Remark 7. Examples of the choice of V_h and Q_h are P2/P1 elements, and P1+/P1 elements [2].

6. Concluding remarks

We have shown finite element schemes for two-fluids flow problems. For a wide class of finite element spaces we have shown the stability of the velocity under a mild assumption on the density. For a mass-conservative upwind scheme we have shown the full stability of the density, the velocity, and the pressure. Numerical results and the discussion on the convergence will be given in a forthcoming paper.

References

- [1] K. Baba and M. Tabata. On a conservative upwind finite element scheme for convective diffusion equations. *R.A.I.R.O., Analyse numérique/Numerical Analysis*, Vol. 15, pp. 3–35, 1981.
- [2] V. Girault and P. A. Raviart. *Finite Element Methods for Navier-Stokes Equations, Theory and Algorithms*. Springer, 1986.
- [3] S. Kaizu and M. Tabata. A finite element analysis of the density-dependent Navier-Stokes equations, to appear.

- [4] K. Ohmori. Convergence of the interface in the finite element approximation for two-fluid flows. In R. Salvi, editor, *The Navier-Stokes Equations: Theory and Numerical Methods*, Vol. 223 of *Lecture Notes in Pure and Applied Mathematics*, pp. 279–293. Marcel Dekker, 2001.
- [5] T. E. Tezduyar, M. Behr, and J. Liou. A new strategy for finite element computations involving boundaries and interfaces - the deforming-spatial-domain/space-time procedure: I. *Computer Methods in Applied Mechanics and Engineering*, Vol. 94, pp. 339–351, 1992.

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Hypergeometric solutions to the q -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE
Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI
Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI
Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA
Ergodicity and exponential β -mixing bounds for a strong solution of Lévy-driven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI
The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI
Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI
On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA
Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV
Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS
Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA
The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON
Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON
Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER
Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift

- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations
- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems