三相分割問題に対する双対定理(非線形解析学と凸解析学の研究)

Kawasaki, Hidefumi
Faculty of Mathematics, Kyushu University

<http://hdl.handle.net/2324/3375>

出版情報 : MHF Preprint Series. 2005-31, 2005-09-15. 九州大学大学院数理学研究院
バージョン :
権利関係 :

MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

A duality theorem for a three-phase partition problem

H. Kawasaki

MHF 2005-31

(Received September 15, 2005)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

A duality theorem for a three-phase partition problem.¹

Hidefumi Kawasaki²

Abstract. In some nonlinear diffusive phenomena, the systems have three or more stable states. Sternberg and Zeimer (Ref. 1) established the existence of local minimizers to the problem of partitioning certain domain $\Omega \subset \mathbb{R}^2$ into three subdomains having least interfacial area. Ikota and Yanagida investigated stability and instability for stationary curves with one triple junction in (Ref. 2) and for stationary binary-tree type interfaces in (Ref. 3). In this paper, we consider a static version of the partitioning problem with a triple junction and present a duality theorem. The novelty of our duality theorem is that it is based on separation of three convex sets by a triangle.

Key words. Duality theorem, Separation, Convex set, Partition problem

1. INTRODUCTION

In some nonlinear diffusive phenomena, e.g., grain growth in annealing pure metal and segregation between biological species, the systems have three or more stable states (Fig.1). Sternberg and Zeimer (Ref. 1) established the existence of local minimizers to the problem of partitioning certain domain $\Omega \subset \mathbb{R}^2$ into three subdomains having least interfacial area. Furthermore, Ikota and Yanagida (Refs. 2, 3) investigated stability for stationary curves with one triple junction and of binary-tree type (Fig. 2).

FIGURE 1. Three-phase partition problem

Although they formulated partitioning problems as variational problems, they can be formulated as extremal problems in $\mathbb{R}^n$, since the shortest curve joining X_0 and X_i is the line segment X_0X_i . From this point of view, the author discussed stability and instability of the

¹This research is partially supported by the Grant-in Aid for General Scientific Research from the Japan Society for the Promotion of Science 14340037.

²Associate Professor, Faculty of Mathematics, Kyushu University, Japan

FIGURE 2. Binary-tree type interface

three-phase partition problem and studied its game-theoretic aspect in Kawasaki (Refs. 4, 5).

In this paper, we formulate the three-phase partition problem as follows. Let C_i ($i = 1, 2, 3$) be closed convex sets with non-empty interior in $\mathbb{R}^2$ such that $\Omega := \cap_{i=1}^3 C_i^c$ is non-empty (Fig. 3).

FIGURE 3. Primal problem

$$\begin{aligned}
 (P) \quad & \text{Minimize} \quad f(X_0, \dots, X_3) := \sum_{i=1}^3 \|X_i - X_0\| \\
 & \text{subject to} \quad X_i \in C_i \ (i = 1, 2, 3) \\
 & \quad \quad \quad X_0 \in \Omega.
 \end{aligned}$$

When we emphasize the domain Ω , we denote (P) by (P_Ω) . Although Ω is not convex, primal problem (P) can be regarded as a convex programming problem if X_0 is restricted to an open convex subset C_0 of Ω . The aims of this paper are to give a dual problem (D) and show duality between (P) and (D) .

This paper is organized as follows. In Section 2, we give first-order optimality conditions for (P) . In Section 3, we briefly review classical duality theorems and introduce a new concept of separation of three convex sets by a triangle. In Section 4, we define the dual problem (D) and show duality.

2. FIRST-ORDER OPTIMALITY CONDITION

In this section, we first give a first-order necessary optimality condition for (P) . Next, we consider the special case that C_i 's are closed half spaces.

A local minimizer $(X_0, \dots, X_3)$ of (P) is said to be non-degenerate if X_0 does not coincide with any X_i ($i = 1, 2, 3$).

Let $N(X_i; C_i)$ denote the normal cone of C_i at X_i , that is,

$$N(X_i; C_i) := \{\xi \in \mathbb{R}^2 \mid \xi^T(X - X_i) \leq 0 \quad \forall X \in C_i\}.$$

Theorem 2.1. *Let $(X_0, \dots, X_3)$ be a non-degenerate minimal solution for (P) , Then it satisfies Young's law*

$$\angle X_i X_0 X_j = 120^\circ \text{ for any } i \neq j$$

and the transversality condition

$$X_0 - X_i \in N(X_i; C_i) \quad (i = 1, 2, 3).$$

FIGURE 4. Young's law and the transversality condition

Proof. According to Kuhn-Tucker's theorem, see e.g. Rockafellar (Ref. 6, Section 28), there exist Kuhn-Tucker multipliers $\lambda_i \geq 0$ ($i = 1, 2, 3$) such that $0 \in \mathbb{R}^8$ belongs to the subdifferential of the Lagrange function

$$L(X_0, \dots, X_3) := f(X_0, \dots, X_3) + \sum_{i=1}^3 \lambda_i \delta(X_i | C_i),$$

where $\delta(X_i | C_i)$ denotes the characteristic function of C_i . Picking up X_0 -component of the subdifferential ∂L , we have

$$\sum_{i=1}^3 \frac{X_i - X_0}{\|X_i - X_0\|} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}. \quad (1)$$

Denoting

$$e_i := (X_i - X_0) / \|X_i - X_0\|, \quad (2)$$

we get from (1) that $\|e_k\|^2 = \|e_i\|^2 + \|e_j\|^2 + 2e_i^T e_j$, where $\{i, j, k\} = \{1, 2, 3\}$. Hence $e_i^T e_j = -1/2$, which implies Young's law (Fig. 4).

Picking up X_i -component ($i = 1, 2, 3$) of ∂L , we have

$$\begin{pmatrix} 0 \\ 0 \end{pmatrix} \in \frac{X_i - X_0}{\|X_i - X_0\|} + \lambda_i N(X_i; C_i). \quad (3)$$

Hence we get the transversality condition. $\square$

Next, we consider the special case that each C_i is a closed half space defined by

$$C_i = \{X_i \mid \xi_i^T X_i \leq \alpha_i\}, \quad (4)$$

where ξ_i is assumed to be a unit vector. Then (P) becomes a convex programming problem

$$(P) \quad \begin{aligned} & \text{Minimize} && \sum_{i=1}^3 \|X_i - X_0\| \\ & \text{subject to} && \xi_i^T X_i \leq \alpha_i \quad (i = 1, 2, 3) \\ & && \xi_i^T X_0 \geq \alpha_i \quad (i = 1, 2, 3), \end{aligned}$$

and optimal solutions are characterized by the first-order optimality condition. Furthermore, $N(X_i; C_i) = \{\lambda_i \geq 0 \mid \lambda_i \xi_i\}$ at $X_i \in \partial C_i$. Hence, if $(X_0, \dots, X_3)$ satisfies both Young's law and the transversality condition, then Ω must be a regular triangle (Fig. 5). Hence we have

$$\min(P) = \Omega\text{'s height}. \quad (5)$$

FIGURE 5. Young's law and the transversality condition are satisfied only when $\theta_1 = \theta_2 = \theta_3$.

Otherwise, for any minimum solution $(X_0, \dots, X_3)$, X_0 must be on the boundary of Ω .

Proposition 2.1. *Assume that C_i 's are defined by (4). If $(X_0, \dots, X_3)$ satisfying $X_0 = X_1 \notin \{X_2, X_3\}$ is a minimum solution for (P), then $\angle X_2 X_0 X_3 \geq 2\pi/3$, the normal vector ξ_1 equally divides angle $\angle X_2 X_0 X_3$, and line segment $X_0 X_i$ ($i = 2, 3$) orthogonally intersects ∂C_i , respectively. Furthermore, Ω is an isosceles triangle and*

$$\min(P) = \text{the smaller height of } \Omega. \quad (6)$$

FIGURE 6. Degenerate case: $X_0 = X_1 \neq X_2, X_3$

Proof. There exist $\lambda_i \geq 0$ ($i = 1, 2, 3$) and $\mu_1 \geq 0$ such that $0 \in \mathbb{R}^8$ belongs to the subdifferential of the Lagrange function

$$L(X_0, \dots, X_3) := \sum_{i=1}^3 \|X_i - X_0\| + \sum_{i=1}^3 \lambda_i (\xi_i^T X_i - \alpha_i) + \mu_1 (\alpha_1 - \xi_1^T X_0). \quad (7)$$

We regard $\|X_1 - X_0\|$ as a function of not (X_0, X_1) but $\mathbf{X} := (X_0, \dots, X_3)$, and denote it by $f_1(X_0, \dots, X_3)$. Then it follows from (Ref. 6, Theorem 23.9) that its subdifferential at $X_0 = X_1$ w.r.t. $\mathbf{X}$ is given by

$$\partial f_1(\mathbf{X}) = \left\{ (s, t) \begin{pmatrix} -1 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix} \mid s^2 + t^2 \leq 1 \right\} \quad (8)$$

$$= \left\{ (-s, -t, s, t, 0, 0, 0, 0) \mid s^2 + t^2 \leq 1 \right\}. \quad (9)$$

So there exists (s, t) such that $s^2 + t^2 \leq 1$,

$$(X_0 - \text{component}) \quad (s, t) + e_2 + e_3 + \mu_1 \xi_1 = (0, 0), \quad (10)$$

$$(X_1 - \text{component}) \quad (s, t) + \lambda_1 \xi_1 = (0, 0), \quad (11)$$

$$(X_i - \text{component}) \quad e_i + \lambda_i \xi_i = (0, 0) \quad (i = 2, 3), \quad (12)$$

where $e_i = (X_i - X_0)/\|X_i - X_0\|$. (12) implies the transversality condition $e_i = -\xi_i$ at X_i . We see from (11) that $0 \leq \lambda_1 \leq 1$. Substituting (11) into (10), we have

$$e_2 + e_3 = (\lambda_1 - \mu_1) \xi_1. \quad (13)$$

We see from (13) that ξ_1 equally divides angle $\angle X_2 X_0 X_3$ and $\lambda_1 - \mu_1 \geq 0$. So, $\|e_2 + e_3\| = \|(\lambda_1 - \mu_1) \xi_1\| \leq 1$. Hence $e_2^T e_3 \leq -1/2$. It follows from the transversality condition that Ω has to be an isosceles triangle (Fig. 6). Since θ_2 in Fig. 6 is greater than or equal to $\pi/3$, $\min(P)$ is equal to the smaller height of the isosceles triangle. $\square$

Finally, we consider the second degenerate case: $X_0 = X_1 = X_2 \neq X_3$.

Proposition 2.2. *Assume that C_i 's are defined by (4). If $(X_0, \dots, X_3)$ satisfying $X_0 = X_1 = X_2 \neq X_3$ is a minimum solution for (P), then line segment $X_0 X_3$ orthogonally intersects ∂C_3 and $X_0 X_3$ is the smallest height of the triangle. Hence*

$$\min(P) = \text{the smallest height of } \Omega. \quad (14)$$

Proof. There exist $\lambda_i \geq 0$ ($i = 1, 2, 3$) and $\mu_i \geq 0$ ($i = 1, 2$) such that $0 \in \mathbb{R}^8$ belongs to the subdifferential of the Lagrange function

$$L(X_0, \dots, X_3) := \sum_{i=1}^3 \|X_i - X_0\| + \sum_{i=1}^3 \lambda_i (\xi_i^T X_i - \alpha_i) + \sum_{i=1}^2 \mu_i (\alpha_i - \xi_i^T X_0). \quad (15)$$

FIGURE 7. Degenerate case: $X_0 = X_1 = X_2 \neq X_3$

Regarding $\|X_2 - X_0\|$ as a function of $\mathbf{X} = (X_0, \dots, X_3)$, we denote it by $f_2(X_0, \dots, X_3)$. Then it follows from (Ref. 6, Theorem 23.9) that its subdifferential at $X_0 = X_1$ w.r.t. $\mathbf{X}$ is given by

$$\partial f_2(\mathbf{X}) = \{(-u, -v, 0, 0, u, v, 0, 0) \mid u^2 + v^2 \leq 1\}. \quad (16)$$

So there exist (s, t) and (u, v) such that $s^2 + t^2 \leq 1$, $u^2 + v^2 \leq 1$,

$$(X_0 - \text{component}) \quad (s, t) + (u, v) + e_3 + \mu_1 \xi_1 + \mu_2 \xi_2 = (0, 0), \quad (17)$$

$$(X_1 - \text{component}) \quad (s, t) + \lambda_1 \xi_1 = (0, 0), \quad (18)$$

$$(X_2 - \text{component}) \quad (u, v) + \lambda_2 \xi_2 = (0, 0), \quad (19)$$

$$(X_3 - \text{component}) \quad e_3 + \lambda_3 \xi_3 = (0, 0). \quad (20)$$

(20) implies the transversality condition $e_3 = -\xi_3$ at X_3 . We see from (18) that $0 \leq \lambda_1 \leq 1$. Substituting (18), (19), and (20) into (17), we have

$$(\lambda_1 - \mu_1)\xi_1 + (\lambda_2 - \mu_2)\xi_2 + \xi_3 = 0. \quad (21)$$

Since each ξ_i is a normal vector of the half space C_i and since C_i 's form a triangle (Fig. 8), it is easily seen that $\sin \theta_i > 0$ and

$$\sin \theta_1 \xi_1 + \sin \theta_2 \xi_2 + \sin \theta_3 \xi_3 = 0. \quad (22)$$

Hence we see from (21) and (22) that $0 \leq \lambda_i - \mu_i \leq 1$, ($i = 1, 2$). It is clear from Fig. 8 that $\xi_i^T \xi_j = -\cos \theta_k$, where $\{i, j, k\} = \{1, 2, 3\}$. Hence, from (21), we get

$$\lambda_1 - \mu_1 - (\lambda_2 - \mu_2) \cos \theta_3 - \cos \theta_2 = 0. \quad (23)$$

$$-(\lambda_1 - \mu_1) \cos \theta_3 + \lambda_2 - \mu_2 - \cos \theta_1 = 0. \quad (24)$$

It follows from (23) and (24) that $\sin \theta_2 = (\lambda_2 - \mu_2) \sin \theta_3 \leq \sin \theta_3$. Similarly, we have $\sin \theta_1 \leq \sin \theta_3$. Since $\theta_1 + \theta_2 + \theta_3 = \pi$, we see that θ_3 is the largest angle. So $X_0 X_3$ is the smallest height of the triangle. $\square$

Unifying (5), (6), and (14), we concludes that

Corollary 2.1. *When Ω is a triangle, $\min(P)$ equals to the smallest height of Ω .*

FIGURE 8. Normal vectors of the triangle

3. CLASSICAL DUALITY THEOREMS AND SEPARATION BY A TRIANGLE

In this section, we first review classical duality theorems in brief. Next, we introduce a new concept of separating three convex sets by a triangle. For the sake of simplicity, we choose $\mathbb{R}^2$ as the stage.

One of the simplest duality theorems is the following. Let C_1 be a non-empty convex set in $\mathbb{R}^2$ and $A \notin C_1$ a point. Then the primal problem is

$$(P_1) \quad \begin{array}{ll} \text{Minimize} & \|X_1 - A\| \\ \text{subject to} & X_1 \in C_1, \end{array}$$

where $\|\cdot\|$ denotes the Euclidean norm.

FIGURE 9. Dual problem (D_1)

Its dual problem is to maximize the distance from A to a hyperplane that separates A and C_1 (Fig. 9).

$$(D_1) \quad \begin{array}{ll} \text{Maximize} & d(A, H) \\ \text{subject to} & H \text{ separates } A \text{ and } C_1, \end{array}$$

where $d(A, H) := \min\{\|X - A\| \mid X \in H\}$.

We can rephrase the dual problem as maximizing the width of a strip that separates A and C_1 (Fig. 10).

If we replace A by a convex set C_2 such that $C_1 \cap C_2 = \emptyset$, then the primal problem is as follows.

$$(P_2) \quad \begin{array}{ll} \text{Minimize} & \|X_1 - X_2\| \\ \text{subject to} & X_i \in C_i \ (i = 1, 2). \end{array}$$

FIGURE 10. Another expression of (D_1)

Its dual problem (D_2) is to minimize the width of a strip that separates C_1 and C_2 (Fig. 11).

FIGURE 11. Dual problem (D_2)

If we take the epigraph of a convex function f and the hypograph of a concave function g as C_1 and C_2 , respectively, and measure the width of the strip in the vertical direction, duality between (P_2) and (D_2) reduces to Fenchel's duality (Fig. 11), see e.g. (Ref. 6, Theorem 31.1).

FIGURE 12. Fenchel's duality

$$(P_F) \quad \text{Minimize} \quad f(x) - g(x),$$

$$(D_F) \quad \text{Maximize} \quad g_*(y) - f^*(y),$$

where $f^*(y) := \sup_x \{y^T x - f(x)\}$ and $g_*(y) := \inf_y \{y^T x - g(x)\}$.

Definition 3.1. Let C_i ($i = 1, 2, 3$) be convex sets in $\mathbb{R}^2$ such that $\Omega = \cap_{i=1}^3 C_i^c$ is not empty, and $\Delta \subset \Omega$ a triangle. Then, we say that Δ separates $\{C_i\}_{i=1}^3$ if there are three closed half spaces $\{H_i^-\}_{i=1}^3$ such that $C_i \subset H_i^-$ for every i and $\Delta = \cap_{i=1}^3 H_i^+$, where H_i^+ denotes the closed half space opposite to H_i^- (Fig. 13).

FIGURE 13. Δ_1 separates C_i 's and Δ_2 does not separate C_i 's.

Lemma 3.1. *Let $(X_0, \dots, X_3)$ be a minimum solution for (P_Ω) and let a triangle Δ separate $\{C_i\}_{i=1}^3$. If X_0 belongs to Δ , then $\min(P_\Delta) \leq \min(P_\Omega)$.*

Proof. Since $X_i \in C_i \subset H_i^-$, we have

$$\min(P_\Omega) = \sum_{i=1}^3 \|X_i - X_0\| \geq \sum_{i=1}^3 d(X_0, H_i^-) \geq \min(P_\Delta).$$

□

4. DUALITY THEOREM

The following (D) is our dual problem. When Ω is bounded, it has a simplified form (D^*) .

(D) Maximize the smallest height of a triangle Δ
 subject to there exists a triangle Δ' such that $\Delta \subset \Delta' \subset \Omega$,
 Δ' separates $\{C_i\}_{i=1}^3$, and $X_0 \in \Delta'$.

Theorem 4.1. *If (X_0, X_1, X_2, X_3) is a non-degenerate minimum for (P_Ω) , then*

$$\min(P_\Omega) = \max(D). \quad (25)$$

Proof. Combining Lemma 3.1 and Corollary 2.1, we have

$$\min(P_\Omega) \geq \min(P_{\Delta'}) \geq \text{the smallest height of } \Delta. \quad (26)$$

By Theorem 2.1, the non-degenerate minimum solution forms a regular triangle Δ^* such that

$$\min(P_\Omega) = \text{the smallest height of } \Delta^*. \quad (27)$$

It follows from definition of the normal cone that Δ^* itself separates $\{C_i\}_{i=1}^3$. Therefore, Δ^* attains the maximum of (D). □

Theorem 4.2. *When Ω is bounded, the dual problem (D) is simplified as follows.*

(D*) Maximize the smallest height of a triangle Δ
 subject to $X_0 \in \Delta \subset \Omega$.

Proof. Assume that $\Delta \subset \Omega$. Then, by separation theorem, there are closed half spaces H_i^- ($i = 1, 2, 3$) such that $C_i \subset H_i^-$ and $\Delta \subset \cap_{i=1}^3 (H_i^+)^c =: \Delta'$. Since Ω is bounded and since $\Delta' \subset \cap_{i=1}^3 C_i^c = \Omega$, Δ' is a triangle separating C_i ($i = 1, 2, 3$), see Fig. 14. $\square$

FIGURE 14. Separating hyperplanes form an unbounded polygon.

5. CONCLUDING REMARK

When Ω is not bounded, separating hyperplanes do not necessarily form a triangle, see Fig. 14. So duality relation $\min(P) = \max(D^*)$ does not always hold. Indeed, since we can enlarge Δ rightward within the dark gray area as we like, $\sup(D^*)$ equals $+\infty$.

References

1. Sternberg, P., and Zeimer W. P., *Local minimizers of a three-phase partition problem with triple junctions*, Proceedings of the Royal Society of Edinburgh, Vol. 124A, pp. 1059–1073, 1994.
2. Ikota, R., and Yanagida, E., *A stability criterion for stationary curves to the curvature-driven motion with a triple junction*, Differential and Integral Equations, Vol. 16, pp. 707–726, 2003.
3. Ikota, R., and Yanagida, E., *Stability of stationary interfaces of binary-tree type*, Partial Differential Equations, Vol. 22, pp. 375–389, 2004.
4. Kawasaki, H., *A game-theoretic aspect of conjugate sets for a nonlinear programming problem*, Proceedings of the third International Conference on Nonlinear Analysis and Convex Analysis, Edited by W. Takahashi and T. Tanaka, Yokohama Publishers, Yokohama, pp. 159–168, 2004.
5. Kawasaki, H., *Conjugate-set game for a nonlinear programming problem*, to appear in Game theory and applications 10, Edited by L. A. Petrosyan, Nova Science Publications, New York, USA, pp. 87–95, 2005.
6. Rockafellar, R. T., *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Hypergeometric solutions to the q -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE
Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI
Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI
Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA
Ergodicity and exponential β -mixing bounds for a strong solution of Lévy-driven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI
The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI
Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI
On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA
Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV
Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS
Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA
The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON
Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON
Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER
Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q -Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift

- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations
- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in $\mathbb{R}^d$
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA
Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS
Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI
Nonlinear regression modeling via regularized wavelets and smoothing parameter selection

- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO,
Yoshihiro MIZOGUCHI & Yasuo KAWAHARA
On reversible cellular automata with finite cell array
- MHF2005-28 Toru KOMATSU
Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU
A computational approach to constructive a priori and a posteriori error
estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO
Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI
A duality theorem for a three-phase partition problem