九州大学学術情報リポジトリ Kyushu University Institutional Repository

Numerical Simulation of Rayleigh-Taylor Problems by an Energy-Stable Finite Element Scheme

Tabata, Masahisa Faculty of Mathematics, Kyushu University

https://hdl.handle.net/2324/3379

出版情報:Proceedings of The Fourth International Workshop on Scientific Computing and Applications, pp.63-73, 2007. Faculty of Mathematics, Kyushu University バージョン: 権利関係:

MHF Preprint Series

Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme

M. Tabata

MHF 2006-1

(Received January 5, 2006)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN

Numerical Simulation of Rayleigh-Taylor Problems by an Energy-Stable Finite Element Scheme

Masahisa TABATA * Faculty of Mathematics, Kyushu University Hakozaki, Fukuoka, 812-8581 JAPAN

January 5, 2006

Abstract

Numerical simulation of some Rayleigh-Taylor problems is performed by an energy-stable finite element scheme. The problems are solved in the frame work of the density-dependent Navier-Stokes equations. Mesh dependency of numerical results is discussed for the Rayleigh-Taylor instability problem and perturbed stable problems.

1 Introduction

The Rayleigh-Taylor problem describes the movement of two fluids subject to an initial state, where a heavy fluid is located in the upper part of a container and a light fluid is in the lower part. It is one of two-fluid flow problems with unknown interfaces. In this paper we simulate such problems by an energystable P1/P2/P1 finite element scheme developed recently by ourselves [6], where the density, the velocity, and the pressure are approximated by linear, quadratic, and linear finite element, respectively.

Many numerical results have been obtained for two-fluid flow problems, e.g. [5], [8], but, to the best of our knowledge, there are no numerical schemes

^{*}Email:tabata@math.kyushu-u.ac.jp

whose solutions are proved to converge to the exact one. Recently we have developed a P0/P1N/P0 finite element scheme for density-dependent Navier-Stokes equations and proved the convergence of the finite element solutions [4]. Here P0 and P1N stand for constant and non-conforming linear element, respectively. After reformulating the original two-fluid flow problem by a density-dependent Navier-Stokes problem, we can apply the P0/P1N/P0 scheme to two-fluid flow problems. Some numerical results for the Rayleigh-Taylor problem by this scheme have been reported in [7].

The P1/P2/P1 finite element scheme is applied to Rayleigh-Taylor problems also in the framework of a density-dependent Navier-Stokes problem, where the density is treated as a field function defined in the whole domain. This scheme has the advantage that the stress interface condition can be treated more easily and the approximation ability is higher than the P0/P1N/P0 scheme. We solve some Rayleigh-Taylor problems subject to an unstable equilibrium and perturbed non-equilibrium initial states and discuss the effect caused by the mesh subdivision.

2 Two-fluid flow problems

We begin by describing the general two-fluid flow problem. Let Ω be a bounded domain in \mathbf{R}^d , d = 2, 3, with piecewise smooth boundary Γ , and Tbe a positive number. At the initial time t = 0 the domain Ω is occupied by two immiscible incompressible viscous fluids; each domain is denoted by Ω_k^0 , k = 1, 2, whose interface $\partial \Omega_1^0 \cap \partial \Omega_2^0$ is denoted by Γ_{12}^0 . At $t \in (0, T)$ the two fluids occupy domains $\Omega_k(t)$, k = 1, 2, and the interface $\partial \Omega_1(t) \cap \partial \Omega_2(t)$ is denoted by $\Gamma_{12}(t)$. Let ρ_k and μ_k , k = 1, 2, be the densities and the viscosities of the two fluids. Let

$$u: \Omega \times (0,T) \to \mathbf{R}^d, \quad p: \Omega \times (0,T) \to \mathbf{R}$$

be the velocity and the pressure to be found. The Navier-Stokes equations are satisfied in each domain $\Omega_k(t)$, $k = 1, 2, t \in (0, T)$,

$$\rho_k \left\{ \frac{\partial u}{\partial t} + (u \cdot \nabla)u \right\} - \nabla \left(2\mu_k D(u) \right) - \nabla p = \rho_k f$$
(1a)

$$\nabla \cdot u = 0, \qquad (1b)$$

where $f: \Omega \times (0,T) \to \mathbf{R}^d$ is a given function, D(u) is the strain tensor defined by

$$D_{ij}(u) = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_j} + \frac{\partial u_j}{\partial x_i} \right)$$

and

$$\left[\nabla \left(2\mu_k D(u)\right)\right]_i = \sum_{j=1}^d \frac{\partial}{\partial x_j} \left(2\mu_k D_{ij}(u)\right).$$

On the boundary Γ , $t \in (0, T)$, the slip conditions

$$u \cdot n = 0, \quad \sigma(\mu, u, p)n \times n = 0$$
 (2)

are imposed, where n is the unit outer normal to Γ and σ is the stress tensor defined by

$$\sigma(\mu, u, p) = -pI + 2\mu D(u).$$

On the interface $\Gamma_{12}(t)$, $t \in (0, T)$, the velocity and the stress vector should be continuous,

$$[u] = 0, \quad [\sigma(\mu, u, p)n_{12}] = 0, \tag{3}$$

where $[\cdot]$ means the difference of the values approaching from the domain Ω_2 and the domain Ω_1 , n_{12} is the unit outer normal to $\Gamma_{12}(t)$ from $\Omega_1(t)$ to $\Omega_2(t)$. The initial condition at t = 0 for the velocity

$$u = u^0 \tag{4}$$

is imposed.

We consider the case where a heavy fluid is located in the upper part and a light fluid is in the lower part as shown in Fig. 1. Given such an initial state, we consider the transitional movement of the fluids.

3 An energy-stable finite element scheme

We have developed a class of energy-stable finite element schemes for problem (1)-(4) in [6]. The main idea is to convert the original problem to a density-dependent Navier-Stokes problem, where the density is treated as a field function of x and t. The viscosity μ is also supposed to be a function of ρ . In the case of two-fluid flow problems μ is defined by

$$\mu(\rho) = \mu_1 \frac{\rho_2 - \rho}{\rho_2 - \rho_1} + \mu_2 \frac{\rho - \rho_1}{\rho_2 - \rho_1}.$$
(5)



Figure 1: The interface $\Gamma_{12}(t)$ (left) and the barycentric domain D_P (right)

Here we only describe a P1/P2/P1 scheme, that is, the density, the velocity, and the pressure are approximated by P1-, P2-, and P1-element, respectively. We consider the 2-dimensional case d = 2. For the extension to the 3-dimensional case and the derivation of the scheme we refer to [6].

Let Φ_h , V_h , and Q_h be P1-, P2-, and P1-finite element space for the density, the velocity, and the pressure. The essential boundary condition is imposed on V_h ,

$$V_h = \{v_h; (v_h \cdot n)(P) = 0 \text{ (node } P \in \Gamma)\},\$$

but is not on Φ_h and Q_h . Let Δt be a time increment and $N_T = \lfloor T/\Delta t \rfloor$. We denote by (ρ_h^n, u_h^n, p_h^n) the value at time $n\Delta t$, and by $\bar{D}_{\Delta t}$ the backward difference operator, e.g.,

$$\bar{D}_{\Delta t} u_h^n = \frac{u_h^n - u_h^{n-1}}{\Delta t}$$

We find $\{(\rho_h^n, u_h^n, p_h^n) \in \Phi_h \times V_h \times Q_h; n = 1, \cdots, N_T\}$ satisfying

$$(\bar{D}_{\Delta t}\bar{\rho}_{h}^{n},\bar{\phi}_{h}) + c_{1h}(u_{h}^{n-1},\rho_{h}^{n},\phi_{h}) = 0, \quad \forall \phi_{h} \in \Phi_{h}$$

$$(6a)
\left(\rho_{h}^{n-1}\bar{D}_{\Delta t}u_{h}^{n} + \frac{1}{2}u_{h}^{n}\bar{D}_{\Delta t}\rho_{h}^{n},v_{h} \right) + a_{1}(\rho_{h}^{n},u_{h}^{n-1},u_{h}^{n},v_{h}) + a_{0}(\rho_{h}^{n},u_{h}^{n},v_{h})
+ b(v_{h},p_{h}^{n}) = (\rho_{h}^{n}\Pi_{h}f^{n},v_{h}), \quad \forall v_{h} \in V_{h}$$

$$(6b)
b(u_{h}^{n},q_{h}) = 0, \quad \forall q_{h} \in Q_{h}$$

$$(6c)$$

subject to the initial conditions

$$\rho_h^0 = \Pi_h \rho^0, \quad u_h^0 = \Pi_h u^0.$$
(7)

Here ρ^0 is a function defined by

$$\rho^0(x) = \begin{cases} \rho_1 & (x \in \Omega_1^0)\\ \rho_2 & (x \in \Omega_2^0), \end{cases}$$

 Π_h is the interpolation operator to the corresponding finite element spaces, (\cdot, \cdot) is the inner product in $L^2(\Omega)$ or $L^2(\Omega)^2$, and linear forms a_1, a_0 , and b are defined by

$$a_1(\rho, w, u, v) = \int_{\Omega} \left(\frac{1}{2} (w \cdot \nabla \rho) u + \frac{1}{2} \rho (\nabla \cdot w) u + \rho (w \cdot \nabla) u \right) \cdot v \, dx$$

$$a_0(\rho, u, v) = \int_{\Omega} \mu(\rho) D(u) : D(v) \, dx, \quad b(v, q) = -\int_{\Omega} (\nabla \cdot v) q \, dx.$$

We now explain (6a), which is based on an upwind approximation developed in [1]. At first we make the dual decomposition $\{D_P\}$ of the domain Ω , where D_P is the barycentric domain of the node P, see Fig. 1. The D_P is made by connecting the midpoints of the sides and the centroids of the triangles around $P \in \Omega$. When the node P is on Γ , some parts of the (approximate) boundary are used. We define the lumping operator $\bar{}$ from Φ_h to $L^2(\Omega)$ by

$$\phi_h \longmapsto \bar{\phi}_h(x) \equiv \sum_P \phi_h(P) \chi_P(x),$$

where χ_P is the characteristic function of D_P . We denote by Λ_P the set of nodes adjacent to P. The form c_{1h} is defined by

$$c_{1h}(u_h,\rho_h,\phi_h) = \sum_P \phi_h(P) \sum_{Q \in \Lambda_P} \beta_{PQ}^-(u_h)(\rho_h(P) - \rho_h(Q)), \qquad (8)$$

where

$$\beta_{PQ}^{-}(u_h) = \max(-\int_{\Gamma_{PQ}} u_h \cdot n_{PQ} \, ds, \, 0), \quad \Gamma_{PQ} = \partial D_P \cap \partial D_Q,$$

and n_{PQ} is the unit outer normal to Γ_{PQ} from D_P to D_Q . (8) is an upwind approximation of the trilinear form

$$c_1(u,\rho,\phi) = \int_{\Omega} (u \cdot \nabla \rho) \phi \ dx,$$

and we refer to (4.13) of [1] for the derivation of (8). Without any restriction on Δt it can be proved that the solution ρ_h of (6) satisfies the maximum principle and that scheme (6) is energy-stable, i.e.,

$$\int_{\Omega} \rho_h^n |u_h^n|^2 dx$$

is uniformly bounded for $n = 0, \dots, N_T$, see [1] and [6].

4 Numerical results

Let $\Omega \equiv (0,1) \times (0,1)$ be a unit square in \mathbb{R}^2 . Letting $\eta : [0,1] \to \mathbb{R}$ be a given continuous function, we define

$$\Omega_2^0 \equiv \{ x \in \Omega; x_2 \ge \eta(x_1) \}, \quad \Omega_1^0 \equiv \{ x \in \Omega; x_2 < \eta(x_1) \}.$$

We take the following values,

$$(\rho_1, \mu_1) = (10, 1), \quad (\rho_2, \mu_2) = (100, 2)$$

and the initial velocity and the outer force,

$$u^0 = (0,0)^T, \quad f = (0,-1)^T.$$

We divide the domain Ω into a union of triangles. We consider three kinds of meshes, Union-Jack type (UJ), Friedricks-Keller type (FK), and FreeFEM type (Free) shown in Fig. 2. The last mesh is made by FreeFEM [2]. Each side of the square is divided into 32 equal segments. We take $\Delta t = \frac{1}{2}$. The function η is chosen to satisfy

$$\int_0^1 \eta(x_1) dx_1 = \frac{1}{2}.$$

In the following figures we show computed interface curves approximating $\Gamma_{12}(t)$. They are determined so that the areas of the two fluids are equal to each other. Driven by the outer force f, the heavy fluid goes down. After a period of time elapses, the heavy fluid finally occupies the lower half part. We observe the transient movement for the period [0, 10].



Figure 2: Meshes of type UJ, FK, and Free

4.1 The Rayleigh-Taylor instability problem

The function η is defined by

$$\eta(x_1) = \frac{1}{2}.\tag{9}$$

The initial state is in equilibrium but unstable as the solution of partial differential equations. Figs. 3, 4, and 5 show the interface curves at t = 0, 2, 4, 6, 8, 10 in meshes UJ, FK, and Free, respectively. Mesh Free is not uniform. Mesh FK is not symmetric with respect to $x_1 = 1/2$, which induces the movement but it differs from the one in mesh Free. The heavy fluid goes down from left. In mesh UJ the interface almost keeps the original position in this period, though a very small change can be observed. Later it develops and finally the heavy fluid goes down. Three movements are completely different from one another and are heavily dependent on the meshes. Such numerical solutions are often called "ghost solutions" because they never appear in the partial differential equation but only in discrete approximations. Applying numerical methods to unstable problems, we may have ghost solutions. In the next subsection, however, our scheme is shown to work well for well-posed problems.

4.2 A perturbed Rayleigh-Taylor problem

The function η is defined by

$$\eta(x_1) = \frac{1}{2} - a\cos 2\pi x_1, \quad a = 0.01.$$
(10)



Figure 3: Interfaces at t = 0, 2, 4, 6, 8, 10. η :(9), mesh UJ

The initial state is a little perturbed, that is, the heavy fluid intrudes a little from both sides near the boundary. Figs. 6, 7, and 8 show the interface curves in meshes UJ, FK, and Free, respectively. These three figures are much similar to one another, and the mesh dependency is small. The differences decrease when the mesh size tends to zero.

4.3 The other problems

In mesh UJ we solve problems with other initial states.

Fig. 9 shows the case where the function η is defined by

$$\eta(x_1) = \frac{1}{2} - a\cos 2\pi x_1, \quad a = -0.01.$$
(11)

The initial state is a little perturbed, that is, the heavy fluid intrudes a little from the center. The small intrusion ignites the fall of the heavy fluid from the center.

Fig. 10 shows the case where the function η is defined by

$$\eta(x_1) = \frac{1}{2} - a\cos 4\pi x_1, \quad a = 0.01.$$
(12)



Figure 4: Interfaces at t = 0, 2, 4, 6, 8, 10. η :(9), mesh FK

The small intrusion of the heavy fluid from both sides and the center ignites the fall of the fluid from these parts.

5 Concluding remarks

We have simulated numerically Rayleigh-Taylor problems by an energy-stable P1/P2/P1 finite element scheme. For Rayleigh-Taylor problems subject to non-equilibrium states numerical interface curves are almost mesh-independent. For the Rayleigh-Taylor instability problem subject to an unstable equilibrium state ghost solutions are obtained. Such phenomenon is often observed in solving numerically unstable problems.

Acknowledgments

This work was supported by the Japan Society for the Promotion of Science under Grant-in-Aid for Scientific Research (S), No.16104001 and by the Ministry of Education, Culture, Sports, Science and Technology of Japan under Kyushu University 21st Century COE Program, Development of Dynamic



Figure 5: Interfaces t t = 0, 2, 4, 6, 8, 10. η :(9), mesh Free

Mathematics with High Functionality.

References

- K. Baba and M. Tabata. On a conservative upwind finite element scheme for convective diffusion equations. R.A.I.R.O., Analyse numérique/Numerical Analysis, Vol. 15, pp. 3–35, 1981.
- [2] http://www.freefem.org/.
- [3] V. Girault and P. A. Raviart. Finite Element Methods for Navier-Stokes Equations, Theory and Algorithms. Springer, 1986.
- [4] S. Kaizu and M. Tabata. A finite element analysis of the densitydependent Navier-Stokes equations, to appear.
- [5] K. Ohmori. Convergence of the interface in the finite element approximation for two-fluid flows. In R. Salvi, editor, *The Navier-Stokes Equations: Theory and Numerical Methods*, Vol. 223 of *Lecture Notes in Pure and Applied Mathematics*, pp. 279–293. Marcel Dekker, 2001.



Figure 6: Interfaces at t = 0, 2, 4, 6, 8, 10. η :(10), mesh UJ

- [6] M. Tabata and S. Kaizu. Finite element schemes for two-fluids flow problems. To appear in *Proceedings of 7th China-Japan Joint Seminar* for Computational Mathematics and Scientific Computing, Science Press, Beijing.
- [7] M. Tabata and Y. Fukushima. A Finite Element Approximation to Density-Dependent Navier-Stokes Equations and Its Application to Rayleigh-Taylor Instability Problem. In S. M. Sivakumar *et al.*, editors, *Advances in Computational & Experimental Engineering and Sciences*, pp. 455–460. Tech Science Press, India, 2005.
- [8] T. E. Tezduyar, M. Behr, and J. Liou. A new strategy for finite element computations involving boundaries and interfaces - the deformingspatial-domain/space-time procedure: I. Computer Methods in Applied Mechanics and Engineering, Vol. 94, pp. 339–351, 1992.



Figure 7: Interfaces at t=0,2,4,6,8,10. $\eta{:}(10).$ mesh FK



Figure 8: Interfaces at $t=0,2,4,6,8,10.~\eta{:}(10),$ mesh Free



Figure 9: Interfaces at t=0,2,4,6,8,10. $\eta{:}(11),$ mesh UJ



Figure 10: Interfaces at t = 0, 2, 4, 6, 8, 10. η :(12), mesh UJ

List of MHF Preprint Series, Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Hypergeometric solutions to the q Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA Ergodicity and exponential β -mixing bounds for a strong solution of Lévydriven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA Simple estimators for non-linear Markovian trend from sampled data: I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA Estimation for a discretely observed small diffusion process with a linear drift

- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA AIC for ergodic diffusion processes from discrete observations
- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI Nonlinear regression modeling via regularized wavelets and smoothing parameter selection

- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO, Yoshihiro MIZOGUCHI & Yasuo KAWAHARA On reversible cellular automata with finite cell array
- MHF2005-28 Toru KOMATSU Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI An explicit formula of the Shapley value for a cooperative game induced from the conjugate point
- MHF2005-34 Hideki MURAKAWA A regularization of a reaction-diffusion system approximation to the two-phase Stefan problem
- MHF2006-1 Masahisa TABATA Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme