

A DUALITY THEORY BASED ON TRIANGULAR CYLINDERS SEPARATING THREE CONVEX SETS IN \mathbb{R}^N

Kawasaki, Hidefumi
Faculty of Mathematics, Kyushu University

<http://hdl.handle.net/2324/3381>

出版情報 : MHF Preprint Series. MHF 2006-5, 2006-01-31. Faculty of Mathematics, Kyushu University
バージョン :
権利関係 :



MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

A duality theory based on
triangular cylinders separating
three convex sets in R^n

H. Kawasaki

MHF 2006-5

(Received January 31, 2006)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

A DUALITY THEORY BASED ON TRIANGULAR CYLINDERS SEPARATING THREE CONVEX SETS IN R^N *

H. KAWASAKI[†]

Abstract. Separation theorems play the central role in the duality theory. Recently, we proposed a duality theorem for a three-phase partition problem in [7]. It is based on triangles separating three convex sets in R^2 . The aim of this paper is to extend the duality theorem to R^n .

Key words. duality theorem, triangular cylinder, partition problem, separation theorem

AMS subject classifications. 49K10, 90C30, 26B99

1. Introduction. The three-phase partition problem is to divide a given domain $\Omega \subset R^2$ into three subdomains with a triple junction having least interfacial area (Fig.1.1).




FIG. 1.1. *Three-phase partition problem*

Sternberg and Zeimer [10] and Ikota and Yanagida [4] formulated this problem as variational problems and discussed stability of stationary solutions. However, since the shortest curve joining two points X_0 and X_i is the line segment X_0X_i , they can be formulated as extremal problems in a Euclidean space. From this point of view, we discussed stability and studied its game-theoretic aspect in [5][6]. Further, we gave a duality theorem for an extremal problem (P_0) below induced from the three-phase partition problem in [7].




FIG. 1.2. *Primal problem (P_0)*

*This research is partially supported by Kyushu University 21st Century COE Program (Development of Dynamic Mathematics with High Functionality) and the Grant-in Aid for General Scientific Research from the Japan Society for the Promotion of Science 14340037.

[†](kawasaki@math.kyushu-u.ac.jp), Faculty of Mathematics, Kyushu University 33, Fukuoka 812-8581, Japan.

$$(P_0) \quad \begin{aligned} & \text{Minimize} && f(X_0, \dots, X_3) := \sum_{i=1}^3 \|X_i - X_0\| \\ & \text{subject to} && X_0 \in \Omega, X_i \in C_i \ (i = 1, 2, 3), \end{aligned}$$

where $\|\cdot\|$ denotes the Euclidean norm and C_i ($i = 1, 2, 3$) are closed convex sets with non-empty interior in R^2 such that $\Omega := \text{cl}(\cap_{i=1}^3 C_i^c)$ is not empty (Fig. 1.2). The main aim of this paper is to extend the duality theorem (Theorem 2.3 below) to R^n .

This paper is organized as follows. In Section 2, we briefly review the first-order optimality condition for the primal problem (P_0) and the duality theorem given in [7]. In Section 3, we give a first-order optimality condition for the primal problem in R^n . In Section 4, we introduce the notion of separating three convex sets by a triangular cylinder in R^n to define the dual problem, and show the strong duality.

We close this section with our notations. For any closed convex sets C_1 and C_2 , we define $d(C_1, C_2) := \inf\{\|X_1 - X_2\| \mid X_i \in C_i \ (i = 1, 2)\}$. We denote by $N(X_i; C_i)$ the normal cone of C_i at X_i , that is, $N(X_i; C_i) := \{Y \in R^n \mid Y^T(X - X_i) \leq 0 \ \forall X \in C_i\}$.

2. Preliminaries. As is easily seen from Fig. 1.2, Ω is not always a convex set. So the primal problem (P_0) is not a convex programming problem. We modify it so that it becomes a convex programming problem.

$$(P) \quad \begin{aligned} & \text{Minimize} && \sum_{i=1}^3 \|X_i - X_0\| \\ & \text{subject to} && X_0 \in R^2, X_i \in C_i \ (i = 1, 2, 3). \end{aligned}$$

The only difference is that Ω is replaced by the whole space R^2 . We say a feasible solution (X_0, \dots, X_3) for (P_0) (or (P)) non-degenerate if $X_i \neq X_j$ for any $i \neq j$. The following is a straightforward consequence of Torricelli's Theorem and the projection theorem on the convex set C_i .

THEOREM 2.1. *Let (X_0, \dots, X_3) be a non-degenerate minimal solution for (P_0) (or (P)). Then it satisfies*

$$(2.1) \quad \angle X_i X_0 X_j = 2\pi/3 \text{ for any } i \neq j$$

and

$$(2.2) \quad X_0 - X_i \in N(X_i; C_i) \quad (i = 1, 2, 3).$$




FIG. 2.1. *First-order optimality conditions*

LEMMA 2.2. ([7]) When Ω is a triangle in R^2 , it holds that

$$\min(P) = \min(P_0) = \text{the smallest height of } \Omega.$$

We say that a triangle $\Delta \subset \Omega$ separates $\{C_i\}_{i=1}^3$ if there are three closed half spaces $\{H_i^-\}_{i=1}^3$ such that $C_i \subset H_i^-$ for every i and $\Delta = \cap_{i=1}^3 H_i^+$, where H_i^+ denotes the closed half space opposite to H_i^- (Fig. 2.2).




FIG. 2.2. Δ_1 separates $\{C_i\}_{i=1}^3$, and Δ_2 does not separate $\{C_i\}_{i=1}^3$.

Then the dual problem is defined as follows.

(D) Maximize the smallest height of a triangle that separates $\{C_i\}_{i=1}^3$.

THEOREM 2.3. ([7]) If (P_0) (or (P)) has a non-degenerate minimal solution, then

$$(2.3) \quad \max(D) = \min(P) = \min(P_0).$$

3. The primal problem in R^n . In this section, we extend Theorem 2.1 to R^n . Let C_i ($i = 1, 2, 3$) be closed convex sets with non-empty interior in R^n such that $\Omega = \text{cl}(\cap_{i=1}^3 C_i^c)$ is not empty. Then (P_0) and (P) are defined as well as in Sections 1 and 2, respectively, where the base space is R^n .




FIG. 3.1. The primal problem (P) in R^3 .

THEOREM 3.1. Let (X_0, \dots, X_3) be a non-degenerate minimal solution for (P_0) . Then X_i 's are on a two-dimensional affine set and (X_0, \dots, X_3) satisfies (2.1) and (2.2). Further, it is a minimum solution for (P) .

Proof. According to Kuhn-Tucker's theorem, see e.g. Rockafellar (Ref. 6, Section 28), there exist multipliers $\lambda_i \geq 0$ ($i = 1, 2, 3$) such that $0 \in R^{4n}$ belongs to the subdifferential of the Lagrange function $L(X_0, \dots, X_3) := \sum_{i=1}^3 \|X_i - X_0\| + \sum_{i=1}^3 \lambda_i \delta(X_i | C_i)$, where $\delta(X_i | C_i)$ denotes the characteristic function of C_i . Picking up X_0 -component of the subdifferential ∂L , we have

$$(3.1) \quad n_1 + n_2 + n_3 = 0 \in R^n,$$

where $n_i := (X_0 - X_i)/\|X_i - X_0\|$, which implies the first assertion. From (3.1), $\|n_k\|^2 = \|n_i\|^2 + \|n_j\|^2 + 2n_i^T n_j$ for any $\{i, j, k\} = \{1, 2, 3\}$. Thus $n_i^T n_j = -1/2$, which implies (2.1). Picking up X_i -component ($i = 1, 2, 3$) of ∂L , we have $0 \in -n_i + \lambda_i N(X_i; C_i)$, which implies (2.2). Next, there exists an open convex neighborhood C_0 of X_0 such that (X_0, \dots, X_3) is a minimum point of f on $C := C_0 \times C_1 \times C_2 \times C_3$. Since f and C are convex, (X_0, \dots, X_3) is a minimum point of f on $R^n \times C_1 \times C_2 \times C_3$. Hence it is a minimum solution for (P). \square

4. Duality theorem. In this section, we first introduce the notion of separation of three convex sets by a triangular cylinder. Next, we define the dual problem and show strong duality.

DEFINITION 4.1. Let n_i ($i = 1, 2, 3$) be nonzero (unit) vectors in R^n satisfying (3.1), and α_i ($i = 1, 2, 3$) negative real numbers. Define

$$(4.1) \quad H_i^+ := \{\xi \in R^n \mid n_i^T \xi \geq \alpha_i\},$$

$H_i^- := \{\xi \in R^n \mid n_i^T \xi \leq \alpha_i\}$, and $H_i := H_i^+ \cap H_i^-$ for any $i = 1, 2, 3$. Then we call a shifted figure of $H_1^+ \cap H_2^+ \cap H_3^+$ a (regular) triangular cylinder, see Fig. 4.1. (Here we remark that the origin of R^n belongs to $H_1^+ \cap H_2^+ \cap H_3^+$.) Further, we say a triangular cylinder $H_1^+ \cap H_2^+ \cap H_3^+ + X$ separates three convex sets $\{C_i\}_{i=1}^3$ if $C_i \subset H_i^- + X$ for any $i = 1, 2, 3$. Let $M := \cap_{i=1}^3 \{\xi \in R^n \mid n_i^T \xi = 0\}$ and

$$(4.2) \quad N := M^\top = \{X \in R^n \mid \xi^T X = 0 \quad \forall \xi \in M\}.$$

Then, by (3.1), N is a 2-dimensional subspace, and

$$(4.3) \quad \Delta := N \cap (H_1^+ \cap H_2^+ \cap H_3^+)$$

is a (regular) triangle. We call the smallest height of Δ the width of the triangular cylinder $H_1^+ \cap H_2^+ \cap H_3^+ + X$.




FIG. 4.1. Regular triangular cylinder in R^3

Our dual problem is defined as follows.

(D) Maximize the width of a triangular cylinder that separates $\{C_i\}_{i=1}^3$.

THEOREM 4.2. (Weak duality)

$$(4.4) \quad \sup(D) \leq \inf(P) \leq \inf(P_0).$$

Proof. Second inequality is trivial. Let (Y_0, \dots, Y_3) be a feasible solution for (P) , and $H_1^+ \cap H_2^+ \cap H_3^+ + X$ a feasible solution for (D) . Then $Y_i \in C_i \subset H_i^- + X$ for any $i = 1, 2, 3$. Hence

$$\|Y_0 - Y_i\| \geq d(Y_0, H_i^- + X) = d(Y_0 - X, H_i^-) = d(P_N(Y_0 - X), H_i^-),$$

where P_N denotes the projection to N defined by (4.2). Hence

$$\begin{aligned} \sum_{i=1}^3 \|Y_0 - Y_i\| &\geq \sum_{i=1}^3 d(P_N(Y_0 - X), H_i^-) \\ &= \min \left\{ \sum_{i=1}^3 \|P_N(Y_0 - X) - X_i\| \mid X_i \in H_i^- \ (i = 1, 2, 3) \right\} \\ &\geq \min \left\{ \sum_{i=1}^3 \|X_0 - X_i\| \mid X_i \in H_i^- \ (i = 1, 2, 3), X_0 \in N \right\} \\ &\geq \text{the smallest height of the triangle } \Delta \text{ defined by (4.3)} \\ &= \text{the width of } H_1^+ \cap H_2^+ \cap H_3^+ + X, \end{aligned}$$

where the last inequality follows from Lemma 2.2. Hence we get $\sup(D) \leq \inf(P)$. \square

THEOREM 4.3. (*Strong duality*) *If (P_0) (or (P)) has a non-degenerate minimal solution, then it holds that*

$$(4.5) \quad \max(D) = \min(P) = \min(P_0).$$

Proof. Second equality follows from Theorem 3.1. Let (X_0, \dots, X_3) be a non-degenerate minimal solution for (P_0) . Then, the regular triangular cylinder $H_1^+ \cap H_2^+ \cap H_3^+ + X_0$ determined by $n_i := (X_0 - X_i)/\|X_i - X_0\|$ and $\alpha_i := -\|X_i - X_0\|$ separates $\{C_i\}_{i=1}^3$. Indeed, it follows from (2.2) that $n_i^T X \leq n_i^T X_i$ for any $X \in C_i$, so that

$$n_i^T (X - X_0) \leq n_i^T (X_i - X_0) = -\|X_i - X_0\| = \alpha_i.$$

Namely, $C_i - X_0$ is contained in H_i^- . Moreover, since X_i ($i = 0, 1, 2, 3$) belong to the regular triangle $\Delta + X_0$, it follows from Lemma 2.2 that $\sum_{i=1}^3 \|X_0 - X_i\|$ is equal to the width of the regular triangular cylinder $H_1^+ \cap H_2^+ \cap H_3^+ + X_0$. Therefore we get (4.5). \square

REMARK 4.1. *We can replace triangular cylinders by regular triangular cylinders in the dual problem (D) , because the maximum value is attained by a regular triangular cylinder. However, it is clear that regular triangular cylinders are not enough when Ω is a non-regular triangular cylinder. That's why we defined the dual problem with (general) triangular cylinders.*

5. Extension to the weighted objective function. In [4], the objective function was weighted. It is not hard to extend the present results to the weighted objective function

$$\sum_{i=1}^3 \sigma_i \|X_i - X_0\|,$$

where $\sigma_i > 0$ ($i = 1, 2, 3$) can be interpreted as interface tension, see Fig. 5.1. We impose an assumption that $\sigma_i < \sigma_j + \sigma_k$ for any $\{i, j, k\} = \{1, 2, 3\}$. If this assumption is violated, the primal problem does not have any non-degenerate minimal solution, see (5.2) below. We denote (P_0) and (P) with weighted objective function by (P_0^σ) and (P^σ) , respectively.




FIG. 5.1. $\sigma_i > 0$ ($i = 1, 2, 3$) can be regarded as interface tensions.

THEOREM 5.1. *Let (X_0, \dots, X_3) be a non-degenerate minimal solution for (P_0^σ) . Then it is a minimum solution for (P^σ) , and satisfies (2.2) and*

$$(5.1) \quad \frac{\sin \theta_1}{\sigma_1} = \frac{\sin \theta_2}{\sigma_2} = \frac{\sin \theta_3}{\sigma_3},$$

where θ_i ($i = 1, 2, 3$) are defined as in Fig. 5.1.

Proof. The first assertion and (2.2) are proved in the same way with Theorem 3.1. As well as (3.1), we get $\sigma_1 n_1 + \sigma_2 n_2 + \sigma_3 n_3 = 0 \in R^n$. Hence we have

$$(5.2) \quad \frac{\sigma_k^2 - \sigma_i^2 - \sigma_j^2}{2\sigma_i\sigma_j} = n_i^T n_j = \cos \theta_k$$

for any $\{i, j, k\} = \{1, 2, 3\}$. So we get

$$\frac{\sin^2 \theta_k}{\sigma_k^2} = \frac{2(\sigma_1^2\sigma_2^2 + \sigma_2^2\sigma_3^2 + \sigma_3^2\sigma_1^2) - \sigma_1^4 - \sigma_2^4 - \sigma_3^4}{4\sigma_1^2\sigma_2^2\sigma_3^2}.$$

Since the right-hand side does not depend on k , we get (5.1). \square

LEMMA 5.2. *When Ω is a triangle in R^2 , it holds that*

$$\min(P^\sigma) = \min(P_0^\sigma) = \min_{i=1,2,3} \sigma_i h_i,$$

where h_i is the height of the triangle with base length a_i .

Proof. It's enough to consider KKT condition for the following extremal problem.

$$\begin{aligned} \text{Minimize} \quad & \sigma_1 l_1 + \sigma_2 l_2 + \sigma_3 l_3 \\ \text{subject to} \quad & a_1 l_1 + a_2 l_2 + a_3 l_3 = 2S \\ & l_i \geq 0 \quad (i = 1, 2, 3). \end{aligned}$$

\square

For a triangular cylinder, let Δ be a triangle defined by (4.3) and σ_i ($i = 1, 2, 3$) positive real numbers. We call the smallest weighted height of Δ the weighted width of the triangular cylinder. Then the dual problem (D^σ) is defined as follows.

Maximize the weighted width of a triangular cylinder that separates $\{C_i\}_{i=1}^3$.

THEOREM 5.3. *If (P_0^σ) (or (P^σ)) has a non-degenerate minimal solution, then it holds that*

$$(5.3) \quad \min(P_0^\sigma) = \min(P^\sigma) = \max(D^\sigma).$$

Proof. Since the proof of this theorem is the same as that of Theorem 4.3 except obvious modifications, we omit it. \square

6. Concluding remark. Gale and Klee [3] gave separation theorems for an infinite number of convex sets with empty intersection in R^n . For example, if a family of open convex proper subsets $\{C_\nu\}$ of R^n has empty intersection, then there are open halfspaces $\{H_\nu^-\}$ with empty intersection such that $C_\nu \subset H_\nu^-$ for any ν , see also a survey paper Klee [8] and Bair [1]. Since their separation theorems are not quantitative but qualitative, they do not seem to derive us to our duality theorems.

Further, a generalization of Duboviskii and Miljutin's theorem [2] asserts that if a finite number of convex sets $\{C_i\}_{i=1}^k$ in R^n has empty intersection, then there exist non-trivial vectors $\xi_i \in R^n$ ($i = 1, \dots, k$) such that

$$(6.1) \quad \xi_1 + \dots + \xi_k = 0$$

and

$$(6.2) \quad \delta^*(\xi_1|C_1) + \dots + \delta^*(\xi_k|C_k) \leq 0,$$

where $\delta^*(\xi_i|C_i)$ denotes the support function $\sup\{\xi_i^T X_i \mid X_i \in C_i\}$. If we directly apply this separation theorem to the linear case $C_i = \{X_i \mid \xi_i^T X_i \leq \alpha_i\}$, (6.1) and (6.2) reduce to $\sum_{i=1}^3 \lambda_i \xi_i = 0$ and $\sum_{i=1}^3 \lambda_i \alpha_i \leq 0$ for some non-trivial $\lambda_i \geq 0$ ($i = 1, 2, 3$), respectively. They do not seem enough to derive our duality theorems, either.

REFERENCES

- [1] J. Bair, *Sur la Séparation de familles finites d'ensembles convexes*, Bulletin de la Société Royale des Sciences de Liège, 41 (1972), pp. 281–291.
- [2] A. Ja. Duboviskii and A. A. Miljutin, *Extremal problems with constraints*, Soviet Math. Dokl. 4 (1963), pp. 452–255.
- [3] D. Gale and V. Klee, *Continuous convex sets*, Mathematics Scandinavica, 7 (1959), pp. 379–391.
- [4] R. Ikota and E. Yanagida, *A stability criterion for stationary curves to the curvature-driven motion with a triple junction*, Differential and Integral Equations, 16 (2003), pp. 707–726.
- [5] H. Kawasaki, *A game-theoretic aspect of conjugate sets for a nonlinear programming problem*, in Proceedings of the third International Conference on Nonlinear Analysis and Convex Analysis, Yokohama Publishers, (2004), pp. 159–168.
- [6] H. Kawasaki, *Conjugate-set game for a nonlinear programming problem*, in *Game theory and applications 10*, eds. L.A. Petrosjan and V.V. Mazalov, Nova Science Publishers, New York, USA, (2005), pp. 87–95.
- [7] H. Kawasaki, *A duality theorem for a three-phase partition problem*, submitted.
- [8] V. Klee, *Separation and support properties of convex sets—A survey*, in *Control Theory and the Calculus of Variations*, ed. A. V. Balakrishnan, Academic Press, New York, (1969), pp. 235–303.
- [9] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, Princeton, New Jersey, (1970).
- [10] P. Sternberg and W. P. Zeimer, *Local minimizers of a three-phase partition problem with triple junctions*, Proc. Royal Soc. Edin., 124A (1994), pp. 1059–1073.

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Hypergeometric solutions to the q -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE
Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI
Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI
Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA
Ergodicity and exponential β -mixing bounds for a strong solution of Lévy-driven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI
The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI
Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI
On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA
Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV
Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS
Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA
The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON
Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON
Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER
Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift

- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations
- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA
Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS
Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI
Nonlinear regression modeling via regularized wavelets and smoothing parameter selection

- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO,
Yoshihiro MIZOGUCHI & Yasuo KAWAHARA
On reversible cellular automata with finite cell array
- MHF2005-28 Toru KOMATSU
Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU
A computational approach to constructive a priori and a posteriori error
estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO
Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI
A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI
A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI
An explicit formula of the Shapley value for a cooperative game induced from
the conjugate point
- MHF2005-34 Hideki MURAKAWA
A regularization of a reaction-diffusion system approximation to the two-phase
Stefan problem
- MHF2006-1 Masahisa TABATA
Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite
element scheme
- MHF2006-2 Ken-ichi MARUNO & G R W QUISPEL
Construction of integrals of higher-order mappings
- MHF2006-3 Setsuo TANIGUCHI
On the Jacobi field approach to stochastic oscillatory integrals with quadratic
phase function
- MHF2006-4 Kouji HASHIMOTO, Kaori NAGATOU & Mitsuhiro T. NAKAO
A computational approach to constructive a priori error estimate for finite
element approximations of bi-harmonic problems in nonconvex polygonal
domains
- MHF2006-5 Hidefumi KAWASAKI
A duality theory based on triangular cylinders separating three convex sets in
 R^n