九州大学学術情報リポジトリ Kyushu University Institutional Repository

Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

Kagei, Yoshiyuki Faculty of Mathematics, Kyushu University

https://hdl.handle.net/2324/3392

出版情報:Funkcialaj Ekvacioj. 50 (2), pp.287-337, 2007-08-14. 日本数学会函数方程式論分科会 バージョン: 権利関係:(c) 2007 the Division of Functional Equations, The Mathematical Society of Japan

MHF Preprint Series

Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

Y. Kagei

MHF 2006-22

(Received June 6, 2006)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN

Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

Yoshiyuki KAGEI

Faculty of Mathematics, Kyushu University, Fukuoka 812-8581, Japan

Abstract

The resolvent problem of the linearized compressible Navier-Stokes equation around a given constant state is considered in an infinite layer $\mathbf{R}^{n-1} \times (0, a)$, $n \geq 2$, under the no slip boundary condition for the momentum. It is proved that the linearized operator is sectorial in $W^{1,p} \times L^p$ for $1 . The <math>L^p$ estimates for the resolvent are established for all $1 \leq p \leq \infty$. The estimates for the *high frequency part* of the resolvent are also derived, which lead to the exponential decay of the corresponding part of the semigroup.

1. Introduction

This paper studies the following resolvent problem

(1.1)
$$(\lambda + L)u = f$$

in an infinite layer $\Omega = \mathbf{R}^{n-1} \times (0, a), n \geq 2$, where $\lambda \in \mathbf{C}$ is a parameter, f = f(x) is a given function with values in $\mathbf{R}^{n+1}, u = \begin{pmatrix} \phi \\ m \end{pmatrix}$ is the unknown function with $\phi = \phi(x) \in \mathbf{R}$ and $m = {}^{T}(m^{1}(x), \cdots, m^{n}(x)) \in \mathbf{R}^{n}$, and L is an operator defined by

$$L = \begin{pmatrix} 0 & \gamma \operatorname{div} \\ \\ \gamma \nabla & -\nu \Delta I_n - \tilde{\nu} \nabla \operatorname{div} \end{pmatrix}$$

with positive constants ν and γ and a nonnegative constant $\tilde{\nu}$. Here $x = \begin{pmatrix} x' \\ x_n \end{pmatrix} \in \Omega$ with $x' \in \mathbf{R}^{n-1}$, $x_n \in (0, a)$; the superscript $^T \cdot$ stands for the transposition; I_n is the $n \times n$ identity matrix; and div, ∇ and Δ are the

usual divergence, gradient and Laplacian with respect to x. We consider (1.1) under the boundary condition

(1.2)
$$m|_{\partial\Omega} = 0.$$

Problem (1.1)–(1.2) is the resolvent problem associated with the linearization of the compressible Navier-Stokes equation around a motionless state with a positive constant density, where ϕ and m are the Laplace transform of the perturbation of the density and the momentum, respectively.

In this paper we establish the L^p estimates for the solution of (1.1)-(1.2)for all $1 \leq p \leq \infty$. The estimates show that -L generates an analytic semigroup in $W^{1,p} \times L^p$ for 1 . We also establish the estimatesfor the*high frequency part*of the resolvent, which lead to the exponentialdecay of the corresponding part of the semigroup. The*low frequency part*isinvestigated in [5]. The precise statement of the main results of this paperwill be given in section 2.

The resolvent problem in an infinite layer was studied in [1, 2, 3] in the case of the incompressible Stokes equation. They established the L^p estimates of the resolvent for $1 , which yields the exponential decay of the Stokes semigroup in <math>L^p$ norms. To analyze the resolvent, they considered the Fourier transform in $x' \in \mathbb{R}^{n-1}$ and derived solution formulae for the resolvent problem. The L^p estimates were then obtained by applying the Fourier multiplier theorem.

In this paper we will also consider the Fourier transform $(\lambda + \hat{L}_{\xi'})^{-1}$ of the resolvent in $x' \in \mathbf{R}^{n-1}$, where $\xi' \in \mathbf{R}^{n-1}$ denotes the phase space variable. In section 3 we derive an integral representation of $(\lambda + \hat{L}_{\xi'})^{-1}$. In the derivation we make use of the invariance of (1.1) under the orthogonal group and Green's formula for (1.1), which naturally leads to a decomposition of the solution into the two parts; one is the solution under the slip boundary condition; and the other is the term arising from the viscous friction stress due to the no slip boundary condition (1.2). (Cf. the solution formula for the half-space problem given in [6, 7].) In section 4 we investigate the resolvent problem based on the integral representation. The L^p estimates for 1are obtained by applying the Fourier multiplier theorem as in [1, 2, 3]. We obtain the L^p estimates for $p = 1, \infty$ based on the Riemann-Lebesgue lemma as in the analysis of the Cauchy problem for (1.1) given in [8]. We also establish the estimates for the high frequency part $|\xi'| >> 1$. In contrast to the incompressible Stokes problem, $L_{\xi'}$ has different characters between the cases $|\xi'| >> 1$ and $|\xi'| << 1$. The spectral analysis near the origin is given in [5]. It is shown in [5] that the continuous spectrum reaches the origin $\lambda = 0.$

2. Main Results

We first introduce some notation which will be used throughout the paper. For a domain D and $1 \leq p \leq \infty$ we denote by $L^p(D)$ the usual Lebesgue space on D and its norm is denoted by $\|\cdot\|_{L^p(D)}$. Let ℓ be a nonnegative integer. The symbol $W^{\ell,p}(D)$ denotes the ℓ th order L^p Sobolev space on D with norm $\|\cdot\|_{W^{\ell,p}(D)}$. When p = 2, the space $W^{\ell,2}(D)$ is denoted by $H^{\ell}(D)$ and its norm is denoted by $\|\cdot\|_{H^{\ell}(D)}$. $C_0^{\ell}(D)$ stands for the set of all C^{ℓ} functions which have compact support in D. We denote by $W_0^{1,p}(D)$ the completion of $C_0^1(D)$ in $W^{1,p}(D)$. In particular, $W_0^{1,2}(D)$ is denoted by $H_0^1(D)$.

We simply denote by $L^p(D)$ (resp., $W^{\ell,p}(D)$, $H^\ell(D)$) the set of all vector fields $m = {}^T(m^1, \dots, m^n)$ on D with $m^j \in L^p(D)$ (resp., $W^{\ell,p}(D)$, $H^\ell(D)$), $j = 1, \dots, n$, and its norm is also denoted by $\|\cdot\|_{L^p(D)}$ (resp., $\|\cdot\|_{W^{\ell,p}(D)}$, $\|\cdot\|_{H^\ell(D)}$). For $u = \begin{pmatrix} \phi \\ m \end{pmatrix}$ with $\phi \in W^{k,p}(D)$ and $m = {}^T(m^1, \dots, m^n) \in$ $W^{\ell,q}(D)$, we define $\|u\|_{W^{k,p}(D) \times W^{\ell,q}(D)}$ by $\|u\|_{W^{k,p}(D) \times W^{\ell,q}(D)} = \|\phi\|_{W^{k,p}(D)} +$ $\|m\|_{W^{\ell,q}(D)}$. When $k = \ell$ and p = q, we simply write $\|u\|_{W^{k,p}(D) \times W^{k,p}(D)} =$ $\|u\|_{W^{k,p}(D)}$.

In case $D = \Omega$ we abbreviate $L^p(\Omega)$ (resp., $W^{\ell,p}(\Omega)$, $H^{\ell}(\Omega)$) as L^p (resp., $W^{\ell,p}$, H^{ℓ}). In particular, the norm $\|\cdot\|_{L^p(\Omega)} = \|\cdot\|_{L^p}$ is denoted by $\|\cdot\|_p$.

In case D = (0, a) we denote the norm of $L^p(0, a)$ by $|\cdot|_p$. The inner product of $L^2(0, a)$ is denoted by

$$(f,g) = \int_0^a f(x_n)\overline{g(x_n)} \, dx_n, \quad f,g \in L^2(0,a).$$

Here \overline{g} denotes the complex conjugate of g. We will also denote the bilinear pairing f and g by

$$((f,g)) = \int_0^a f(x_n)g(x_n)\,dx_n.$$

The norms of $W^{\ell,p}(0,a)$ and $H^{\ell}(0,a)$ are denoted by $|\cdot|_{W^{\ell,p}}$ and $|\cdot|_{H^{\ell}}$, respectively.

We often write $x \in \Omega$ as $x = \begin{pmatrix} x' \\ x_n \end{pmatrix}$, $x' = {}^T(x_1, \cdots, x_{n-1}) \in \mathbf{R}^{n-1}$.

Partial derivatives of a function u in x, x', x_n and t are denoted by $\partial_x u$, $\partial_{x'} u$, $\partial_{x_n} u$ and $\partial_t u$, respectively. We also write higher order partial derivatives of u in x as $\partial_x^k u = (\partial_x^{\alpha} u; |\alpha| = k)$.

We denote the $k \times k$ identity matrix by I_k . In particular, when k = n+1, we simply write I for I_{n+1} . We also define $(n+1) \times (n+1)$ diagonal matrices

 Q_0, \tilde{Q}, Q' and Q_n by

$$Q_{0} = \operatorname{diag} (1, 0, \dots, 0, 0), \quad \widetilde{Q} = \operatorname{diag} (0, 1, \dots, 1, 1),$$
$$Q' = \operatorname{diag} (0, 1, \dots, 1, 0), \quad Q_{n} = \operatorname{diag} (0, 0, \dots, 0, 1).$$
For $u = \begin{pmatrix} \phi \\ m \end{pmatrix} \in \mathbf{R}^{n+1}$ with $m = \begin{pmatrix} m' \\ m^{n} \end{pmatrix} \in \mathbf{R}^{n}$, we have
$$Q_{0}u = \begin{pmatrix} \phi \\ 0 \end{pmatrix}, \quad \widetilde{Q}u = \begin{pmatrix} 0 \\ m \end{pmatrix}, \quad Q'u = \begin{pmatrix} 0 \\ m' \\ 0 \end{pmatrix}, \quad Q_{n}u = \begin{pmatrix} 0 \\ 0 \\ m^{n} \end{pmatrix}.$$

We next introduce some notation about integral operators. For a function f = f(z) ($z \in \mathbf{R}^k$), we denote its Fourier transform by $\mathscr{F}_z f$:

$$(\mathscr{F}_z f)(\zeta) = \int_{\mathbf{R}^k} f(z) e^{-i\zeta \cdot z} dz$$

In particular, when k = n - 1 $(x' \in \mathbf{R}^{n-1})$, we denote $\mathscr{F}_{x'}f$ by \hat{f} ,

$$\widehat{f} = (\mathscr{F}_{x'}f)(\xi') = \int_{\mathbf{R}^{n-1}} f(x')e^{-i\xi'\cdot x'} \, dx'.$$

The inverse Fourier transform is denoted by \mathscr{F}_{ζ}^{-1} :

$$(\mathscr{F}_{\zeta}^{-1}f)(z) = (2\pi)^{-k} \int_{\mathbf{R}^k} f(\zeta) e^{i\zeta \cdot z} \, d\zeta.$$

For a function $K(x_n, y_n)$ on $(0, a) \times (0, a)$ we will denote by Kf the integral operator $\int_0^a K(x_n, y_n) f(y_n) dy_n$. Similarly, for a function $K(x', x_n, y_n)$ on $\mathbf{R}^{n-1} \times (0, a) \times (0, a)$ we will denote by Kf the integral operator $\int_{\mathbf{R}^{n-1}} \int_0^a K(x'-y', x_n, y_n) f(y_n) dy' dy_n$.

We denote the resolvent set of a closed operator A by $\rho(A)$ and the spectrum of A by $\sigma(A)$. For $\Lambda \in \mathbf{R}$ and $\theta \in (\frac{\pi}{2}, \pi)$ we will denote the set $\{\lambda \in \mathbf{C}; |\arg(\lambda - \Lambda)| \leq \theta\}$ by $\Sigma(\Lambda, \theta)$:

$$\Sigma(\Lambda, \theta) = \{\lambda \in \mathbf{C}; |\arg(\lambda - \Lambda)| \le \theta\}.$$

We now state the main results of this paper.

Theorem 2.1. Let $1 . There exists a number <math>\theta \in (\frac{\pi}{2}, \pi)$ such that for any $\eta > 0$ problem (1.1)–(1.2) has a unique solution $u \in W^{1,p} \times$

 $(W^{2,p} \cap W_0^{1,p})$ for any $f \in W^{1,p} \times L^p$, provided that $\lambda \in \Sigma(\eta, \theta)$. Furthermore, $u = (\lambda + L)^{-1}f$ satisfies the following estimates uniformly in $\lambda \in \Sigma(\eta, \theta)$:

$$\|\partial_x^k (\lambda + L)^{-1} f\|_p \le C \left\{ \frac{\|Q_0 f\|_{W^{k,p}}}{|\lambda| + 1} + \frac{\|\widetilde{Q} f\|_p}{(|\lambda| + 1)^{1 - \frac{k}{2}}} \right\}, \quad k = 0, 1,$$

and

$$\|\partial_x^2 \widetilde{Q}(\lambda + L)^{-1} f\|_2 \le C \|f\|_{W^{1,p} \times L^p}.$$

Furthermore, if $\tilde{Q}f\Big|_{x_n=0,a}=0$, then there holds

$$\|\partial_x \widetilde{Q}(\lambda+L)^{-1}f\|_p \le \frac{C}{|\lambda|+1} \|f\|_{W^{1,p}}.$$

In the application to the nonlinear problem we will also use the following $H^s \to L^\infty$ estimates of the resolvent.

Theorem 2.2. Let θ be the number given in Theorem 2.1 and let η be a positive number. Then the following estimates hold uniformly in $\lambda \in \Sigma(\eta, \theta)$:

$$\|\partial_x^k Q_0(\lambda+L)^{-1}f\|_{\infty} \le C \left\{ \frac{\|Q_0f\|_{H^{[\frac{n}{2}]+1+k}}}{|\lambda|+1} + \frac{\|\tilde{Q}f\|_{H^{[\frac{n}{2}]+k}}}{(|\lambda|+1)^{\frac{3}{4}}} \right\}, \quad k = 0, 1,$$

and

$$\|\partial_x^k \widetilde{Q}(\lambda+L)^{-1} f\|_{\infty} \le C \left\{ \frac{\|Q_0 f\|_{H^{[\frac{n}{2}]+k}}}{(|\lambda|+1)^{\frac{3}{4}}} + \frac{\|\widetilde{Q} f\|_{H^{[\frac{n}{2}]-1+k}}}{(|\lambda|+1)^{\frac{\varepsilon}{4}}} \right\}, \quad k = 0, 1.$$

Here ε is some number satisfying $0 < \varepsilon < \frac{1}{3}$.

As for the L^p estimates for $p = 1, \infty$, we have the following result.

Theorem 2.3. Let θ be the number given in Theorem 2.1 and let η be a positive number. Then the following estimates hold uniformly in $\lambda \in \Sigma(\eta, \theta)$: for $p = 1, \infty$,

$$\|\partial_x^k Q_0(\lambda+L)^{-1}f\|_p \le \frac{C}{|\lambda|+1} \|f\|_{W^{k+1,p} \times W^{k,p}}, \quad k = 0, 1,$$

and

$$\|\partial_x^k \widetilde{Q}(\lambda+L)^{-1} f\|_p \le C \left\{ \frac{\|Q_0 f\|_{W^{k,p}}}{|\lambda|+1} + \frac{\|\widetilde{Q} f\|_p}{(|\lambda|+1)^{1-\frac{k}{2}}} \right\}, \quad k = 0, 1.$$

Furthermore, if $\tilde{Q}f\Big|_{x_n=0,a}=0$, then, there hold, for $p=1,\infty$,

$$\|\partial_x \widetilde{Q}(\lambda + L)^{-1} f\|_p \le \frac{C}{|\lambda| + 1} \|f\|_{W^{1,p}}.$$

We see from Theorem 2.1 that -L generates the analytic semigroup $\mathscr{U}(t)$ in $W^{1,p} \times L^p$ for 1 . Based on Theorems 2.1–2.3 we have the following $estimates of <math>\mathscr{U}(t)$ for $0 < t \leq 1$.

Corollary 2.4. Let $\ell = 0, 1$. Then there hold the estimates

$$\begin{aligned} \|\partial_x^{\ell} \mathscr{U}(t) u_0\|_p &\leq Ct^{-\frac{\ell}{2}} \|u_0\|_{W^{\ell,p} \times L^p}, \quad 1$$

and

$$\|\partial_x^\ell \mathscr{U}(t)u_0\|_p \le Ct^{-\frac{\ell}{2}} \|u_0\|_{W^{\ell+1,p} \times W^{\ell,p}}, \quad p = 1, \infty,$$

for $0 < t \leq 1$ with some constant $0 < \varepsilon < 1$, provided that u_0 belongs to the Sobolev spaces indicated on the right-hand side of each inequality above. Furthermore, if $\tilde{Q}u_0|_{x_n=0,a} = 0$, then

$$\|\partial_x \mathscr{U}(t)u_0\|_1 \le C \|u_0\|_{W^{2,1} \times W^{1,1}}$$

holds for $0 \le t \le 1$.

To investigate problem (1.1)–(1.2) we consider the Fourier transform in $x' \in \mathbf{R}^{n-1}$. Applying the Fourier transform, we have the following boundary value problem for functions $\phi(x_n)$ and $m(x_n)$ on the interval (0, a):

(2.1)
$$\lambda u + \hat{L}_{\xi'} u = f,$$

where $u = \begin{pmatrix} \phi(x_n) \\ m'(x_n) \\ m^n(x_n) \end{pmatrix}$, $f = \begin{pmatrix} f^0(x_n) \\ f'(x_n) \\ f^n(x_n) \end{pmatrix}$ and $\hat{L}_{\xi'}$ is a closed operator on $H^1(0, a) \times L^2(0, a)$ defined by $\hat{L}_{\xi'} = \hat{A}_{\xi'} + \hat{B}_{\xi'}$ with domain of definition

 $H^{1}(0,a) \times L^{2}(0,a)$ defined by $L_{\xi'} = A_{\xi'} + B_{\xi'}$ with domain of definition $D(\hat{L}_{\xi'}) = H^{1}(0,a) \times (H^{2}(0,a) \cap H^{1}_{0}(0,a)).$ Here

$$\widehat{A}_{\xi'} = \begin{pmatrix} 0 & 0 & 0 \\ 0 & \nu(|\xi'|^2 - \partial_{x_n}^2)I_{n-1} + \widetilde{\nu}\xi'^T\xi' & -i\widetilde{\nu}\xi'\partial_{x_n} \\ 0 & -i\widetilde{\nu}^T\xi'\partial_{x_n} & \nu(|\xi'|^2 - \partial_{x_n}^2) - \widetilde{\nu}\partial_{x_n}^2 \end{pmatrix}$$

and

$$\widehat{B}_{\xi'} = \begin{pmatrix} 0 & i\gamma^T \xi' & \gamma \partial_{x_n} \\ i\gamma \xi' & 0 & 0 \\ \gamma \partial_{x_n} & 0 & 0 \end{pmatrix}.$$

In the analysis of the semigroup generated by -L, it is convenient to decompose the phase space $\{\xi' \in \mathbf{R}^{n-1}\}$ into the parts $|\xi'| << 1$ and $|\xi'| >> 1$, since $\hat{L}_{\xi'}$ has different characters between the cases $|\xi'| << 1$ and $|\xi'| >> 1$. Motivated by this we introduce the following decomposition. Let r be a positive number. We take a function $\chi(\xi') \in C^{\infty}(\mathbf{R}^{n-1})$ satisfying $0 \le \chi \le 1$ on \mathbf{R}^{n-1} , $\chi(\xi') = 0$ for $|\xi'| \le \frac{r}{2}$ and $\chi(\xi') = 1$ for $|\xi'| \ge r$. We define the operator $R^{(1)}(\lambda)$ by

$$R^{(1)}(\lambda)f = \mathscr{F}_{\xi'}^{-1}\left[\chi(\xi')(\lambda + \widehat{L}_{\xi'})^{-1}\widehat{f}\right].$$

The estimates in Theorems 2.1–2.3 hold for a negative η with $(\lambda + L)^{-1}$ replaced by $R^{(1)}(\lambda)$.

Theorem 2.5. Let r be a positive number.

(i) There exist positive numbers $\tilde{\eta}$ and $\tilde{\theta}$ with $\tilde{\theta} \in (\frac{\pi}{2}, \pi)$ such that $\Sigma(-\tilde{\eta}, \tilde{\theta}) \subset \rho(-\hat{L}_{\xi'})$ for $|\xi'| \geq r$.

(ii) Let $1 and define <math>R^{(1)}(\lambda)$ as above. Then the following estimates hold uniformly in $\lambda \in \Sigma(-\tilde{\eta}, \tilde{\theta})$:

$$\|\partial_x^k R^{(1)}(\lambda)f\|_p \le \left\{\frac{\|Q_0 f\|_{W^{k,p}}}{|\lambda|+1} + \frac{\|\tilde{Q}f\|_p}{(|\lambda|+1)^{1-\frac{k}{2}}}\right\}, \quad k = 0, 1,$$

and

$$\|\partial_x^2 \widetilde{Q} R^{(1)}(\lambda) f\|_2 \le C \|f\|_{W^{1,p} \times L^p}.$$

Theorem 2.6 Let $\tilde{\eta}$ and $\tilde{\theta}$ be the numbers as in Theorem 2.5. Then the following estimates hold uniformly in $\lambda \in \Sigma(-\tilde{\eta}, \tilde{\theta})$:

$$\|\partial_x^k Q_0 R^{(1)}(\lambda) f\|_{\infty} \le C \left\{ \frac{\|Q_0 f\|_{H^{[\frac{n}{2}]+1+k}}}{|\lambda|+1} + \frac{\|\tilde{Q}f\|_{H^{[\frac{n}{2}]+k}}}{(|\lambda|+1)^{\frac{3}{4}}} \right\}, \quad k = 0, 1,$$

and

$$\|\partial_x^k \widetilde{Q} R^{(1)}(\lambda) f\|_{\infty} \le C \left\{ \frac{\|Q_0 f\|_{H^{[\frac{n}{2}]+k}}}{(|\lambda|+1)^{\frac{3}{4}}} + \frac{\|\widetilde{Q} f\|_{H^{[\frac{n}{2}]-1+k}}}{(|\lambda|+1)^{\frac{\varepsilon}{4}}} \right\}, \quad k = 0, 1.$$

Here ε is some number satisfying $0 < \varepsilon < \frac{1}{3}$.

Theorem 2.7. Let $p = 1, \infty$. Let $\tilde{\eta}$ and $\tilde{\theta}$ be the numbers as in Theorem 2.5. Then the following estimates hold uniformly in $\lambda \in \Sigma(-\tilde{\eta}, \tilde{\theta})$:

$$\|\partial_x^k Q_0 R^{(1)}(\lambda) f\|_p \le \frac{C}{|\lambda|+1} \|f\|_{W^{k+1,p} \times W^{k,p}}, \quad k = 0, 1,$$

and

$$\|\partial_x^k \widetilde{Q}R^{(1)}(\lambda)f\|_p \le C\left\{\frac{\|Q_0 f\|_{W^{k,p}}}{|\lambda|+1} + \frac{\|\widetilde{Q}f\|_p}{(|\lambda|+1)^{1-\frac{k}{2}}}\right\}, \quad k = 0, 1.$$

3. An integral representation of the resolvent

In this section we derive an integral representation of the resolvent. For this purpose we take the Fourier transform of (1.1)–(1.2) in $x' \in \mathbb{R}^{n-1}$ and consider the boundary value problem (2.1).

We begin with the following observation on the resolvent of $-\hat{L}_{\xi'}$. If $\lambda \neq 0$, then, by the first row of equation (2.1), ϕ is written as

(3.1)
$$\phi = \frac{1}{\lambda} \left\{ f^0 - i\gamma\xi' \cdot m' - \gamma\partial_{x_n}m^n \right\}.$$

Substituting (3.1) into the second and third rows of (2.1), we obtain

(3.2)
$$\mathcal{L}(\lambda,\xi')m = F,$$

where
$$\mathcal{L}(\lambda,\xi') = \mathcal{A}(\lambda,\xi') + \mathcal{B}(\lambda,\xi')$$
 with domain of definition $D(\tilde{\mathcal{L}}(\lambda,\xi')) = H^2(0,a) \cap H^1_0(0,a)$ and $F = \lambda \begin{pmatrix} f' \\ f^n \end{pmatrix} - \gamma \begin{pmatrix} i\xi' \\ \partial_{x_n} \end{pmatrix} f^0$. Here
 $\mathcal{A}(\lambda,\xi') = \begin{pmatrix} \{\lambda^2 + \nu\lambda(|\xi'|^2 - \partial_{x_n}^2)\}I_{n-1} & 0 \\ 0 & \lambda^2 + \nu\lambda(|\xi'|^2 - \partial_{x_n}^2) \end{pmatrix},$
 $\mathcal{B}(\lambda,\xi') = (\tilde{\nu}\lambda + \gamma^2) \begin{pmatrix} \xi'^T\xi' & -i\xi'\partial_{x_n} \\ -i^T\xi'\partial_{x_n} & -\partial_{x_n}^2 \end{pmatrix}.$

We thus deduce that (2.1) is equivalent to (3.1) and (3.2) if $\lambda \neq 0$. We also write (3.2) as

(3.3)
$$\hat{\mathcal{L}}(\lambda,\xi')m = \hat{F},$$

where

$$\widetilde{\mathcal{L}}(\lambda,\xi') = -\partial_{x_n}^2 I_n + \widetilde{\mathcal{A}}(\lambda,\xi') + \widetilde{\mathcal{B}}(\lambda,\xi')\partial_{x_n}$$

with domain of definition $D(\tilde{\mathcal{L}}(\lambda,\xi')) = H^2(0,a) \cap H^1_0(0,a)$ and $\tilde{F} = \begin{pmatrix} \frac{1}{\nu\lambda}F' \\ \frac{1}{\nu_1\lambda+\gamma^2}F^n \end{pmatrix}$. Here

$$\widetilde{\mathcal{A}}(\lambda,\xi') = \begin{pmatrix} \frac{1}{\nu}(\lambda+\nu|\xi'|^2)I_{n-1} + \frac{1}{\nu\lambda}(\widetilde{\nu}\lambda+\gamma^2)\xi'^T\xi' & 0\\ 0 & \frac{\lambda(\lambda+\nu|\xi'|^2)}{\nu_1\lambda+\gamma^2} \end{pmatrix}$$

and

$$\widetilde{\mathcal{B}}(\lambda,\xi') = \begin{pmatrix} 0 & -i\frac{\widetilde{\nu}\lambda+\gamma^2}{\nu\lambda}\xi' \\ -i\frac{\widetilde{\nu}\lambda+\gamma^2}{\nu_1\lambda+\gamma^2}T\xi' & 0 \end{pmatrix}.$$

Here and in what follows we write ν_1 for $\nu + \tilde{\nu}$:

$$\nu_1 = \nu + \widetilde{\nu}.$$

Lemma 3.1. Assume that $\lambda \neq 0$ and $\nu_1 \lambda + \gamma^2 \neq 0$. Then $\lambda \in \rho(-\hat{L}_{\xi'})$ if and only if Ker $\mathcal{L}(\lambda, \xi') = \{0\}$.

Proof. Suppose that $\lambda \in \rho(-\hat{L}_{\xi'})$. Let $m \in H^2(0, a) \cap H^1_0(0, a)$ satisfy $\mathcal{L}(\lambda, \xi')m = 0$, namely, $\tilde{\mathcal{L}}(\lambda, \xi')m = 0$. We define ϕ as in (3.1) with $f^0 = 0$ and set $u = \begin{pmatrix} \phi \\ m \end{pmatrix}$. It then follows that $(\lambda + \hat{L}_{\xi'})u = 0$. Since $\lambda \in \rho(-\hat{L}_{\xi'})$, we see that u = 0, in particular, m = 0. This shows that Ker $\mathcal{L}(\lambda, \xi') = \{0\}$. Conversely, suppose that Ker $\mathcal{L}(\lambda, \xi') = \{0\}$, namely, Ker $\tilde{\mathcal{L}}(\lambda, \xi') = \{0\}$. We define \tilde{F} as in (3.3). Since (2.1) is equivalent to (3.1) and (3.3), it suffices to show the unique existence of the solution $u = \begin{pmatrix} \phi \\ m \end{pmatrix}$ of (3.1) and (3.3) satisfying $|u|_{H^1 \times H^2} \leq C|f|_{H^1 \times L^2}$ with some C > 0. Since Ker $\tilde{\mathcal{L}}(\lambda, \xi') = \{0\}$, the standard theory of elliptic equations shows that there exists a unique solution $m \in H^2(0, a) \cap H^1_0(0, a)$ of (3.3) and m satisfies

$$|m|_{H^2} \le C(\lambda,\xi') \left| \widetilde{F} \right|_2 \le C(\lambda,\xi') |f|_{H^1 \times L^2}.$$

It then follows from (3.1) that $\phi \in H^1(0, a)$ and $|\phi|_{H^1} \leq C(\lambda, \xi')|f|_{H^1 \times L^2}$. We thus conclude that $\lambda \in \rho(-\hat{L}_{\xi'})$. This completes the proof.

We next give a fundamental set of solutions of the ordinary differential equation (3.2) with F = 0. For this purpose we introduce some notation. We set

$$\lambda_{1,0} = -\nu |\xi'|^2$$

and

$$\lambda_{\pm,0} = -\frac{\nu_1}{2} |\xi'|^2 \pm \frac{1}{2} \sqrt{\nu_1^2 |\xi'|^4 - 4\gamma^2 |\xi'|^2},$$

and define

$$\mu_1 = \mu_1(\lambda, |\xi'|^2) = \sqrt{\frac{\lambda + \nu |\xi'|^2}{\nu}}$$

and

$$\mu_2 = \mu_2(\lambda, |\xi'|^2) = \sqrt{\frac{\lambda^2 + \nu_1 |\xi'|^2 \lambda + \gamma^2 |\xi'|^2}{\nu_1 \lambda + \gamma^2}}.$$

Remark 3.2. We observe that $\mu_1 = \sqrt{\frac{\lambda - \lambda_{1,0}}{\nu}}$ and $\mu_2 = \sqrt{\frac{(\lambda - \lambda_{+,0})(\lambda - \lambda_{-,0})}{(\nu_1 \lambda + \gamma^2)}}$. Furthermore, $\lambda_{-,0} = \overline{\lambda_{+,0}}$ with $\operatorname{Im} \lambda_{+,0} = \gamma |\xi'| \sqrt{1 - \frac{\nu_1^2}{4\gamma^2} |\xi'|^2}$ when $|\xi'| < 2\gamma/\nu_1$ and $\lambda_{\pm,0} \in \mathbf{R}$ when $|\xi'| > 2\gamma/\nu_1$, and it holds that

$$\lambda_{\pm,0} = -\frac{\nu_1}{2} |\xi'|^2 \pm i |\xi'| + O(|\xi'|^3)$$

as $|\xi'| \to 0$, and

$$\lambda_{+,0} = -\frac{\gamma^2}{\nu_1} + O(|\xi'|^{-2}), \quad \lambda_{-,0} = -\nu_1 |\xi'|^2 + O(1)$$

as $|\xi'| \to \infty$.

Proposition 3.3. Assume that $\lambda \neq 0$, $\tilde{\nu}\lambda + \gamma^2 \neq 0$, $\nu_1\lambda + \gamma^2 \neq 0$, $\lambda \neq \lambda_{1,0}$ and $\lambda \neq \lambda_{\pm,0}$. Then the following functions v_1, \dots, v_{2n} form a fundamental set of solutions of the ordinary differential equation in (3.2) with F = 0:

$$v_j(x_n) = \begin{pmatrix} \mathbf{e}'_j \cosh \mu_1 x_n \\ -i\frac{\xi_j}{\mu_1} \sinh \mu_1 x_n \end{pmatrix}, \quad v_n(x_n) = \begin{pmatrix} i\frac{\xi'}{\mu_2} \sinh \mu_2 x_n \\ \cosh \mu_2 x_n \end{pmatrix},$$
$$v_{n+j}(x_n) = \begin{pmatrix} \mathbf{e}'_j \mu_1 \sinh \mu_1 x_n \\ -i\xi_j \cosh \mu_1 x_n \end{pmatrix}, \quad v_{2n}(x_n) = \begin{pmatrix} i\xi' \cosh \mu_2 x_n \\ \mu_2 \sinh \mu_2 x_n \end{pmatrix},$$

where $j = 1, \dots, n-1$ and $e'_j = T(0, \dots, 1, \dots, 0) \in \mathbf{R}^{n-1}$.

Remark. We note that v_1, \dots, v_{2n} are analytic in λ and $|\xi'|^2$.

Proof. Setting $w = {}^{T}(w^{1}, \dots, w^{2n})$ with $w^{j} = m^{j}$, $w^{n+j} = \partial_{x_{n}}m^{j}$, $j = 1, \dots, n$, we see that the ordinary differential equation in (3.2) with F = 0 is equivalent to

(3.4)
$$\frac{dw}{dx_n} = \mathcal{M}(\lambda, \xi')w,$$

where

$$\mathcal{M}(\lambda,\xi') = \begin{pmatrix} 0 & I_n \\ \widetilde{\mathcal{A}}(\lambda,\xi') & \widetilde{\mathcal{B}}(\lambda,\xi') \end{pmatrix}$$

with $\widetilde{\mathcal{A}}(\lambda,\xi')$ and $\widetilde{\mathcal{B}}(\lambda,\xi')$ defined in (3.3).

To obtain a fundamental set of solutions of (3.4) we first consider the characteristic equation of $\mathcal{M}(\lambda, \xi')$. Let T' be an $(n-1) \times (n-1)$ orthogonal matrix and set

$$\mathcal{T} = \begin{pmatrix} T' & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & T' & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

It then follows that $\mathcal{M}(\lambda,\xi') = \mathcal{T}^{-1}\mathcal{M}(\lambda,T'\xi')\mathcal{T}$.

We take T' in such a way that $T'\xi' = |\xi'|e'_{n-1}$. With this T' we find that

$$\det (\mu I_{2n} - \mathcal{M}(\lambda, \xi')) = \det (\mu I_{2n} - \mathcal{M}(\lambda, T'\xi'))$$
$$= \det (\mu^2 I_n - \mu \widetilde{\mathcal{B}}(\lambda, T'\xi') - \widetilde{\mathcal{A}}(\lambda, T'\xi'))$$
$$= (\mu^2 - \mu_1^2)^{n-1}(\mu^2 - \mu_2^2).$$

Therefore, the eigenvalues of $\mathcal{M}(\lambda,\xi')$ are given by $\pm \mu_1$ and $\pm \mu_2$. Note that $\mu_1 \neq \pm \mu_2$ since $\lambda \neq 0$ and $\tilde{\nu}\lambda + \gamma^2 \neq 0$. Furthermore, $\mu_j \neq 0$, j=1,2, since $\lambda \neq \lambda_{1,0}$ and $\lambda \neq \lambda_{\pm,0}$.

We next look for eigenvectors associated with $\pm \mu_j$. To obtain eigenvectors for μ_1 we consider the problem

$$\mathcal{M}(\lambda,\xi')\begin{pmatrix} X\\ Y \end{pmatrix} = \mu_1\begin{pmatrix} X\\ Y \end{pmatrix},$$

where $X, Y \in \mathbb{C}^n$. This is equivalent to

$$(\widetilde{\mathcal{A}}(\lambda,\xi') + \mu_1 \widetilde{\mathcal{B}}(\lambda,\xi') - \mu_1^2 I_n) X = 0.$$

We write $X = \begin{pmatrix} X' \\ X_n \end{pmatrix}$. Since $\tilde{\nu}\lambda + \gamma^2 \neq 0$, we have $\xi' \cdot X' - i\mu_1 X_n = 0$. This implies that $X_{j,+} = \begin{pmatrix} \mu_1 \mathbf{e}'_j \\ -i\xi_j \end{pmatrix}$, $j = 1, \dots, n-1$, are eigenvectors for μ_1 . Similarly, one can see that $X_{j,-} = \begin{pmatrix} -\mu_1 \mathbf{e}'_j \\ -i\xi_j \end{pmatrix}$, $j = 1, \dots, n-1$, are eigenvectors for $-\mu_1$.

We next look for eigenvector for μ_2 . As in the case of μ_1 , it suffices to find a nontrivial solution $X \in \mathbb{C}^n$ of

$$(\widetilde{\mathcal{A}}(\lambda,\xi') + \mu_2 \widetilde{\mathcal{B}}(\lambda,\xi') - \mu_2^2 I_n) X = 0.$$

Since $\tilde{\nu}\lambda + \gamma^2 \neq 0$, it follows that

$$\frac{\lambda}{\nu_1 \lambda + \gamma^2} X_j + \frac{1}{\lambda} (\xi' \cdot X') \xi_j - i\mu_2 \frac{\xi_j}{\lambda} X_n = 0, \quad j = 1, \cdots, n-1, -i\mu_2 (\xi' \cdot X') - |\xi'|^2 X_n = 0.$$

This implies that $X_{n,+} = \begin{pmatrix} \xi' \\ -i\mu_2 \end{pmatrix}$ is an eigenvector for μ_2 . Similarly, we can find that $X_{n,-} = \begin{pmatrix} \xi' \\ i\mu_2 \end{pmatrix}$ is an eigenvector for $-\mu_2$.

Since $\mu_1 \neq \pm \mu_2$ and $\mu_j \neq 0$, j = 1, 2, we have a fundamental set of solutions of the ordinary differential equation in (3.2) with F = 0: $u_{j,\pm} = X_{j,\pm}e^{\pm\mu_1x_n}$ $(j = 1, \dots, n-1)$, $u_{n,\pm} = X_{n,\pm}e^{\pm\mu_2x_n}$. One can now obtain the basis v_1, \dots, v_{2n} by setting $v_j = \frac{1}{2\mu_1}(u_{j,+} - u_{j,-})$, $v_{n+j} = \frac{1}{2}(u_{j,+} + u_{j,-})$ $(j = 1, \dots, n-1)$, $v_n = \frac{i}{2\mu_2}(u_{n,+} - u_{n,-})$ and $v_{2n} = \frac{i}{2}(u_{n,+} + u_{n,-})$. This completes the proof.

We next give a characterization of the resolvent set $\rho(-\hat{L}_{\xi'})$. We define complex valued functions b_j (j = 1, 2, 3) by

$$b_1(\lambda, \xi', x_n) = b_1(\lambda, |\xi'|^2, x_n) = \cosh \mu_1 x_n - \cosh \mu_2 x_n,$$

$$b_2(\lambda, \xi', x_n) = b_2(\lambda, |\xi'|^2, x_n) = \mu_1 \sinh \mu_1 x_n - \frac{|\xi'|^2}{\mu_2} \sinh \mu_2 x_n,$$

$$b_3(\lambda, \xi', x_n) = b_3(\lambda, |\xi'|^2, x_n) = \mu_2 \sinh \mu_2 x_n - \frac{|\xi'|^2}{\mu_1} \sinh \mu_1 x_n,$$

with $\mu_j = \mu_j(\lambda, \xi'), \ j = 1, 2$. We set

$$D(\lambda,\xi') = D(\lambda,|\xi|^2) = b_3(\lambda,\xi',a)b_2(\lambda,\xi',a) + |\xi'|^2b_1(\lambda,\xi',a)^2.$$

In the following we will frequently abbreviate $b_j(\lambda, \xi', x_n)$ to $b_j(x_n)$. Note that b_j (j = 1, 2, 3) are analytic in λ and $|\xi'|^2$, and hence, so is D. We also set

$$\lambda_{1,k} = -\nu |\xi^{(k)}|^2$$

for $\xi' \in \mathbf{R}^{n-1}$ and $k = 1, 2, \cdots$, where

$$|\xi^{(k)}|^2 = |\xi'|^2 + a_k^2, \quad a_k = \frac{k\pi}{a}.$$

Lemma 3.4. Assume that $\lambda \neq 0$, $\tilde{\nu}\lambda + \gamma^2 \neq 0$, $\nu_1\lambda + \gamma^2 \neq 0$, $\lambda \neq \lambda_{\pm,0}$ and $\lambda \neq \lambda_{1,k}$ for any $k = 0, 1, 2, \cdots$. Then $\lambda \in \rho(-\hat{L}_{\xi'})$ if and only if $D(\lambda, \xi') \neq 0$. **Proof.** By Proposition 3.3 the solution of the equation in (3.2) with F = 0 is written as

$$m = c_1 v_1 + \dots + c_{2n} v_{2n}$$

for some $c_j \in \mathbf{C}$, $j = 1, \dots, 2n$. This *m* satisfies the boundary condition $m|_{x_n=0,a} = 0$ if and only if

(3.5)
$$A\begin{pmatrix}c_1\\\vdots\\c_{2n}\end{pmatrix} = 0 \text{ with } A = \begin{pmatrix}A_1(0) & A_2(0)\\A_1(a) & A_2(a)\end{pmatrix},$$

where $A_1(x_n) = (v_1(x_n), \dots, v_n(x_n))$ and $A_2(x_n) = (v_{n+1}(x_n), \dots, v_{2n}(x_n))$. Note that $A_1(0) = I_n$. In view of Lemma 3.1, $\lambda \in \rho(-\hat{L}_{\xi'})$ if and only if (3.5) has only the trivial solution $c_j = 0, j = 1, \dots, 2n$, namely, det $A \neq 0$.

Let us compute det A. Since $A_1(0) = I_n$, by a well known formula, we have

$$\det A = \det (A_2(a) - A_1(a)A_2(0)).$$

A direct calculation gives

$$A_{2}(a) - A_{1}(a)A_{2}(0)$$

$$= \begin{pmatrix} \mu_{1} \sinh \mu_{1} a I_{n-1} - \frac{\xi'^{T}\xi'}{\mu_{2}} \sinh \mu_{2} a & -i\xi' b_{1}(a) \\ -i^{T}\xi' b_{1}(a) & b_{3}(a) \end{pmatrix}$$

•

Let T' be the $(n-1) \times (n-1)$ orthogonal matrix given in the proof of Proposition 3.3 such as $T'\xi' = |\xi'|e'_{n-1}$ and set

$$T = \left(\begin{array}{cc} T' & 0\\ \\ 0 & 1 \end{array}\right).$$

Then

$$T(A_2(a) - A_1(a)A_2(0))T^{-1}$$

$$= \begin{pmatrix} \mu_1 \sinh \mu_1 a I_{n-2} & 0 & 0\\ 0 & b_2(a) & -i |\xi'| b_1(a) \\ 0 & -i |\xi'| b_1(a) & b_3(a) \end{pmatrix}.$$

It follows that

(3.6)
$$\det A = \det \left(T(A_2(a) - A_1(a)A_2(0))T^{-1} \right) \\ = \{ \mu_1 \sinh \mu_1 a \}^{n-2} D(\lambda, \xi').$$

Therefore, we see that $\lambda \in \rho(-\hat{L}'_{\xi})$ if and only if $D(\lambda, \xi') \neq 0$. This completes the proof.

To obtain a solution formula for (2.1) we next consider the Fourier series expansion of the solution $u = \begin{pmatrix} \phi \\ m \end{pmatrix}$ of (2.1). We expand ϕ and m into the Fourier series as

$$\phi = \frac{1}{2}\phi_0 + \sum_{k=1}^{\infty} \phi_k \cos a_k x_n, \quad m' = \frac{1}{2}m'_0 + \sum_{k=1}^{\infty} m'_k \cos a_k x_n$$

and

$$m^n = \frac{1}{2}m_0^n + \sum_{k=1}^{\infty} m_k^n \sin a_k x_n,$$

where $a_k = k\pi/a$; and $m_0^n = 0$,

Similarly, we expand

$$\phi_k = \frac{2}{a} \int_0^a \phi(x_n) \cos a_k x_n \, dx_n, \quad m'_k = \frac{2}{a} \int_0^a m'(x_n) \cos a_k x_n \, dx_n$$

and

$$m_k^n = \frac{2}{a} \int_0^a m^n(x_n) \sin a_k x_n \, dx_n.$$
$$f = \begin{pmatrix} f^0 \\ f' \\ f^n \end{pmatrix} \text{ as }$$

$$f^{0} = \frac{1}{2}f_{0}^{0} + \sum_{k=1}^{\infty} f_{k}^{0} \cos a_{k}x_{n}, \quad f' = \frac{1}{2}f'_{0} + \sum_{k=1}^{\infty} f'_{k} \cos a_{k}x_{n}$$

and

$$f^{n} = \frac{1}{2}f_{0}^{n} + \sum_{k=1}^{\infty} f_{k}^{n} \sin a_{k}x_{n},$$

where the Fourier coefficients are defined as above.

Similarly to $\lambda_{1,k}$, we also introduce $\lambda_{\pm,k}$ by

$$\lambda_{\pm,k} = -\frac{\nu_1}{2} |\xi^{(k)}|^2 \pm \frac{1}{2} \sqrt{\nu_1^2 |\xi^{(k)}|^4 - 4\gamma^2 |\xi^{(k)}|^2},$$

for $\xi' \in \mathbf{R}^{n-1}$ and $k = 1, 2, \cdots$. Here, as above, $|\xi^{(k)}|^2 = |\xi'|^2 + a_k^2$.

Remark 3.5. We note that $\lambda_{\pm,k}$ have properties similar to those of $\lambda_{\pm,0}$, namely, $\lambda_{\pm,k}$ are the two roots of $\lambda^2 + \nu_1 |\xi^{(k)}|^2 \lambda + \gamma^2 |\xi^{(k)}|^2 = 0$; $\lambda_{-,k} = \overline{\lambda_{+,k}}$

with $\operatorname{Im} \lambda_{+,k} = \gamma |\xi^{(k)}| \sqrt{1 - \frac{\nu_1^2}{4\gamma^2} |\xi^{(k)}|^2}$ when $|\xi^{(k)}| < 2\gamma/\nu_1$ and $\lambda_{\pm,k} \in \mathbf{R}$ when $|\xi^{(k)}| > 2\gamma/\nu_1$; and

$$\lambda_{+,k} = -\frac{\gamma^2}{\nu_1} + O(|\xi^{(k)}|^{-2}), \quad \lambda_{-,k} = -\nu_1 |\xi^{(k)}|^2 + O(1)$$

as $|\xi^{(k)}| \to \infty$. In contrast to the case k = 0, we find that there exists a positive number η_0 such that

$$\lambda_{1,k}, \ \lambda_{\pm,k} \notin \{\lambda; \operatorname{Re} \lambda \ge -\eta_0\}$$

for all $k \geq 1$ and $\xi' \in \mathbf{R}^{n-1}$.

The Fourier expansions described above are based on the reflection symmetry of equation (1.1), but they do not fit in the boundary condition (1.2). This leads to a decomposition of the Fourier coefficients ϕ_k , m'_k and m^n_k into two parts, one of which involves the boundary values of the x_n derivative of m' at $x_n = 0, a$.

Proposition 3.6. Let $\xi' \neq 0$. Assume that $\lambda \neq 0$, $\tilde{\nu}\lambda + \gamma^2 \neq 0$, $\nu_1\lambda + \gamma^2 \neq 0$, $\lambda \neq \lambda_{1,k}$ and $\lambda \neq \lambda_{\pm,k}$ for any $k = 0, 1, 2, \cdots$. Then the Fourier coefficients $u_k = \begin{pmatrix} \phi_k \\ m_k \end{pmatrix}$, $k = 0, 1, 2, \cdots$, are given by

$$u_k = \widehat{L}_k(\lambda, \xi')^{-1} f_k + \widehat{L}_k(\lambda, \xi')^{-1} Y_k.$$

Here

$$\widehat{L}_{k}(\lambda,\xi')^{-1} = \frac{1}{\lambda - \lambda_{1,k}} \begin{pmatrix} 0 & 0 \\ 0 & P_{0,k} \end{pmatrix}$$

$$+ \frac{1}{(\lambda - \lambda_{+,k})(\lambda - \lambda_{-,k})} \begin{pmatrix} \lambda + \nu_{1} |\xi^{(k)}|^{2} & -i\gamma^{T}\xi' & -\gamma a_{k} \\ -i\gamma\xi' & \lambda \frac{\xi'^{T}\xi'}{|\xi^{(k)}|^{2}} & -i\lambda \frac{a_{k}\xi'}{|\xi^{(k)}|^{2}} \\ \gamma a_{k} & i\lambda \frac{a_{k}^{T}\xi'}{|\xi^{(k)}|^{2}} & \lambda \frac{a_{k}^{2}}{|\xi^{(k)}|^{2}} \end{pmatrix}$$

where $P_{0,k}$ is an $n \times n$ matrix defined by

$$P_{0,k} = I_n - \begin{pmatrix} \frac{\xi'^T \xi'}{|\xi^{(k)}|^2} & -i\frac{a_k \xi'}{|\xi^{(k)}|^2} \\ i\frac{a_k^T \xi'}{|\xi^{(k)}|^2} & \frac{a_k^2}{|\xi^{(k)}|^2} \end{pmatrix}$$

and Y_k is given by

$$Y_k = \begin{pmatrix} 0 \\ \frac{2\nu}{a} \{ (-1)^k \partial_{x_n} m'(a) - \partial_{x_n} m'(0) \} \\ 0 \end{pmatrix}.$$

Proof. Since

$$\frac{2}{a} \int_0^a \partial_{x_n} w(x_n) \cos a_k x_n \, dx_n = \frac{2}{a} \{ (-1)^k w(a) - w(0) \} \\ + a_k \frac{2}{a} \int_0^a w(x_n) \sin a_k x_n \, dx_n$$

and

$$\frac{2}{a}\int_0^a \partial_{x_n} w(x_n) \sin a_k x_n \, dx_n = -a_k \frac{2}{a}\int_0^a w(x_n) \cos a_k x_n \, dx_n,$$

it is not so difficult to obtain the desired expression of u_k from (2.1). We omit the details.

We next compute the term Y_k which involves the boundary values of the x_n derivative of m'. To do so, we make use of Green's formula for (3.2). We define $(n-1) \times (n-1)$ matrices

$$P'_{1,0} = \frac{\xi'^T \xi'}{|\xi'|^2}, \quad P'_{0,0} = I_{n-1} - P'_{1,0}.$$

Proposition 3.7. Let the assumption of Proposition 3.6 be satisfied. Then Y_k has the form

$$Y_k = \frac{2}{a} \int_0^a \{ (-1)^k B(\lambda, \xi', y_n) + \check{B}(\lambda, \xi', a - y_n) \} f(y_n) \, dy_n,$$

where $B(\lambda,\xi',y_n)$ and $\check{B}(\lambda,\xi',y_n)$ are $(n+1)\times(n+1)$ matrices of the form

$$B(\lambda,\xi',y_n) = \begin{pmatrix} 0 & 0 & 0 \\ \mathbf{b}_0(\lambda,\xi',y_n) & B'(\lambda,\xi',y_n) & \mathbf{b}_n(\lambda,\xi',y_n) \\ 0 & 0 & 0 \end{pmatrix}$$

and

$$\check{B}(\lambda,\xi',y_n) = B(\lambda,\xi',y_n) \operatorname{diag}(1,\cdots,1,-1).$$

Here $\mathbf{b}_0(\lambda, \xi', y_n)$ and $\mathbf{b}_n(\lambda, \xi', y_n)$ are (n-1)-vectors of the form

 $\boldsymbol{b}_0(\lambda,\xi',y_n) = i\xi'\beta_0(\lambda,\xi',y_n)$

with

$$\beta_0(\lambda,\xi',y_n) = \frac{\gamma\lambda}{\nu_1\lambda + \gamma^2} \frac{1}{D(\lambda,\xi')} \{b_3(a)\frac{1}{\mu_2}\sinh\mu_2 y_n + b_1(a)\cosh\mu_2 y_n\}$$

and

$$\boldsymbol{b}_{n}(\lambda,\xi',y_{n}) = -\frac{i\xi'}{D(\lambda,\xi')} \{b_{3}(a)b_{1}(y_{n}) - b_{1}(a)b_{3}(y_{n})\};$$

and $B'(\lambda,\xi',y_n)$ is an $(n-1)\times(n-1)$ matrix defined by

$$B'(\lambda,\xi',y_n) = -\frac{\sinh \mu_1 y_n}{\sinh \mu_1 a} P'_{0,0} - \beta_1(\lambda,\xi',y_n) P'_{1,0}$$

with

$$\beta_1(\lambda,\xi',y_n) = \frac{1}{D(\lambda,\xi')} \{ b_3(a)b_2(y_n) + |\xi'|^2 b_1(a)b_1(y_n) \}.$$

Proof. In the proof we denote $Y_k = \begin{pmatrix} 0 \\ \tilde{Y}_k \end{pmatrix}$. Since $\lambda \neq 0$, (2.1) is equivalent to (3.1) and (3.2). Consider the following equation

(3.7)
$$\widetilde{M}(\lambda,\xi')m = 0,$$

where

$$\widetilde{M}(\lambda,\xi') = \begin{pmatrix} \{\lambda^2 + \nu\lambda(|\xi'|^2 - \partial_{x_n}^2)\}I_{n-1} + (\tilde{\nu}\lambda + \gamma^2)\xi'^T\xi' & i(\tilde{\nu}\lambda + \gamma^2)\xi'\partial_{x_n} \\ \\ i(\tilde{\nu}\lambda + \gamma^2)^T\xi'\partial_{x_n} & \lambda^2 + \nu\lambda|\xi'|^2 - \{\nu_1\lambda + \gamma^2\}\partial_{x_n}^2 \end{pmatrix}.$$

As in the proof of Proposition 3.3, one can see that the following functions $\tilde{v}_1, \dots, \tilde{v}_{2n}$ form a fundamental set of solutions of (3.7):

$$\widetilde{v}_j(x_n) = \begin{pmatrix} \mathbf{e}'_j \cosh \mu_1 x_n \\ i\frac{\xi_j}{\mu_1} \sinh \mu_1 x_n \end{pmatrix}, \quad \widetilde{v}_n(x_n) = \begin{pmatrix} -i\frac{\xi'}{\mu_2} \sinh \mu_2 x_n \\ \cosh \mu_2 x_n \end{pmatrix},$$
$$\widetilde{v}_{n+j}(x_n) = \begin{pmatrix} \mathbf{e}'_j \mu_1 \sinh \mu_1 x_n \\ i\xi_j \cosh \mu_1 x_n \end{pmatrix}, \quad \widetilde{v}_{2n}(x_n) = \begin{pmatrix} -i\xi' \cosh \mu_2 x_n \\ \mu_2 \sinh \mu_2 x_n \end{pmatrix},$$

where $j = 1, \dots, n-1$. We note that $\tilde{v}_1, \dots, \tilde{v}_{2n}$ are analytic in λ and $|\xi'|^2$. By Green's formula for (3.2), we see that

$$((F, v_j)) = ((m, \widetilde{M}(\lambda, \xi')\widetilde{v}_j)) - \lambda \nu \left[\partial_{x_n} m' \cdot \widetilde{v}_j'\right]_{x_n=0}^{x_n=a} - \{\nu_1 \lambda + \gamma^2\} \left[\partial_{x_n} m^n \widetilde{v}_j^n\right]_{x_n=0}^{x_n=a} = -\partial_{x_n} V(a) \cdot \widetilde{v}_j(a) + \partial_{x_n} V(0) \cdot \widetilde{v}_j(0),$$

where

$$V(x_n) = \left(\begin{array}{c} \nu\lambda m'(x_n)\\ \{\nu_1\lambda + \gamma^2\}m^n(x_n) \end{array}\right).$$

We thus obtain

$$\tilde{A} \begin{pmatrix} \partial_{x_n} V(0) \\ \partial_{x_n} V(a) \end{pmatrix} = \begin{pmatrix} \tilde{F}^{(1)} \\ \tilde{F}^{(2)} \end{pmatrix}$$

with

$$\widetilde{A} = \begin{pmatrix} A^{(1)}(0) & -A^{(1)}(a) \\ A^{(2)}(0) & -A^{(2)}(a) \end{pmatrix}, \quad F^{(j)} = ((A^{(j)}, F)) \quad (j = 1, 2),$$
$$A^{(1)}(x_n) = \begin{pmatrix} ^T \widetilde{v}_1(x_n) \\ \vdots \\ ^T \widetilde{v}_n(x_n) \end{pmatrix}, \quad A^{(2)}(x_n) = \begin{pmatrix} ^T \widetilde{v}_{n+1}(x_n) \\ \vdots \\ ^T \widetilde{v}_{2n}(x_n) \end{pmatrix}.$$

For a moment we assume that \widetilde{A} is invertible. We write the inverse of \widetilde{A} as

$$\widetilde{A}^{-1} = \left(\begin{array}{cc} S_1 & S_2 \\ \\ S_3 & S_4 \end{array}\right).$$

Then we have

$$\begin{pmatrix} \partial_{x_n} V(0) \\ \partial_{x_n} V(a) \end{pmatrix} = \int_0^a \begin{pmatrix} S_1 A^{(1)}(y_n) + S_2 A^{(2)}(y_n) \\ S_3 A^{(1)}(y_n) + S_4 A^{(2)}(y_n) \end{pmatrix} F(y_n) \, dy_n,$$

and, hence,

$$\begin{split} \widetilde{Y}_k &= \frac{2}{a\lambda} \widetilde{Q}'\{(-1)^k \partial_{x_n} V(a) - \partial_{x_n} V(0)\} \\ &= \frac{2}{a\lambda} \int_0^a \widetilde{Q}'\{(-1)^k (S_3 A^{(1)}(y_n) + S_4 A^{(2)}(y_n)) \\ &- (S_1 A^{(1)}(y_n) + S_2 A^{(2)}(y_n))\} F(y_n) \, dy_n. \end{split}$$

Here and in what follows we denote the $n \times n$ matrix diag $(1, \dots, 1, 0)$ by \tilde{Q}' :

$$\widetilde{Q}' = \operatorname{diag}(1, \cdots, 1, 0).$$

By a direct calculation we have

$$A^{(1)}(x_n) = \begin{pmatrix} \cosh \mu_1 x_n I_{n-1} & i\frac{\xi'}{\mu_1} \sinh \mu_1 x_n \\ -i\frac{\tau_{\xi'}}{\mu_2} \sinh \mu_2 x_n & \cosh \mu_2 x_n \end{pmatrix}$$

and

$$A^{(2)}(x_n) = \begin{pmatrix} \mu_1 \sinh \mu_1 x_n I_{n-1} & i\xi' \cosh \mu_1 x_n \\ -i^T \xi' \cosh \mu_2 x_n & \mu_2 \sinh \mu_2 x_n \end{pmatrix}.$$

We thus obtain

$$\widetilde{A} = \begin{pmatrix} I_n & -A^{(1)}(a) \\ A^{(2)}(0) & -A^{(2)}(a) \end{pmatrix}$$

We define $A(y_n)$ by

$$A(y_n) = A^{(2)}(0)A^{(1)}(y_n) - A^{(2)}(y_n)$$

=
$$\begin{pmatrix} -\mu_1 \sinh \mu_1 y_n I_{n-1} + \frac{|\xi'|^2}{\mu_2} \sinh \mu_2 y_n P'_{1,0} & -ib_1(y_n)\xi' \\ & -ib_1(y_n)^T\xi' & -b_3(y_n) \end{pmatrix}$$

One can see that if A(a) is invertible, then so is \tilde{A} , and \tilde{A}^{-1} is given by

$$\tilde{A}^{-1} = \begin{pmatrix} I_n - A^{(1)}(a)A(a)^{-1}A^{(2)}(0) & A^{(1)}(a)A(a)^{-1} \\ -A(a)^{-1}A^{(2)}(0) & A(a)^{-1} \end{pmatrix}.$$

We now verify that A(a) is invertible under the assumption of Proposition 3.7. We denote the $n \times n$ diagonal matrix diag $(1, \dots, 1, -1)$ by \check{I}_n . We observe that $A^{(j)}(x_n) = {}^T \left[\check{I}_n A_j(x_n)\check{I}_n\right], j = 1, 2$, where $A_j(x_n)$ are the matrices given in the proof of Lemma 3.4. Therefore, we have

$$\widetilde{A} = {}^{T} \left[\operatorname{diag} \left(\check{I}_{n}, -\check{I}_{n} \right) A \operatorname{diag} \left(\check{I}_{n}, \check{I}_{n} \right) \right].$$

It then follows from (3.6) that

det
$$\tilde{A} = (-1)^n \det A = (-1)^n \{\mu_1 \sinh \mu_1 a\}^{n-2} D(\lambda, \xi'),$$

and hence, by Lemma 3.4, A(a) is invertible under the assumption of Proposition 3.7.

We thus obtain

(3.8)
$$\widetilde{Y}_k = \frac{2}{a\lambda} \int_0^a \{(-1)^k \widetilde{B}^{(1)}(\lambda, \xi', y_n) + \widetilde{B}^{(2)}(\lambda, \xi', y_n)\} F(y_n) \, dy_n,$$

where $\tilde{B}^{(1)}(\lambda,\xi,y_n) = -\tilde{Q}'A(a)^{-1}A(y_n)$ and $\tilde{B}^{(2)}(\lambda,\xi',y_n) = \tilde{Q}'(A^{(1)}(y_n) - A^{(1)}(a)A(a)^{-1}A(y_n)).$

We next show that

(3.9)
$$-A(a)^{-1}A(y_n) = K^{(1)}(\lambda, \xi', y_n) + K^{(2)}(\lambda, \xi', y_n),$$

where

$$K^{(1)}(\lambda,\xi',y_n) = -\frac{\sinh\mu_1 y_n}{\sinh\mu_1 a} \begin{pmatrix} P'_{0,0} & 0\\ 0 & 0 \end{pmatrix}$$

and

$$K^{(2)}(\lambda,\xi',y_n) = -\frac{1}{D(\lambda,\xi')} \begin{pmatrix} \{b_3(a)b_2(y_n) + |\xi'|^2b_1(a)b_1(y_n)P'_{1,0} & i\{b_3(a)b_1(y_n) - b_1(a)b_3(y_n)\}\xi'\\ \\ i\{b_2(a)b_1(y_n) - b_1(a)b_2(y_n)\}^T\xi' & b_2(a)b_3(y_n) + |\xi'|^2b_1(a)b_1(y_n) \end{pmatrix}.$$

Let us prove (3.9). Noting that $P'_{1,0}$ is an orthogonal projection onto the subspace spanned by ξ' , namely, $P_{1,0}\xi' = \xi'$, ${}^{T}\xi'P'_{1,0} = {}^{T}\xi'$ and ${P'_{1,0}}^2 = P'_{1,0}$, we see that

$$A(a)^{-1} = -\frac{1}{\mu_1 \sinh \mu_1 a} \begin{pmatrix} P'_{0,0} & 0\\ 0 & 0 \end{pmatrix} + \frac{1}{D(\lambda,\xi')} \begin{pmatrix} -b_3(a)P'_{1,0} & ib_1(a)\xi'\\ ib_1(a)^T\xi' & -b_2(a) \end{pmatrix},$$

since $\mu_1 \sinh \mu_1 a \neq 0$ and $D \neq 0$. We now obtain (3.9) by a direct computation.

We next prove $\tilde{B}^{(2)}(\lambda,\xi',a-y_n) = \tilde{B}^{(1)}(\lambda,\xi',y_n)\check{I}_n$. To do so, we write $\tilde{B}^{(j)}(\lambda,\xi',y_n)$ as

(3.10)
$$\widetilde{B}^{(j)}(\lambda,\xi',y_n) = \begin{pmatrix} B^{\prime(j)}(\lambda,\xi',y_n) & \boldsymbol{b}_n^{(j)}(\lambda,\xi',y_n) \\ 0 & 0 \end{pmatrix}$$

for j = 1, 2. Then we need to show

(3.11)
$$B'^{(1)}(\lambda,\xi',y_n) = B'^{(2)}(\lambda,\xi',a-y_n)$$

and

(3.12)
$$\boldsymbol{b}_n^{(1)}(\lambda,\xi',y_n) = -\boldsymbol{b}_n^{(2)}(\lambda,\xi',a-y_n).$$

To prove (3.11) and (3.12) we make use of the following reflection symmetry of problem (3.2). For $m(x_n) = \begin{pmatrix} m'(x_n) \\ m^n(x_n) \end{pmatrix}$ we define $\check{m}(x_n)$ by $\check{m}(x_n) = \begin{pmatrix} m'(a - x_n) \\ -m^n(a - x_n) \end{pmatrix}$.

Let $m(x_n)$ be a solution of problem (3.2) with $F(x_n) = \lambda \begin{pmatrix} f'(x_n) \\ f^n(x_n) \end{pmatrix} - \begin{pmatrix} i\xi' \\ - k \end{pmatrix}$

$$\gamma \begin{pmatrix} i\zeta \\ \partial_{x_n} \end{pmatrix} f^0(x_n)$$
. Then $\check{m}(x_n)$ is a solution of problem (3.2) with $F(x_n)$

replaced by
$$\check{F}(x_n) = \begin{pmatrix} F'(a-x_n) \\ -F^n(a-x_n) \end{pmatrix}$$
. Since $\partial_{x_n}(\check{m})'(a) = -\partial_{x_n}m'(0)$ and $\partial_{x_n}(\check{m})'(0) = -\partial_{x_n}m'(a)$, we have

$$(3.13) \ \ (-1)^k \partial_{x_n}(\check{m})'(a) - \partial_{x_n}(\check{m})'(0) = (-1)^k \{ (-1)^k \partial_{x_n} m'(a) - \partial_{x_n} m'(0) \}.$$

We next define $B'_k(\lambda,\xi',y_n)$ and $\boldsymbol{b}_{n,k}(\lambda,\xi',y_n)$ by

(3.14)
$$B'_k(\lambda,\xi',y_n) = (-1)^k B'^{(1)}(\lambda,\xi',y_n) + B'^{(2)}(\lambda,\xi',y_n)$$

and

(3.15)
$$\boldsymbol{b}_{n,k}(\lambda,\xi',y_n) = (-1)^k \boldsymbol{b}_n^{(1)}(\lambda,\xi',y_n) + \boldsymbol{b}_n^{(2)}(\lambda,\xi',y_n).$$

Since $\mathbf{b}_{n,k}(\lambda, \xi', 0) = \mathbf{b}_{n,k}(\lambda, \xi', a) = 0$, integrating by parts, we see from (3.8) that

$$(3.16)$$

$$\frac{2}{a}\lambda\nu\{(-1)^{k}\partial_{x_{n}}(\check{m})'(a) - \partial_{x_{n}}(\check{m})'(0)\}$$

$$= \frac{2}{a}\int_{0}^{a} [B'_{k}(\lambda,\xi',y_{n})\{\lambda f'(a-y_{n}) - i\gamma\xi'f^{0}(a-y_{n})\}\}$$

$$-\lambda \boldsymbol{b}_{n,k}(\lambda,\xi',y_{n})f^{n}(a-y_{n})$$

$$+\gamma\partial_{y_{n}}\boldsymbol{b}_{n,k}(\lambda,\xi',y_{n})f^{0}(a-y_{n})]\,dy_{n}$$

$$= \frac{2}{a}\int_{0}^{a} [B'_{k}(\lambda,\xi',a-y_{n})\{\lambda f'(y_{n}) - i\gamma\xi'f^{0}(y_{n})\}\}$$

$$-\lambda \boldsymbol{b}_{n,k}(\lambda,\xi',a-y_{n})f^{n}(y_{n})$$

$$+\gamma\partial_{y_{n}}\boldsymbol{b}_{n,k}(\lambda,\xi',a-y_{n})f^{0}(y_{n})]\,dy_{n}$$

and

$$(3.17) \qquad \frac{2}{a}\lambda\nu\{(-1)^{k}\partial_{x_{n}}m'(a) - \partial_{x_{n}}m'(0)\}$$
$$= \frac{2}{a}\int_{0}^{a} [B'_{k}(\lambda,\xi',y_{n})\{\lambda f'(y_{n}) - i\gamma\xi'f^{0}(y_{n})\} + \lambda \boldsymbol{b}_{n,k}(\lambda,\xi',y_{n})f^{n}(y_{n}) + \gamma\partial_{y_{n}}\boldsymbol{b}_{n,k}(\lambda,\xi',y_{n})f^{0}(y_{n})] dy_{n}.$$

Combining (3.13), (3.16) and (3.17) we obtain $B'_k(\lambda, \xi', a-y_n) = (-1)^k B'_k(\lambda, \xi', y_n)$ and $\boldsymbol{b}_{n,k}(\lambda, \xi', a-y_n) = (-1)^{k+1} \boldsymbol{b}_{n,k}(\lambda, \xi', y_n)$. This, together with (3.14) and (3.15), implies that

(3.18)
$$(-1)^{k} B'^{(1)}(\lambda,\xi',a-y_{n}) + B'^{(2)}(\lambda,\xi',a-y_{n})$$
$$= B'^{(1)}(\lambda,\xi',y_{n}) + (-1)^{k} B'^{(2)}(\lambda,\xi',y_{n})$$

and

(3.19)
$$(-1)^{k} \boldsymbol{b}_{n}^{(1)}(\lambda,\xi',a-y_{n}) + \boldsymbol{b}_{n}^{(2)}(\lambda,\xi',a-y_{n}) \\ = -\boldsymbol{b}_{n}^{(1)}(\lambda,\xi',y_{n}) - (-1)^{k} \boldsymbol{b}_{n}^{(2)}(\lambda,\xi',y_{n}).$$

It follows from (3.18), with k being odd and even respectively, that

$$\begin{cases} -B'^{(1)}(\lambda,\xi',a-y_n) + B'^{(2)}(\lambda,\xi',a-y_n) = B'^{(1)}(\lambda,\xi',y_n) - B'^{(2)}(\lambda,\xi',y_n), \\ B'^{(1)}(\lambda,\xi',a-y_n) + B'^{(2)}(\lambda,\xi',a-y_n) = B'^{(1)}(\lambda,\xi',y_n) + B'^{(2)}(\lambda,\xi',y_n), \end{cases}$$

which implies that $B'^{(2)}(\lambda,\xi',a-y_n) = B'^{(1)}(\lambda,\xi',y_n)$. This shows (3.11). Similarly, one can prove (3.12) by using (3.19).

We finally set $B'(\lambda, \xi', y_n) = B'^{(1)}(\lambda, \xi', y_n)$, $\boldsymbol{b}_n(\lambda, \xi', y_n) = \boldsymbol{b}_n^{(1)}(\lambda, \xi', y_n)$ and $\boldsymbol{b}_0(\lambda, \xi', y_n) = -\frac{\gamma}{\lambda} \{ iB'(\lambda, \xi', y_n)\xi' - \partial_{y_n}\boldsymbol{b}_n(\lambda, \xi', y_n) \}$. Then, by a direct calculation, we see that

$$\boldsymbol{b}_0 = \frac{i\gamma\lambda\xi'}{\nu_1\lambda + \gamma^2} \frac{1}{D(\lambda,\xi')} \left\{ b_3(a) \frac{1}{\mu_2} \sinh\mu_2 y_n + b_1(a) \cosh\mu_2 y_n \right\},\,$$

and the desired expression of Y_k is obtained. This completes the proof.

We now give an integral representation of $(\lambda + \hat{L}_{\xi'})^{-1}f$. To do so, we introduce some functions. We define $g^{D}_{\mu_{j}}(x_{n}, y_{n})$ (j = 1, 2) by

$$g_{\mu_j}^D(x_n, y_n) = \frac{1}{\mu_j \sinh \mu_j a} \sinh \mu_j (a - x_n) \sinh \mu_j y_n, \quad y_n \le x_n,$$

and x_n, y_n exchanged for $x_n \leq y_n$. Similarly, we define $g_{\mu_j}^N(x_n, y_n)$ by

$$g_{\mu_j}^N(x_n, y_n) = \frac{1}{\mu_j \sinh \mu_j a} \cosh \mu_j (a - x_n) \cosh \mu_j y_n, \quad y_n \le x_n,$$

and x_n, y_n exchanged for $x_n \leq y_n$. We set

$$g_{\mu_1,\mu_2}^M(x_n,y_n) = g_{\mu_1}^M(x_n,y_n) - g_{\mu_2}^M(x_n,y_n), \quad M = D, N.$$

Note that $g_{\mu_j}^D$ and $g_{\mu_j}^N$ are the Green functions of the equation $\mu_j^2 v - \partial_{x_n}^2 v = 0$ under the Dirichlet and Neumann boundary conditions at $\{x_n = 0, a\}$, respectively. We also define $h_{\mu_j}(x_n)$ and $h_{\mu_1,\mu_2}(x_n)$ by

$$h_{\mu_j}(x_n) = \frac{1}{\mu_j \sinh \mu_j a} \cosh \mu_j x_n$$

and

$$h_{\mu_1,\mu_2}(x_n) = h_{\mu_1}(x_n) - h_{\mu_2}(x_n)$$

We will denote the Dirac measure with point mass at x_n by δ_{x_n} , namely,

$$\delta_{x_n} f = \int_0^a \delta(x_n - y_n) f(y_n) \, dy_n = f(x_n).$$

Theorem 3.8. Let λ satisfy $\lambda \neq 0$, $\tilde{\nu}\lambda + \gamma^2 \neq 0$, $\nu_1\lambda + \gamma^2 \neq 0$. Assume that $\lambda \neq \lambda_{1,k}$ and $\lambda \neq \lambda_{\pm,k}$ for any $k = 0, 1, 2, \cdots$. Assume also that $\lambda \in \rho(-\hat{L}'_{\xi})$. Then the solution $(\lambda + \hat{L}_{\xi'})^{-1}f$ of (2.1) is written as

$$(\lambda + \widehat{L}_{\xi'})^{-1}f = \widehat{G}(\lambda, \xi')f + \widehat{K}(\lambda, \xi')f$$

with integral operators $\widehat{G}(\lambda,\xi')$ and $\widehat{K}(\lambda,\xi')$ defined by

$$(\widehat{G}(\lambda,\xi')f)(x_n) = \int_0^a \widehat{G}(\lambda,\xi',x_n,y_n)f(y_n)\,dy_n$$

and

$$(\widehat{K}(\lambda,\xi')f)(x_n) = \int_0^a \widehat{K}(\lambda,\xi',x_n,y_n)f(y_n)\,dy_n.$$

Here $\widehat{G}(\lambda,\xi',x_n,y_n)$ is an $(n+1) \times (n+1)$ matrix of the form

$$\begin{split} \hat{G}(\lambda,\xi',x_n,y_n) &= \frac{\nu_1}{d(\lambda)} \delta(x_n - y_n) Q_0 \\ &+ \frac{\gamma}{d(\lambda)} \begin{pmatrix} \frac{\gamma \lambda}{d(\lambda)} g_{\mu_2}^N(x_n,y_n) & -i^T \xi' g_{\mu_2}^N(x_n,y_n) & -\partial_{x_n} g_{\mu_2}^D(x_n,y_n) \\ -i\xi' g_{\mu_2}^N(x_n,y_n) & 0 & 0 \\ -\partial_{x_n} g_{\mu_2}^N(x_n,y_n) & 0 & 0 \\ &+ \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{\nu} g_{\mu_1}^N(x_n,y_n) I_{n-1} & 0 \\ 0 & 0 & \frac{1}{\nu} g_{\mu_1}^D(x_n,y_n) \end{pmatrix} \\ &+ \begin{pmatrix} 0 & 0 & 0 \\ 0 & \frac{\xi'^T \xi'}{\lambda} g_{\mu_1,\mu_2}^N(x_n,y_n) & -\frac{i\xi'}{\lambda} \partial_{x_n} g_{\mu_1,\mu_2}^D(x_n,y_n) \\ 0 & -\frac{i\xi'}{\lambda} \partial_{x_n} g_{\mu_1,\mu_2}^N(x_n,y_n) & -\frac{1}{\lambda} \partial_{x_n}^2 g_{\mu_1,\mu_2}^D(x_n,y_n) \end{pmatrix}, \end{split}$$
where $d(\lambda) = \nu_1 \lambda + \gamma^2$ and $\mu_j = \mu_j(\lambda,\xi'), \ j = 1,2;$ and

$$\widehat{K}(\lambda,\xi',x_n,y_n) = \widehat{H}(\lambda,\xi',x_n,y_n) + \check{H}(\lambda,\xi',a-x_n,a-y_n),$$

where

$$\check{H}(\lambda,\xi',a-x_n,a-y_n) = \widehat{H}(\lambda,\xi',a-x_n,a-y_n) \operatorname{diag}\left(I_n,-1\right)$$

$$\begin{aligned} and \\ H(\lambda,\xi',x_n,y_n) \\ &= \begin{pmatrix} 0 & 0 & 0 \\ \frac{i\xi'}{\nu}h_{\mu_1}(x_n)\beta_0(y_n) & \frac{1}{\nu}h_{\mu_1}(x_n)B'(y_n) & \frac{1}{\nu}h_{\mu_1}(x_n)\mathbf{b}_n(y_n) \\ 0 & 0 & 0 \end{pmatrix} \\ &+ \begin{pmatrix} \frac{\gamma|\xi'|^2}{d(\lambda)}h_{\mu_2}(x_n)\beta_0(y_n) & \frac{i\gamma^T\xi'}{d(\lambda)}h_{\mu_2}(x_n)\beta_1(y_n) & -\frac{i\gamma^T\xi'}{d(\lambda)}h_{\mu_2}(x_n)\mathbf{b}_n(y_n) \\ \frac{i|\xi'|^2\xi'}{\lambda}h_{\mu_1,\mu_2}(x_n)\beta_0(y_n) & 0 & 0 \\ -\frac{|\xi'|^2}{\lambda}\partial_{x_n}h_{\mu_1,\mu_2}(x_n)\beta_0(y_n) & 0 & 0 \end{pmatrix} \\ &+ \begin{pmatrix} 0 & 0 & 0 \\ 0 & -\frac{\xi'^T\xi'}{\lambda}h_{\mu_1,\mu_2}(x_n)\beta_1(y_n) & \frac{|\xi'|^2}{\lambda}h_{\mu_1,\mu_2}(x_n)\mathbf{b}_n(y_n) \\ 0 & -\frac{i'^T\xi'}{\lambda}\partial_{x_n}h_{\mu_1,\mu_2}(x_n)\beta_1(y_n) & \frac{i^T\xi'}{\lambda}\partial_{x_n}h_{\mu_1,\mu_2}(x_n)\mathbf{b}_n(y_n) \end{pmatrix} \\ with d(\lambda) = w_{\lambda} + \gamma^2 \quad w = w_{\lambda}(\lambda,\xi') \quad i = 1, 2, \beta_1(w_{\lambda}) = \beta_1(\lambda,\xi',w_{\lambda}) \quad i = 0, 1 \end{aligned}$$

with $d(\lambda) = \nu_1 \lambda + \gamma^2$, $\mu_j = \mu_j(\lambda, \xi')$, j = 1, 2, $\beta_j(y_n) = \beta_j(\lambda, \xi', y_n)$, j = 0, 1, $B'(y_n) = B'(\lambda, \xi', y_n)$ and $\mathbf{b}_n(y_n) = \mathbf{b}_n(\lambda, \xi', y_n)$.

Proof. By assumption we see that the Fourier series of $(\lambda + \hat{L}_{\xi'})^{-1}f$ takes the form as in Proposition 3.6 with Y_k given in Proposition 3.7.

The Fourier sine and cosine series of $w^M(x_n) = \int_0^a g^M_\mu(x_n, y_n) v(y_n) dy_n$ with M = D, N are given by

$$w^{D}(x_{n}) = \sum_{k=1}^{\infty} \frac{1}{\mu^{2} + a_{k}^{2}} v_{s,k} \sin a_{k} x_{n}$$

and

$$w^{N}(x_{n}) = \frac{1}{2}v_{c,0} + \sum_{k=1}^{\infty} \frac{1}{\mu^{2} + a_{k}^{2}} v_{c,k} \cos a_{k} x_{n},$$

respectively. Here $v_{s,k}$ and $v_{c,k}$ are the Fourier sine and cosine coefficients of v respectively. Since

$$\frac{1}{\lambda - \lambda_{1,k}} = \frac{1}{\nu} \frac{1}{\mu_1^2 + a_k^2},$$
$$\frac{1}{(\lambda - \lambda_{+,k})(\lambda - \lambda_{-,k})} = \frac{1}{\nu_1 \lambda + \gamma^2} \frac{1}{\mu_2^2 + a_k^2}$$

and

$$\frac{1}{\lambda - \lambda_{1,k}} \frac{1}{|\xi^{(k)}|^2} - \frac{\lambda}{(\lambda - \lambda_{+,k})(\lambda - \lambda_{-,k})} \frac{1}{|\xi^{(k)}|^2} = -\frac{1}{\lambda} \left(\frac{1}{\mu_1^2 + a_k^2} - \frac{1}{\mu_2^2 + a_k^2} \right),$$

we see that the Fourier coefficients of $\widehat{G}(\lambda,\xi')f$ are equal to $\widehat{L}_k(\lambda,\xi')^{-1}f_k$.

Also, since the Fourier cosine coefficients of $e^{\mu x_n}$ are given by

$$\frac{2}{a}\frac{\mu\{(-1)^k e^{\mu a} - 1\}}{\mu^2 + a_k^2}, \quad k = 0, 1, 2, \cdots,$$

we see from Propositions 3.6 and 3.7 that the Fourier coefficients of $\widehat{K}(\lambda,\xi')f$ are equal to $\widehat{L}_k(\lambda,\xi')^{-1}Y_k$. This completes the proof.

4. Preliminary estimates

In this section we prepare some estimates for the analysis of the integral kernel given in Theorem 3.8.

We first give an estimate of the resolvent set by the energy method.

Proposition 4.1. (i) There exists a positive constant c_1 such that the set

$$\Sigma_0 \equiv \{\lambda; \operatorname{Re} \lambda + c_1 |\operatorname{Im} \lambda|^2 > 0\}$$

is in the resolvent set $\rho(-\hat{L}_{\xi'})$ for all ξ' .

(ii) There exists a positive number η_1 such that the set

$$\{\lambda; \operatorname{Re} \lambda \ge -\eta_1\} \cap \{\lambda; \operatorname{Re} \lambda \le 0, \operatorname{Im} \lambda = 0\}^c$$

is in the resolvent set $\rho(-\hat{L}_{\xi'})$ for all ξ' . Here and in what follows, for a set E, the symbol E^c denotes the complementary set of E.

Proof. Let us consider problem (2.1). We note that $\widehat{A}_{\xi'}$ is self adjoint in $L^2(0, a)$ and $\widehat{B}_{\xi'}$ is skew-symmetric and the following relations hold:

(4.1)

$$\begin{aligned}
(\widehat{A}_{\xi'}u, u) &= \nu |\xi'|^2 |m|_2^2 + \nu |\partial_{x_n}m|_2^2 + \widetilde{\nu} |i\xi' \cdot m' + \partial_{x_n}m^n|_2^2, \\
(\widehat{B}_{\xi'}u, u) &= 2i\gamma \operatorname{Im} (i\xi' \cdot m' + \partial_{x_n}m^n, \phi). \\
\text{Let } u &= \begin{pmatrix} \phi \\ m \end{pmatrix} \text{ be a solution of (2.1). Taking the } L^2 \text{-inner product of} \\
(2.1) \text{ with } u \text{ we see from (4.1) that}
\end{aligned}$$

(4.2)
$$\lambda |u|_{2}^{2} + \nu |\xi'|^{2} |m|_{2}^{2} + \nu |\partial_{x_{n}}m|_{2}^{2} + \widetilde{\nu} |i\xi' \cdot m' + \partial_{x_{n}}m^{n}|_{2}^{2} + 2i\gamma \operatorname{Im}\left(i\xi' \cdot m' + \partial_{x_{n}}m^{n}, \phi\right) = (f, u).$$

The real part of (4.2) gives

(4.3)

$$\operatorname{Re} \lambda |u|_{2}^{2} + \nu |\xi'|^{2} |m|_{2}^{2} + \nu |\partial_{x_{n}}m|_{2}^{2} + \widetilde{\nu} |i\xi' \cdot m' + \partial_{x_{n}}m^{n}|_{2}^{2}$$

$$= \operatorname{Re} (f, u) \leq \varepsilon |u|_{2}^{2} + \frac{1}{\varepsilon} |f|_{2}^{2}$$

for any $\varepsilon > 0$. On the other hand, the imaginary part of (4.2) gives

$$\operatorname{Im} \lambda |u|_{2}^{2} = \operatorname{Im} (f, u) - 2\gamma \operatorname{Im} (i\xi' \cdot m' + \partial_{x_{n}} m^{n}, \phi),$$

from which we obtain

(4.4)
$$|\operatorname{Im} \lambda|^2 |u|_2^2 \le 2\left\{ |f|_2^2 + \gamma^2 |i\xi' \cdot m' + \partial_{x_n} m^n|_2^2 \right\}.$$

It then follows from (4.3) and (4.4) that

(4.5)
$$\left(\operatorname{Re}\lambda + c_1 |\operatorname{Im}\lambda|^2 - \varepsilon\right) |u|_2^2 + \frac{\nu}{2} \left(|\xi'|^2 |m|_2^2 + |\partial_{x_n}m|_2^2\right) \le C_{\varepsilon} |f|_2^2$$

for any $\varepsilon > 0$ with some constant $c_1 = c_1(\nu, \gamma) > 0$.

We next estimate $|\partial_{x_n}\phi|_2$. Differentiating the first row of (2.1) with respect to x_n we have

(4.6)
$$\lambda \partial_{x_n} \phi + \gamma \partial_{x_n}^2 m^n = \partial_{x_n} f^0 - i\gamma \xi' \cdot \partial_{x_n} m'$$

We also see from the third row of (2.1) that

(4.7)
$$-\nu_1 \partial_{x_n}^2 m^n + \gamma \partial_{x_n} \phi = g$$

where $g = f^n - \{\lambda m^n + \nu |\xi'|^2 m^n - i \tilde{\nu} \xi' \cdot \partial_{x_n} m'\}$. By adding (4.7) $\times \frac{\gamma}{\nu_1}$ to (4.6) we obtain

(4.8)
$$\left(\lambda + \frac{\gamma^2}{\nu_1}\right)\partial_{x_n}\phi = \partial_{x_n}f^0 - i\gamma\xi'\cdot\partial_{x_n}m' + \frac{\gamma}{\nu_1}g.$$

This implies that if $\lambda \neq -\frac{\gamma^2}{\nu_1}$, then

(4.9)
$$|\partial_{x_n}\phi|_2 \leq \frac{C}{\left|\lambda + \frac{\gamma^2}{\nu_1}\right|} \left\{ |f|_{H^1 \times L^2} + |\lambda| |m|_2 + |\xi'|^2 |m|_2 + |\xi'| |\partial_{x_n} m|_2 \right\}.$$

We thus deduce from (4.5) and (4.9) that

$$(4.10) |u|_{H^1 \times L^2} \le C |f|_{H^1 \times L^2}$$

for some C > 0, provided that $\lambda \in \Sigma_0$. This, together with (2.1), yields

(4.11)
$$|m|_{H^2} \le \tilde{C}|f|_{H^1 \times L^2}$$

with some constant $\tilde{C} = \tilde{C}(\eta_0, \theta_0, \xi') > 0$. Since $\lambda \neq 0$ and $\nu_1 \lambda + \gamma^2 \neq 0$ when $\lambda \in \Sigma_0$, it follows from (4.10) and (4.11) that Ker $\mathcal{L} = \{0\}$ for $\lambda \in \Sigma_0$. Lemma 3.1 then implies that $\Sigma_0 \subset \rho(-\hat{L}_{\xi'})$. This completes the proof of (i). We next prove (ii). In view of Lemma 3.1 it suffices to prove that (3.2) is uniquely solvable for any $F \in L^2(0, a)$ if $\lambda \in \{\lambda; \operatorname{Re} \lambda \geq -\eta_1\} \cap \{\lambda; \operatorname{Re} \lambda \leq 0, \operatorname{Im} \lambda = 0\}^c$ with some $\eta_1 > 0$.

We take the L^2 -inner product of (3.2) with m. Then, integrating by parts, we have

(4.12)
$$(\lambda^2 + \nu\lambda|\xi'|^2)|m|_2^2 + \nu\lambda|\partial_{x_n}m|_2^2 + (\tilde{\nu}\lambda + \gamma^2)|i\xi' \cdot m' + \partial_{x_n}m^n|_2^2 = (F, m).$$

The imaginary part of (4.12) gives

$$\operatorname{Im} \lambda \left\{ (2\operatorname{Re} \lambda + \nu |\xi'|^2) |m|_2^2 + \nu |\partial_{x_n} m|_2^2 + \widetilde{\nu} |i\xi' \cdot m' + \partial_{x_n} m^n|_2^2 \right\} = \operatorname{Im} (F, m).$$

It follows that if $\operatorname{Im} \lambda \neq 0$, then

(4.13)
$$(2\operatorname{Re}\lambda + \nu|\xi'|^2) |m|_2^2 + \nu|\partial_{x_n}m|_2^2 + \tilde{\nu}|i\xi' \cdot m' + \partial_{x_n}m^n|_2^2 = \frac{\operatorname{Im}(F,m)}{\operatorname{Im}\lambda}$$

By the Poincaré inequality, there exists a positive constant η_1 such that the left-hand side of (4.13) is bounded from below by

$$\left(2\operatorname{Re}\lambda + \nu|\xi'|^2 + 4\eta_1\right)|m|_2^2 + \frac{\nu}{2}|\partial_{x_n}m|_2^2,$$

while the right-hand side of (4.13) is bounded from above by

$$\eta_1 |m|_2^2 + \frac{C}{|\mathrm{Im}\,\lambda|^2} |F|_2^2.$$

We thus conclude that

$$\left(2\operatorname{Re}\lambda + \nu|\xi'|^2 + 3\eta_1\right)|m|_2^2 + \frac{\nu}{2}|\partial_{x_n}m|_2^2 \le \frac{C}{|\operatorname{Im}\lambda|^2}|F|_2^2,$$

in particular, if $\lambda \in \operatorname{Re} \lambda \geq -\eta_1 - \frac{\nu |\xi'|^2}{2}$, $\operatorname{Im} \lambda \neq 0$ and F = 0, then m = 0. This, together with (i), implies that $\{\lambda; \operatorname{Re} \lambda \geq -\eta_1 - \frac{\nu |\xi'|^2}{2}\} \cap \{\lambda; \operatorname{Re} \lambda \leq 0, \operatorname{Im} \lambda = 0\}^c \subset \rho(-\widehat{L}_{\xi'})$. This completes the proof.

We next investigate $D(\lambda, \xi')$. We first estimate $|D(\lambda, \xi')|$ from below for small λ .

Proposition 4.2. (i) Let r_1 and R_1 be any positive numbers with $r_1 < R_1$. Then there exists a positive number $\Lambda_1 = \Lambda_1(r_1, R_1)$ such that the inequality

$$|D(\lambda,\xi')| \ge \frac{a^4}{24\nu^2} r_1^2 |\lambda|^2$$

holds for $|\lambda| \leq \Lambda_1$ and $r_1 \leq |\xi'| \leq R_1$.

(ii) There are positive numbers Λ_2 and $R_2 = R_2(\Lambda_2)$ such that the inequality

$$|D(\lambda,\xi')| \ge \frac{1}{16\nu^2} \left(\frac{\sinh|\xi'|a}{|\xi'|}\right)^2 |\lambda|^2$$

holds for $|\lambda| \leq \Lambda_2$ and $|\xi'| \geq R_2$.

Proof. Assume that $|\nu_1 \lambda + \gamma^2| \ge \frac{\gamma^2}{2\nu_1} \equiv c_2$. Let Λ and r be positive numbers satisfying

(4.14)
$$\frac{\Lambda}{\nu r^2} < 1 \text{ and } \frac{\Lambda^2}{c_2 r^2} < 1.$$

Then for $|\lambda| \leq \Lambda$ and $|\xi'| \geq r$, we have

$$\left|\frac{\lambda}{\nu|\xi'|^2}\right| \le \frac{\Lambda}{\nu r^2} < 1 \quad \text{and} \quad \left|\frac{\lambda^2}{\nu_1 \lambda + \gamma^2} \frac{1}{|\xi'|^2}\right| \le \frac{\Lambda^2}{c_2 r^2} < 1,$$

and hence,

$$\mu_1 = |\xi'| + \frac{\lambda}{2\nu} \frac{1}{|\xi'|} - \frac{\lambda^2}{8\nu^2} \frac{1}{|\xi'|^3} + O\left(|\lambda|^3 |\xi'|^{-5}\right)$$

and

$$\mu_2 = |\xi'| + \mu_2^{(1)} \frac{1}{|\xi'|} + \mu_2^{(2)} \frac{1}{|\xi'|^3} + O\left(|\lambda|^6 |\xi'|^{-5}\right).$$

Here

$$\mu_2^{(1)} = \frac{\lambda^2}{2(\nu_1 \lambda + \gamma^2)}$$
 and $\mu_2^{(2)} = -\frac{\lambda^4}{8(\nu_1 \lambda + \gamma^2)^2}$.

It then follows that

$$\sinh \mu_j a = \sinh |\xi'| a + \frac{e_j^{(1)}}{|\xi'|} \cosh |\xi'| a + \frac{e_j^{(2)}}{|\xi'|^2} \sinh |\xi'| a + \frac{e_j^{(3)}}{|\xi'|^3} \cosh |\xi'| a + O\left(|\lambda|^3 |\xi'|^{-4} e^{|\xi'|a}\right)$$

and

$$\cosh \mu_{j}a = \cosh |\xi'|a + \frac{e_{j}^{(1)}}{|\xi'|} \sinh |\xi'|a + \frac{e_{j}^{(2)}}{|\xi'|^{2}} \cosh |\xi'|a + \frac{e_{j}^{(3)}}{|\xi'|^{3}} \sinh |\xi'|a + O\left(|\lambda|^{3}|\xi'|^{-4}e^{|\xi'|a}\right),$$

where j = 1, 2,

$$e_1^{(1)} = \frac{\lambda a}{2\nu}, \quad e_2^{(1)} = \frac{\lambda^2 a}{2(\nu_1 \lambda + \gamma^2)}, \quad e_1^{(2)} = \frac{\lambda^2 a^2}{8\nu^2}, \quad e_2^{(2)} = \frac{\lambda^4 a^2}{8(\nu_1 \lambda + \gamma^2)^2}$$

and

$$e_1^{(3)} = O(|\lambda|^2), \quad e_2^{(3)} = O(|\lambda|^4).$$

We thus obtain

(4.15)
$$\frac{1}{\lambda^2} D(\lambda, \xi') = -\frac{a^2}{4\nu^2} + d_0(\lambda) + \left(\frac{1}{4\nu^2} + d_1(\lambda)\right) \left(\frac{\sinh|\xi'|a}{|\xi'|}\right)^2 + d_2(\lambda, \xi'),$$

where $d_0(\lambda)$, $d_1(\lambda)$ and $d_2(\lambda, \xi')$ are some functions satisfying

(4.16)
$$|d_0(\lambda)| \le C|\lambda|, \quad |d_1(\lambda)| \le C|\lambda|, \quad |d_2(\lambda,\xi')| \le C \frac{|\lambda|}{|\xi'|^3} e^{2|\xi'|a}$$

uniformly in λ and ξ' with $|\lambda| \leq \Lambda$ and $|\xi'| \geq r$ for some $C = C(r_1) > 0$. Let $r_1 > 0$ and $R_1 > 0$ be fixed. Since

$$\left(\frac{\sinh|\xi'|a}{|\xi'|}\right)^2 - a^2 = \left(\frac{\sinh|\xi'|a}{|\xi'|} + a\right) \left(\frac{\sinh|\xi'|a}{|\xi'|} - a\right) \ge \frac{a^4}{3} |\xi'|^2$$

for $|\xi'| > 0$, we see from (4.15) and (4.16) that if $r_1 \le |\xi'| \le R_1$, then

$$\left|\frac{1}{\lambda^2}D(\lambda,\xi')\right| \ge \frac{a^4}{12\nu^2}|\xi'|^2 - C|\lambda| \left\{1 + \frac{e^{2R_1a}}{|\xi'|^3}\right\},\,$$

provided that $|\lambda| \leq \Lambda_1$ for some Λ_1 satisfying (4.14) with $\Lambda = \Lambda_1$ and $r = r_1$. Here $C = C(r_1) > 0$. Therefore, there exists a constant $\Lambda_1 = \Lambda(r_1, R_1) > 0$ such that

$$|D(\lambda,\xi')| \ge \frac{a^4}{24\nu^2} r_1^2 |\lambda|^2$$

for λ and ξ' with $|\lambda| \leq \Lambda_1$ and $r_1 \leq |\xi'| \leq R_1$. This proves (i).

We next prove (ii). From (4.16) we see that there exists a positive number Λ_2 such that

$$\left|\frac{1}{4\nu^2} + d_1(\lambda)\right| \ge \frac{1}{8\nu^2}$$

for $|\lambda| \leq \Lambda_2$. Furthermore, there exists a positive number $R_2 = R_2(\Lambda_2)$ such that (4.14) is satisfied with $\Lambda = \Lambda_2$ and $r = R_2$, and the inequalities

$$\left| -\frac{a^2}{4\nu^2} + d_0(\lambda) \right| \le \frac{1}{32\nu^2} \left(\frac{\sinh |\xi'|a}{|\xi'|} \right)^2, \quad |d_2(\lambda,\xi')| \le \frac{1}{32\nu^2} \left(\frac{\sinh |\xi'|a}{|\xi'|} \right)^2$$

hold for all $|\lambda| \leq \Lambda_2$ and $|\xi'| \geq R_2$. It then follows from (4.15) that

$$\left|\frac{1}{\lambda^2}D(\lambda,\xi')\right| \ge \frac{1}{16\nu^2} \left(\frac{\sinh|\xi'|a}{|\xi'|}\right)^2$$

for all $|\lambda| \leq \Lambda_2$ and $|\xi'| \geq R_2$. This completes the proof.

By Propositions 4.1 and 4.2 we have the following consequence on $\rho(-L_{\xi'})$.

Lemma 4.3. (i) There exists a number $\theta_1 \in (\frac{\pi}{2}, \pi)$ such that $\Sigma(\eta, \theta_1) \subset \rho(-\hat{L}_{\xi'})$ for any $\eta > 0$ and $\xi' \in \mathbf{R}^{n-1}$. Furthermore, the integral representation of $(\lambda + \hat{L}_{\xi'})^{-1}$ in Theorem 3.8 holds for $\lambda \in \Sigma(\eta, \theta_1)$ for any $\eta > 0$ and $\xi' \in \mathbf{R}^{n-1}$.

(ii) For any r > 0 there exist positive numbers η_2 and θ_2 with $\theta_2 \in (\frac{\pi}{2}, \pi)$ such that $\Sigma(-\eta_2, \theta_2) \subset \rho(-\hat{L}_{\xi'})$ for $|\xi'| \geq r$. Furthermore, the integral representation of $(\lambda + \hat{L}_{\xi'})^{-1}$ in Theorem 3.8 holds for $\lambda \in \Sigma(-\eta_2, \theta_2)$ and $|\xi'| \geq r$.

Proof. The first assertion of (i) is an easy consequence of Proposition 4.1. By Remarks 3.2 and 3.5, changing θ_1 suitably if necessary, we see that $\lambda_{1,k}, \lambda_{\pm,k} \notin \Sigma(\eta, \theta_1)$ for any $k = 0, 1, 2, \cdots$. Therefore, the integral representation in Theorem 3.8 holds for $\lambda \in \Sigma(\eta, \theta_1)$.

Similarly, we see from Lemma 3.4 and Propositions 4.1 and 4.2 that for any r > 0 there exist positive numbers η_2 and θ_2 with $\theta_2 \in (\frac{\pi}{2}, \pi)$ such that $\Sigma(-\eta_2, \theta_2) - \{0\} \subset \rho(-\hat{L}_{\xi'})$ for $|\xi'| \geq r$. Furthermore, by Remarks 3.2 and 3.5, changing η_2 and θ_2 suitably if necessary, we deduce that $\lambda_{1,k}, \lambda_{\pm,k} \notin$ $\Sigma(-\eta_2, \theta_2)$ for any $k = 0, 1, 2, \cdots$, and the integral representation in Theorem 3.8 holds for $\lambda \in \Sigma(-\eta_2, \theta_2) - \{0\}$ and $|\xi'| \geq r$.

Let us prove $0 \in \rho(-\widehat{L}_{\xi'})$. Assume that ξ' satisfies $|\xi'| \geq r$. We note that $(\lambda + \widehat{L}_{\xi'})^{-1}$ is analytic in $\Sigma(-\eta_2, \theta_2) - \{0\}$. Furthermore, $\widehat{G}(\lambda, \xi')$ and $\widehat{K}(\lambda, \xi')$ are also analytic in $\Sigma(-\eta_2, \theta_2) - \{0\}$. Therefore, it suffices to prove that $\widehat{G}(\lambda, \xi')$ and $\widehat{K}(\lambda, \xi')$ are bounded in $\Sigma(-\eta_2, \theta_2) - \{0\}$.

Since

(4.17)
$$\mu_1 - \mu_2 = \frac{\lambda}{\mu_1 + \mu_2} \left\{ \frac{1}{\nu} + \frac{\lambda}{\nu_1 \lambda + \gamma^2} \right\},$$

we see that $\tilde{Q}\hat{G}(\lambda,\xi')\tilde{Q}$ is bounded in $\Sigma(-\eta_2,\theta_2) - \{0\}$. It is easy to see that the other components of $\hat{G}(\lambda,\xi')$ are bounded in $\Sigma(-\eta_2,\theta_2) - \{0\}$, and hence, $\hat{G}(\lambda,\xi')$ is bounded in $\Sigma(-\eta_2,\theta_2) - \{0\}$.

As for $K(\lambda, \xi')$, we easily see that $b_j(\lambda, \xi', y_n) = O(\lambda)$ as $\lambda \to 0$ for j = 1, 2, 3. This, together with Proposition 4.2, implies that $\beta_j(\lambda, \xi', y_n) = O(1)$ $(j = 0, 1), B'(\lambda, \xi', y_n) = O(1)$ and $\mathbf{b}_n(\lambda, \xi', y_n) = O(1)$ as $\lambda \to 0$. Furthermore, using (4.17), we also see that $h_{\mu_1,\mu_2}(x_n) = O(\lambda)$ as $\lambda \to 0$. It then follows that $\widehat{K}(\lambda, \xi')$ is bounded in $\Sigma(-\eta_2, \theta_2) - \{0\}$ and assertion (ii) is proved. This completes the proof. We next derive estimates for $|D(\lambda,\xi')|$ from below when $|\lambda| + |\xi'|^2$ is large. In the following we specify branches of $\mu_j(\lambda,\xi')$, j = 1, 2, as a function of λ . As for $\mu_1(\lambda,\xi')$ we take the principal branch of the square root of $\frac{\lambda+\nu|\xi'|^2}{\nu}$, i.e., for $\lambda \notin (-\infty, -\nu|\xi'|^2]$ we denote by $\mu_1(\lambda,\xi')$ the square root of $\frac{\lambda+\nu|\xi'|^2}{\nu}$ with $\operatorname{Re} \mu_1(\lambda,\xi') > 0$. As for $\mu_2(\lambda,\xi')$ we take the branch in the following way. When $|\xi'| < \frac{2\gamma}{\nu_1}$, we use the branch specified by the requirement

$$\arg(\lambda - \lambda_{\pm,0}) = \mp \frac{\pi}{2} \text{ at } \lambda = \operatorname{Re} \lambda_{+,0} \text{ and } \arg(\lambda + \frac{\gamma^2}{\nu_1}) = 0 \text{ at } \lambda = 0$$

and take the branch cut

$$\{\lambda; \operatorname{Re} \lambda \leq -\frac{\gamma^2}{\nu_1}, \operatorname{Im} \lambda = 0\} \cup \{\lambda \in \Gamma_{\gamma,\nu_1}; \operatorname{Re} \lambda \leq \operatorname{Re} \lambda_{\pm,0}\}.$$

Here Γ_{γ,ν_1} is the circle defined by $\Gamma_{\gamma,\nu_1} = \{\lambda; \left|\lambda + \frac{\gamma^2}{\nu_1}\right| = \frac{\gamma^2}{\nu_1}\}$. When $|\xi'| \ge \frac{2\gamma}{\nu_1}$, we use the branch specified by $\arg(\lambda - \lambda_{\pm,0}) = \arg(\lambda + \frac{\gamma^2}{\nu_1}) = 0$ at $\lambda = 0$ and take the branch cut

$$\{\lambda; \operatorname{Re} \lambda \leq -\frac{\gamma^2}{\nu_1}, \operatorname{Im} \lambda = 0\}$$

As shown in [6], it holds $\operatorname{Re} \mu_2(\lambda, \xi') > 0$ for λ outside the branch cut.

We introduce a function $D_1(\lambda, \xi')$ defined by

$$D_1(\lambda, \xi') = e^{-(\mu_1 + \mu_2)a} D(\lambda, \xi').$$

To investigate the resolvent $(\lambda + L)^{-1} = \mathscr{F}^{-1} \left[(\lambda + L'_{\xi})^{-1} \right]$ for large $\lambda + |\xi'|^2$, it is convenient to consider $D_1(\lambda, \xi')$ rather than $D(\lambda, \xi')$.

Lemma 4.4. (i) There are positive numbers η_3 and θ_3 with $\theta_3 \in (\frac{\pi}{2}, \pi)$ such that the inequality

$$|D_1(\lambda,\xi')| \ge C \frac{|\lambda|^2}{|\lambda| + 1 + |\xi'|}$$

holds uniformly in $\lambda \in \Sigma(-\eta_3, \theta_3) \cap \{\lambda; |\lambda| \ge \delta\}$ and $|\xi'| \ge R_3$ for any $\delta > 0$ with some constant $R_3 = R_3(\delta) > 0$.

(ii) There are positive numbers η_4 and θ_4 with $\theta_4 \in (\frac{\pi}{2}, \pi)$ such that the inequality

$$|D_1(\lambda, \xi')| \ge C \frac{|\lambda|^2}{|\lambda| + 1 + |\xi'|^2}$$

holds uniformly in $\lambda \in \Sigma(\eta_4, \theta_4)$ and $\xi' \in \mathbf{R}^{n-1}$.

Proof. We first observe that there are positive numbers η_3 , $\theta_3 \in (\frac{\pi}{2}, \pi)$ and \widetilde{R}_3 such that

(4.18)
$$\operatorname{Re} \mu_j \ge C(|\lambda| + 1 + |\xi'|^2)^{\frac{1}{2}} \quad (j = 1, 2)$$

for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \geq \tilde{R}_3$. Furthermore, there exist positive numbers $\tilde{\eta}_4$ and $\theta_4 \in (\frac{\pi}{2}, \pi)$ such that (4.18) also holds for $\lambda \in \Sigma(\tilde{\eta}_4, \theta_4)$ and $\xi' \in \mathbf{R}^{n-1}$. In fact, it is easy to show the inequalities (4.18) for j = 1 with appropriate η_3 , $\tilde{\eta}_4$, θ_3 , θ_4 and \tilde{R}_3 . As for j = 2 one can find (4.18) by using the observation in Remark 3.2. It is also possible to see that

(4.19)
$$|\mu_j| \le C(|\lambda| + 1 + |\xi'|^2)^{\frac{1}{2}}$$

for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and ξ' with $|\xi'| \ge \widetilde{R}_3$, and for $\lambda \in \Sigma(\widetilde{\eta}_4, \theta_4)$ and $\xi' \in \mathbf{R}^{n-1}$.

Consider next the quadratic equation $\omega^2 + (2\nu + \tilde{\nu})|\xi'|^2\omega + \gamma^2|\xi'|^2 = 0$ for ω . This equation has two roots

$$\omega_{\pm} = -\frac{1}{2}(2\nu + \tilde{\nu})|\xi'|^2 \pm \frac{1}{2}\sqrt{(2\nu + \tilde{\nu})^2|\xi'|^4 - 4\gamma^2|\xi'|^2}.$$

Therefore, as in Remark 3.2, we see that

$$\omega_{\pm} = -\frac{1}{2}(2\nu + \tilde{\nu})|\xi'|^2 \pm i\gamma|\xi'| + O(|\xi'|^3) \text{ as } |\xi'| \to 0,$$

and $\omega_{\pm} \in \mathbf{R}$ for $|\xi'| \geq \frac{2\gamma}{2\nu + \tilde{\nu}}$, and

$$\omega_{+} = -\frac{\gamma^{2}}{2\nu + \tilde{\nu}} + O(|\xi'|^{-2}), \quad \omega_{-} = -(2\nu + \tilde{\nu})|\xi'|^{2} + O(1) \quad \text{as} \quad |\xi'| \to \infty.$$

We thus deduce, by suitably changing $\theta_3, \theta_4, \widetilde{R}_3, \eta_3$ and $\widetilde{\eta}_4$ if necessary, that

(4.20)
$$\begin{aligned} |\lambda^2 + (2\nu + \tilde{\nu})|\xi'|^2\lambda + \gamma^2|\xi'|^2| &= |(\lambda - \omega_+)(\lambda - \omega_-)| \\ &\geq C(|\lambda| + 1 + |\xi'|^2)(|\lambda| + 1) \end{aligned}$$

for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and ξ' with $|\xi'| \ge \tilde{R}_3$, and for $\lambda \in \Sigma(\tilde{\eta}_4, \theta_4)$ and $\xi' \in \mathbf{R}^{n-1}$. We now estimate $D_1(\lambda, \xi')$. We write $D_1(\lambda, \xi')$ as

(4.21)
$$D_1(\lambda,\xi') = \frac{1}{4} \left(\frac{|\xi'|^4}{\mu_1 \mu_2} - 2|\xi'|^2 + \mu_1 \mu_2 \right) + \widetilde{D}_1(\lambda,\xi'),$$

where (4.22)

$$\widetilde{D}_{1}(\lambda,\xi') = \frac{1}{4} \left\{ 8|\xi'|^{2} e^{-(\mu_{1}+\mu_{2})a} + \left(\frac{|\xi'|^{4}}{\mu_{1}\mu_{2}} - 2|\xi'|^{2} + \mu_{1}\mu_{2}\right) e^{-2(\mu_{1}+\mu_{2})a} - \left(\frac{|\xi'|^{4}}{\mu_{1}\mu_{2}} + 2|\xi'|^{2} + \mu_{1}\mu_{2}\right) \left(e^{-2\mu_{1}a} + e^{-2\mu_{2}a}\right) \right\}.$$

Since

$$\frac{|\xi'|^4}{\mu_1\mu_2} - 2|\xi'|^2 + \mu_1\mu_2 = \left|\frac{1}{\mu_1\mu_2}\right| \left|\frac{(\mu_1\mu_2)^2 - |\xi'|^4}{\mu_1\mu_2 + |\xi'|^2}\right|^2$$

and

$$(\mu_1 \mu_2)^2 - |\xi'|^4 = \frac{\lambda \{\lambda^2 + (2\nu + \tilde{\nu}) |\xi'|^2 \lambda + \gamma^2 |\xi'|^2\}}{\nu_1 \lambda + \gamma^2},$$

we see from (4.19) and (4.20) that

(4.23)
$$\frac{1}{4} \left| \frac{|\xi'|^4}{\mu_1 \mu_2} - 2|\xi'|^2 + \mu_1 \mu_2 \right| \ge C_0 \frac{|\lambda|^2}{|\lambda| + 1 + |\xi'|^2}$$

uniformly for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and ξ' with $|\xi'| \ge \widetilde{R}_3$, and for $\lambda \in \Sigma(\widetilde{\eta}_4, \theta_4)$ and $\xi' \in \mathbf{R}^{n-1}$.

We also see from (4.18) and (4.19) that

$$\left|e^{-\mu_{j}a}\right| \le C_{k} \left(\operatorname{Re} \mu_{j}a\right)^{-k} \le C_{k} \left(|\lambda|+1+|\xi'|^{2}\right)^{-\frac{k}{2}}$$

for any k. This, together with (4.22), implies that for any $\delta > 0$ there exists a positive number $R_3 = R_3(\delta)$ with $R_3 \ge \tilde{R}_3$ such that if $\lambda \in \Sigma(-\eta_3, \theta_3) \cap \{\lambda; |\lambda| \ge \delta\}$ and $|\xi'| \ge R_3$, then

$$\left|\widetilde{D}_{1}(\lambda,\xi')\right| \leq \frac{C|\lambda|^{2}}{\delta^{2}(|\lambda|+1+|\xi'|^{2})^{2}} \leq \frac{C_{0}}{2} \frac{|\lambda|^{2}}{|\lambda|+1+|\xi'|^{2}}.$$

Combining this with (4.23), we have

$$|D_1(\lambda,\xi')| \ge \frac{C_0}{2} \frac{|\lambda|^2}{|\lambda| + 1 + |\xi'|^2}$$

for $\lambda \in \Sigma(-\eta_3, \theta_3) \cap \{\lambda; |\lambda| \ge \delta\}$ and ξ' with $|\xi'| \ge R_3$. This proves (i). We can also find a positive number η_4 with $\eta_4 \ge \tilde{\eta}_4$ such that

$$\left|\widetilde{D}_{1}(\lambda,\xi')\right| \leq \frac{C|\lambda|^{2}}{\eta_{4}^{2}(|\lambda|+1+|\xi'|^{2})^{2}} \leq \frac{C_{0}}{2} \frac{|\lambda|^{2}}{|\lambda|+1+|\xi'|^{2}}$$

for all $\lambda \in \Sigma(\eta_4, \theta_4)$ and $\xi' \in \mathbf{R}^{n-1}$, and hence,

$$|D_1(\lambda,\xi')| \ge \frac{C_0}{2} \frac{|\lambda|^2}{|\lambda| + 1 + |\xi'|^2}$$

for all such λ and ξ' . This completes the proof.

To estimate each component of the integral kernel in Theorem 3.8, we will use the following lemma.

Lemma 4.5. Let $\alpha' \in \mathbb{Z}^{n-1}$ be any multi-index with $|\alpha'| \leq n$ and let θ_4 be the number given in Lemma 4.4. Then there exists a positive number η_4 such that the following estimates hold uniformly in $\lambda \in \Sigma(\eta_4, \theta_4)$ and $\xi' \in \mathbb{R}^{n-1}$:

(i)
$$\left|\partial_{\xi'}^{\alpha'}\mu_{j}\right| \leq C(|\lambda|+1+|\xi'|^{2})^{\frac{1}{2}-\frac{|\alpha'|}{2}} \quad (j=1,2),$$

(ii)
$$\left| \partial_{\xi'}^{\alpha'} \left(\mu_1 - \mu_2 \right) \right| \le C |\lambda| (|\lambda| + 1 + |\xi'|^2)^{-\frac{1}{2} - \frac{|\alpha'|}{2}},$$

(iii)
$$\left| \partial_{\xi'}^{\alpha'} \left(\mu_1 \mu_2 - |\xi'|^2 \right) \right| \le C |\lambda| (|\lambda| + 1 + |\xi'|^2)^{-\frac{|\alpha'|}{2}},$$

(iv)
$$\left|\partial_{\xi'}^{\alpha'} e^{-\mu_j x_n}\right| \le C(|\lambda| + 1 + |\xi'|^2)^{-\frac{|\alpha'|}{2}} e^{-\frac{1}{2}\operatorname{Re}\mu_j x_n} \quad (j = 1, 2)$$

for all $0 \leq x_n \leq a$,

(v)
$$\left| \partial_{\xi'}^{\alpha'} \left(\frac{1}{D_1(\lambda,\xi')} \right) \right| \leq C |\lambda|^{-2} (|\lambda| + 1 + |\xi'|^2)^{1 - \frac{|\alpha'|}{2}},$$

(vi)
$$\begin{aligned} & \left| \partial_{\xi'}^{\alpha'} \left(e^{-\mu_1 x_n} - e^{-\mu_2 x_n} \right) \right| \\ & \leq C |\lambda| (|\lambda| + 1 + |\xi'|^2)^{-1 - \frac{|\alpha'|}{2}} \left\{ e^{-\frac{1}{8} \operatorname{Re} \mu_1 x_n} + e^{-\frac{1}{8} \operatorname{Re} \mu_2 x_n} \right\} \end{aligned}$$

for all $0 \leq x_n \leq a$.

The inequalities (i)–(vi) also hold uniformly in $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$ for some $R_3 > 0$, where η_3 and θ_3 are the numbers given in Lemma 4.4.

Proof. Let η_4 and θ_4 are the numbers given in Lemma 4.4 (ii). Then a direct calculation gives the inequality in (i). The inequality in (ii) follows from (i), (4.17) and (4.18).

Let us prove (iii). Suppose first that $|\lambda| \leq C_0 |\xi'|^2$, where C_0 is a positive constant to be determined later. We write $\mu_1 \mu_2 - |\xi'|^2$ as

(4.24)
$$\mu_1 \mu_2 - |\xi'|^2 = \frac{(\mu_1 \mu_2)^2 - |\xi'|^4}{\mu_1 \mu_2 + |\xi'|^2}.$$

Since

$$(\mu_1\mu_2)^2 - |\xi'|^4 = \frac{\lambda^3}{\nu(\nu_1\lambda + \gamma^2)} + \lambda\left(\frac{1}{\nu} + \frac{\lambda}{\nu_1\lambda + \gamma^2}\right)|\xi'|^2,$$

we see that

(4.25)
$$\left| \partial_{\xi'}^{\alpha'} \left((\mu_1 \mu_2)^2 - |\xi'|^4 \right) \right| \le C |\lambda| (|\lambda| + |\xi'|^2)^{1 - \frac{|\alpha'|}{2}}.$$

On the other hand, since $\left|\frac{\lambda^2}{\nu_1\lambda+\gamma^2}\right| \leq C|\lambda|$ for $\lambda \in \Sigma(\eta_4, \theta_4)$, taking C_0 sufficiently small, we find that

$$\left|\frac{\lambda}{\nu}\right| \le \frac{|\xi'|^2}{2}, \quad \left|\frac{\lambda^2}{\nu_1\lambda + \gamma^2}\right| \le \frac{|\xi'|^2}{2}$$

for λ and ξ' with $|\lambda| \leq C_0 |\xi'|^2$. Applying the mean value theorem to $\mu_1 = |\xi'| \sqrt{1 + \frac{\lambda}{\nu |\xi'|^2}}$ and $\mu_2 = |\xi'| \sqrt{1 + \frac{\lambda^2}{(\nu_1 \lambda + \gamma^2) |\xi'|^2}}$, we obtain

$$\mu_1 \mu_2 + |\xi'|^2 = 2|\xi'|^2 + \frac{\lambda}{\nu} q_1(\lambda,\xi') + \frac{\lambda^2}{\nu_1 \lambda + \gamma^2} q_2(\lambda,\xi') + \frac{\lambda^3}{\nu(\nu_1 \lambda + \gamma^2)|\xi'|^2} q_3(\lambda,\xi'),$$

where $q_j(\lambda, \xi'), j = 1, 2, 3$, are some functions satisfying

$$|q_j(\lambda, \xi')| \le C, \quad j = 1, 2, 3,$$

for λ and ξ' with $|\lambda| \leq C_0 |\xi'|^2$. Therefore, taking C_0 smaller if necessary, we see that

(4.26)
$$\left| \mu_1 \mu_2 + |\xi'|^2 \right| \ge 2|\xi'|^2 - C|\lambda| \ge |\xi'|^2 \ge C(|\lambda| + 1 + |\xi'|^2)$$

uniformly for $\lambda \in \Sigma(\eta_4, \theta_4)$ and ξ' with $|\lambda| \leq C_0 |\xi'|^2$. It then follows from (i) and (4.24)–(4.26) that

$$\left|\partial_{\xi'}^{\alpha'} \left(\mu_1 \mu_2 - |\xi'|^2\right)\right| \le C |\lambda| (|\lambda| + 1 + |\xi'|^2)^{-\frac{|\alpha'|}{2}}$$

uniformly for $\lambda \in \Sigma(\eta_4, \theta_4)$ and ξ' with $|\lambda| \leq C_0 |\xi'|^2$. As for the case $|\lambda| \geq C_0 |\xi'|^2$, we easily see that

$$\left|\partial_{\xi'}^{\alpha'} \left(\mu_1 \mu_2 - |\xi'|^2\right)\right| \le C(|\lambda| + |\xi'|^2)^{1 - \frac{|\alpha'|}{2}} \le C|\lambda|(|\lambda| + 1 + |\xi'|^2)^{-\frac{|\alpha'|}{2}}.$$

This proves (iii).

We next consider (iv). The case $\alpha' = 0$ is trivial. Let $|\alpha'| \ge 1$. Then, by (i) and (4.18), we have

$$\begin{aligned} \partial_{\xi'}^{\alpha'} e^{-\mu_{j}x_{n}} \Big| &\leq C \sum_{\ell=1}^{|\alpha'|} e^{-\operatorname{Re}\mu_{j}x_{n}} \prod_{\alpha_{1}+\dots+\alpha_{\ell}=\alpha'} \left| \partial_{\xi'}^{\alpha_{1}}\mu_{j}x_{n} \right| \cdots \left| \partial_{\xi'}^{\alpha_{\ell}}\mu_{j}x_{n} \right| \\ &\leq C \sum_{\ell=1}^{|\alpha'|} x_{n}^{\ell} e^{-\operatorname{Re}\mu_{j}x_{n}} \left(|\lambda|+1+|\xi'|^{2} \right)^{\frac{\ell}{2}-\frac{|\alpha'|}{2}} \\ &\leq C \sum_{\ell=1}^{|\alpha'|} (\operatorname{Re}\mu_{j})^{-\ell} e^{-\frac{1}{2}\operatorname{Re}\mu_{j}x_{n}} \left(|\lambda|+1+|\xi'|^{2} \right)^{\frac{\ell}{2}-\frac{|\alpha'|}{2}} \\ &\leq C \left(|\lambda|+1+|\xi'|^{2} \right)^{-\frac{|\alpha'|}{2}} e^{-\frac{1}{2}\operatorname{Re}\mu_{j}x_{n}}. \end{aligned}$$

This shows (iv).

As for (v), since $\frac{|\xi'|^4}{\mu_1\mu_2} - 2|\xi'|^2 + \mu_1\mu_2 = \frac{1}{\mu_1\mu_2}(\mu_1\mu_2 - |\xi'|^2)^2$, we see from (i) and (iii) that

$$\left|\partial_{\xi'}^{\alpha'}\left(\frac{|\xi'|^4}{\mu_1\mu_2} - 2|\xi'|^2 + \mu_1\mu_2\right)\right| \le C|\lambda|^2(|\lambda| + 1 + |\xi'|^2)^{-1 - \frac{|\alpha'|}{2}}.$$

This, together with (4.21) and (4.22), implies that

(4.27)
$$\left| \partial_{\xi'}^{\alpha'} D_1(\lambda, \xi') \right| \le C |\lambda|^2 (|\lambda| + 1 + |\xi'|^2)^{-1 - \frac{|\alpha'|}{2}}$$

Combining Lemma 4.4 (ii) and (4.27), we obtain the desired inequality in (v).

We finally prove (vi). Since

$$e^{-\mu_1 x_n} - e^{-\mu_2 x_n} = (\mu_2 - \mu_1) x_n \int_0^1 e^{-\{\theta \mu_1 + (1-\theta)\mu_2\} x_n} d\theta,$$

we see from (ii) and (iv) that

$$\begin{aligned} \left| \partial_{\xi'}^{\alpha'} \left(e^{-\mu_{1}x_{n}} - e^{-\mu_{2}x_{n}} \right) \right| \\ &\leq Cx_{n} |\lambda| (|\lambda| + 1 + |\xi'|^{2})^{-\frac{1}{2} - \frac{|\alpha'|}{2}} \int_{0}^{1} e^{-\frac{1}{2}\operatorname{Re}\left\{\theta\mu_{1} + (1-\theta)\mu_{2}\right\}x_{n}} d\theta \\ &\leq Cx_{n} |\lambda| (|\lambda| + 1 + |\xi'|^{2})^{-\frac{1}{2} - \frac{|\alpha'|}{2}} \left(e^{-\frac{1}{4}\operatorname{Re}\mu_{1}x_{n}} + e^{-\frac{1}{4}\operatorname{Re}\mu_{2}x_{n}} \right) \\ &\leq C |\lambda| (|\lambda| + 1 + |\xi'|^{2})^{-1 - \frac{|\alpha'|}{2}} \left(e^{-\frac{1}{8}\operatorname{Re}\mu_{1}x_{n}} + e^{-\frac{1}{8}\operatorname{Re}\mu_{2}x_{n}} \right). \end{aligned}$$

The desired inequalities are thus proved for $\lambda \in \Sigma(\eta_4, \theta_4)$.

Let us consider the case $\lambda \in \Sigma(-\eta_3, \theta_3)$. In the same way as above, one can show the inequalities (i)–(iv) and (vi) for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$, where R_3 is the positive number given in Lemma 4.4 (i). As for (v), we take $\delta = \Lambda_2$ in Lemma 4.4 (i) with Λ_2 being the number given in Proposition 4.2 (ii). Similarly to the case $\lambda \in \Sigma(\eta_4, \theta_4)$, by using Lemma 4.4 (i), we can also prove the inequality (v) for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$, provided that $|\lambda| \ge \delta = \Lambda_2$. In case $|\lambda| \le \Lambda_2$, since $e^{-(\mu_1 + \mu_2)a} e^{2|\xi'|a} = e^{O(|\xi'|^{-1})}$, we see from Proposition 4.2 (ii) that

$$|D_1(\lambda,\xi')| \ge C \frac{|\lambda|^2}{|\xi'|^2}$$

for large $|\xi'|$. Furthermore, similarly to the case $\lambda \in \Sigma(\eta_4, \theta_4)$, one can obtain (4.27) with a more detailed computation for $\widetilde{D}_1(\lambda, \xi')$ given in (4.22)

for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and large $|\xi'|$. The desired inequality (v) then follows by changing R_3 suitably large if necessary. This completes the proof.

We next derive the estimates for the integral kernel of $\widehat{G}(\lambda,\xi')$. We set

$$(4.28) \begin{cases} g_{\mu_{j}}^{(1)}(x_{n}, y_{n}) = \frac{1}{2\mu_{j}}e^{-\mu_{j}|x_{n}-y_{n}|}, \\ g_{\mu_{j}}^{(2)}(x_{n}, y_{n}) = \frac{1}{2\mu_{j}}\left(e^{-\mu_{j}(x_{n}+y_{n})} + e^{-\mu_{j}\{(a-x_{n})+(a-y_{n})\}}\right), \\ g_{\mu_{j}}^{(3)}(x_{n}, y_{n}) = \frac{1}{2\mu_{j}}\frac{e^{-2\mu_{j}a}}{1-e^{-2\mu_{j}}}\left(e^{-\mu_{j}|x_{n}-y_{n}|} + e^{\mu_{j}|x_{n}-y_{n}|}\right), \\ g_{\mu_{j}}^{(4)}(x_{n}, y_{n}) = \frac{1}{2\mu_{j}}\frac{e^{-2\mu_{j}a}}{1-e^{-2\mu_{j}}}\left(e^{-\mu_{j}(x_{n}+y_{n})} + e^{-\mu_{j}\{(a-x_{n})+(a-y_{n})\}}\right), \end{cases}$$

where $\mu_j = \mu_j(\lambda, \xi'), \ j = 1, 2$. In what follows we will denote $|\lambda| + 1 + |\xi'|^2$ by $\sigma(\lambda, \xi')$:

$$\sigma(\lambda,\xi') = |\lambda| + 1 + |\xi'|^2.$$

Lemma 4.6. Let $g_{\mu_j}^{(k)}(x_n, y_n)$, $j = 1, 2, k = 1, \dots, 4$, be defined in (4.28). Then

$$g_{\mu_j}^D(x_n, y_n) = \sum_{k=1}^4 (-1)^{k+1} g_{\mu_j}^{(k)}(x_n, y_n), \quad g_{\mu_j}^N(x_n, y_n) = \sum_{k=1}^4 g_{\mu_j}^{(k)}(x_n, y_n),$$

and the following estimates hold for any multi-index $\alpha' \in \mathbb{Z}^{n-1}$ with $|\alpha'| \leq [\frac{n-1}{2}]+1$ and any nonnegative integer ℓ uniformly in $\lambda \in \Sigma(\eta_4, \theta_4), \xi' \in \mathbb{R}^{n-1}$ and $x_n, y_n \in [0, a]$:

(i)
$$\left|\partial_{\xi'}^{\alpha'}\left[\left(\partial_{x_n}^{\ell}g_{\mu_j}^{(k)}\right)(x_n, y_n)\right]\right| \le C_{\ell}\sigma(\lambda, \xi')^{\frac{\ell}{2} - \frac{1}{2} - \frac{|\alpha'|}{2}}E^{(k)}(x_n, y_n)$$

for k = 1, 2 and j = 1, 2, and

(ii)
$$\begin{aligned} \left| \partial_{\xi'}^{\alpha'} \left[(\partial_{x_n}^{\ell} g_{\mu_1}^{(k)})(x_n, y_n) - (\partial_{x_n}^{\ell} g_{\mu_2}^{(k)})(x_n, y_n) \right] \right| \\ &\leq C |\lambda| \sigma(\lambda, \xi')^{\frac{\ell}{2} - \frac{3}{2} - \frac{|\alpha'|}{2}} E^{(k)}(x_n, y_n), \end{aligned}$$

for k = 1, 2. Here $E^{(k)}(x_n, y_n)$, k = 1, 2, are the functions defined by

$$E^{(1)}(x_n, y_n) = e^{-c\sigma(\lambda, \xi')^{\frac{1}{2}}|x_n - y_n|},$$

$$E^{(2)}(x_n, y_n) = e^{-c\sigma(\lambda, \xi')^{\frac{1}{2}}(x_n + y_n)} + e^{-c\sigma(\lambda, \xi')^{\frac{1}{2}}\{(a - x_n) + (a - y_n)\}}$$

with some constant c > 0 independent of $\lambda \in \Sigma(\eta_4, \theta_4)$, $\xi' \in \mathbb{R}^{n-1}$ and $x_n, y_n \in [0, a]$. Furthermore, for any nonnegative integer q, there hold the estimates

(iii)
$$\left|\partial_{\xi'}^{\alpha'}\left[\left(\partial_{x_n}^{\ell}g_{\mu_j}^{(k)}\right)(x_n, y_n)\right]\right| \le C_{\ell,q}\sigma(\lambda, \xi')^{-\frac{q+|\alpha'|}{2}}, \quad k = 3, 4, \ j = 1, 2.$$

The inequalities (i)–(iii) also hold uniformly in $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$ for some $R_3 > 0$, where η_3 and θ_3 are the numbers given in Lemma 4.4.

Proof. A direct computation shows that $g^M_{\mu_j}(x_n, y_n)$, M = D, N, are written as above. One can prove inequalities (i) and (iii) by a direct application of Lemma 4.5.

As for (ii), we write $g_{\mu_1}^{(1)}(x_n, y_n) - g_{\mu_2}^{(1)}(x_n, y_n)$ as

$$g_{\mu_1}^{(1)}(x_n, y_n) - g_{\mu_2}^{(1)}(x_n, y_n) = \frac{1}{2} \left(\frac{1}{\mu_1} - \frac{1}{\mu_2} \right) e^{-\mu_1 |x_n - y_n|} + \frac{1}{2\mu_2} \left(e^{-\mu_1 |x_n - y_n|} - e^{-\mu_2 |x_n - y_n|} \right).$$

Since $\partial_{x_n}^{\ell} e^{-\mu_j |x_n - y_n|} = (-\mu_j \operatorname{sgn} (x_n - y_n))^{\ell} e^{-\mu_j |x_n - y_n|}$, we can obtain inequality (ii) for k = 1. The case k = 2 can be proved similarly. This completes the proof.

In the same manner we can estimate $h_{\mu_j}(x_n)$.

Lemma 4.7. Let $\alpha' \in \mathbb{Z}^{n-1}$ be a multi-index with $|\alpha'| \leq \left[\frac{n-1}{2}\right] + 1$ and let ℓ be a nonnegative integer. Then the following estimates hold uniformly in $\lambda \in \Sigma(\eta_4, \theta_4), \xi'$ and $x_n \in [0, a]$:

(i)
$$\left|\partial_{\xi'}^{\alpha'}\partial_{x_n}^{\ell}h_{\mu_j}(x_n)\right| \leq C_\ell \sigma(\lambda,\xi')^{\frac{\ell}{2}-\frac{1}{2}-\frac{|\alpha'|}{2}}e^{-c\sigma(\lambda,\xi')^{\frac{1}{2}}(a-x_n)}$$

for j = 1, 2, and

(ii)
$$\left|\partial_{\xi'}^{\alpha'}\partial_{x_n}^{\ell}h_{\mu_1,\mu_2}(x_n)\right| \leq C_\ell \sigma(\lambda,\xi')^{\frac{\ell}{2}-\frac{3}{2}-\frac{|\alpha'|}{2}}e^{-c\sigma(\lambda,\xi')^{\frac{1}{2}}(a-x_n)}.$$

The inequalities (i) and (ii) also hold uniformly in $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$ for some $R_3 > 0$, where η_3 and θ_3 are the numbers given in Lemma 4.4.

Proof. The proof is similar to that of Lemma 4.6. We omit it.

We next estimate $\beta_j(\lambda, \xi', y_n)$, $B'(\lambda, \xi', y_n)$ and $\boldsymbol{b}_n(\lambda, \xi', y_n)$.

Lemma 4.8. Let $\alpha' \in \mathbb{Z}^{n-1}$ be a multi-index with $|\alpha'| \leq [\frac{n-1}{2}] + 1$. Then the following estimates hold uniformly in $\lambda \in \Sigma(\eta_4, \theta_4), \xi' \in \mathbb{R}^{n-1}$ and $y_n \in [0, a]$:

(i)
$$\left|\partial_{\xi'}^{\alpha'}\beta_0(\lambda,\xi',y_n)\right| \leq C_\ell \left(|\lambda|+1\right)^{-1} \sigma(\lambda,\xi')^{-\frac{|\alpha'|}{2}} e^{-c\sigma(\lambda,\xi')^{\frac{1}{2}}(a-y_n)},$$

(ii)
$$\left|\partial_{\xi'}^{\alpha'}\beta_1(\lambda,\xi',y_n)\right| \le C\sigma(\lambda,\xi')^{-\frac{|\alpha'|}{2}}e^{-c\sigma(\lambda,\xi')^{\frac{1}{2}}(a-y_n)},$$

(iii)
$$\left|\partial_{\xi'}^{\alpha'}B'(\lambda,\xi',y_n)\right| \le C\sigma(\lambda,\xi')^{-\frac{|\alpha'|}{2}}e^{-c\sigma(\lambda,\xi')^{\frac{1}{2}}(a-y_n)}$$

(iv)
$$\left|\partial_{\xi'}^{\alpha'} \boldsymbol{b}_n(\lambda,\xi',y_n)\right| \leq C\sigma(\lambda,\xi')^{-\frac{|\alpha'|}{2}} e^{-c\sigma(\lambda,\xi')^{\frac{1}{2}}(a-y_n)}.$$

The inequalities (i)–(iv) also hold uniformly in $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$ for some $R_3 > 0$, where η_3 and θ_3 are the numbers given in Lemma 4.4.

Proof. A direct calculation shows that

(4.29)
$$\beta_0(\lambda,\xi',y_n) = \frac{\gamma\lambda}{\nu_1\lambda+\gamma^2} \frac{1}{4D_1(\lambda,\xi')} \frac{\mu_1\mu_2 - |\xi'|^2}{\mu_1\mu_2} e^{-\mu_2(a-y_n)} + \frac{1}{4D_1(\lambda,\xi')} \widetilde{\beta}_0(\lambda,\xi',y_n),$$

where $\tilde{\beta}_0(\lambda, \xi', y_n)$ is written as the sum of terms of the form

$$e^{-\mu_k a - \mu_1 z_n - \mu_2 w_n} \times \left\{ \text{polynomial in } \mu_1, \mu_2, \frac{1}{\mu_1}, \frac{1}{\mu_2} \text{ and } \xi' \right\}, \ k = 1, 2,$$

with z_n and w_n being linear functions of y_n that satisfy $z_n \ge 0$, $w_n \ge 0$ and Re $(\mu_k a + \mu_1 z_n + \mu_2 w_n) \ge \text{Re } \mu_2(a - y_n)$ for $y_n \in [0, a]$. The desired inequality in (i) is obtained by applying Lemma 4.5 to (4.29). Similarly, by a direct computation, we see that $\beta_1(\lambda, \xi', y_n)$ and $\boldsymbol{b}_n(\lambda, \xi', y_n)$ can be written as

$$\beta_{1}(\lambda,\xi',y_{n}) = \frac{\mu_{1}\mu_{2}-|\xi'|^{2}}{4D_{1}(\lambda,\xi')} \left\{ \frac{\mu_{1}\mu_{2}-|\xi'|^{2}}{\mu_{1}\mu_{2}} e^{-\mu_{2}(a-y_{n})} - \left(e^{-\mu_{1}(a-y_{n})} - e^{-\mu_{2}(a-y_{n})}\right) \right\} + \widetilde{\beta}_{1}(\lambda,\xi',y_{n})$$

and

$$\boldsymbol{b}_{n}(\lambda,\xi',y_{n}) = -\frac{i\xi'}{4D_{1}(\lambda,\xi')} \frac{\mu_{1}\mu_{2} - |\xi'|^{2}}{\mu_{1}} \left(e^{-\mu_{1}(a-y_{n})} - e^{-\mu_{2}(a-y_{n})} \right)$$

+ $i\xi' \widetilde{\beta}_{n}(\lambda,\xi',y_{n}),$

where $\tilde{\beta}_j(\lambda, \xi', y_n)$, j = 1, n, are written as the sum of terms of the same form as that for $\tilde{\beta}_0(\lambda, \xi', y_n)$ with z_n and w_n satisfying $z_n \ge 0$, $w_n \ge 0$ and $\operatorname{Re}(\mu_k a + \mu_1 z_n + \mu_2 w_n) \ge \operatorname{Re} \mu_\ell(a - y_n)$ for $y_n \in [0, a]$, k = 1, 2 and $\ell = 1$ or 2. The desired inequalities in (ii) and (iv) now follow from Lemma 4.5.

We next consider inequality (iii). We write $B'(\lambda, \xi', y_n)$ as

$$B'(\lambda,\xi',y_n) = -\frac{\sinh\mu_1 y_n}{\sinh\mu_1 a}I_{n-1} - \beta(\lambda,\xi',y_n)P'_{1,0}$$

where

$$\beta(\lambda,\xi',y_n) = \frac{\sinh \mu_1 y_n}{\sinh \mu_1 a} - \beta_1(\lambda,\xi',y_n).$$

A direct computation shows that $\beta(\lambda, \xi', y_n)$ is written as

$$\beta(\lambda,\xi',y_n) = -\frac{|\xi'|^2}{4D_1(\lambda,\xi')} \frac{\mu_1\mu_2 - |\xi'|^2}{\mu_1\mu_2} \left(e^{-\mu_1(a-y_n)} - e^{-\mu_2(a-y_n)} \right) \\ + \frac{|\xi'|^2}{4D_1(\lambda,\xi')(1-e^{-2\mu_1a})} \widetilde{\beta}(\lambda,\xi',y_n),$$

where $\tilde{\beta}(\lambda, \xi', y_n)$ is written as the sum of the same form as that for $\tilde{\beta}_1(\lambda, \xi', y_n)$. Consequently we have

$$B'(\lambda,\xi',y_n) = \frac{1}{\mu_1} e^{-\mu_1(a-y_n)} I_{n-1} + \frac{\xi'^T \xi'}{4D_1(\lambda,\xi')} \frac{\mu_1 \mu_2 - |\xi'|^2}{\mu_1 \mu_2} \left(e^{-\mu_1(a-y_n)} - e^{-\mu_2(a-y_n)} \right) \\ + \frac{1}{\mu_1} \frac{e^{-\mu_1 a}}{1 - e^{-2\mu_1 a}} \left(e^{-\mu_1(2a-y_n)} - e^{-\mu_1 y_n} \right) I_{n-1} + \frac{\xi'^T \xi'}{4D_1(\lambda,\xi')(1 - e^{-2\mu_1 a})} \widetilde{\beta}(\lambda,\xi',y_n).$$

The desired inequality in (iii) now follows from Lemma 4.5.

Similarly to above, one can prove the desired inequalities for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \ge R_3$ with $|\lambda| \ge \delta$, where δ is any positive number.

Let us consider the case $|\lambda| \leq \delta$. In this case a direct computation shows that

$$\left|\partial_{\xi'}^{\alpha'}\widetilde{\beta}_j(\lambda,\xi',y_n)\right| \le C|\lambda|^2 e^{-c|\xi'|a}, \quad j=0,1,n,$$

for small $|\lambda|$ and large $|\xi'|$. Combining this with Lemma 4.5, we can obtain the desired inequalities for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \geq R_3$ with some large $R_3 > 0$. This completes the proof.

5. Proof of the main results

In this section we prove the main results stated in section 2.

To prove Theorem 2.1 we will apply the following Fourier multiplier theorem. **Lemma 5.1.** (Fourier multiplier theorem) Let $1 and let s be an integer satisfying <math>s \ge \left[\frac{k}{2}\right] + 1$. Suppose that $\Psi(\zeta) \in C^s(\mathbf{R}^k - \{0\}) \cap L^{\infty}(\mathbf{R}^k)$ and that there exists a positive constant C_0 such that

$$\left|\zeta\right|^{|\alpha|} \left|\partial_{\zeta}^{\alpha} \Psi(\zeta)\right| \le C_0$$

for all $\zeta \in \mathbf{R}^k - \{0\}$ and $|\alpha| \leq s$. Then the operator $\mathscr{F}_{\zeta}^{-1}[\Psi(\zeta)(\mathscr{F}f)(\zeta)]$ is extended to a bounded linear operator on $L^p(\mathbf{R}^k)$ and there holds the estimate

$$\|\mathscr{F}_{\zeta}^{-1}\left[\Psi(\zeta)(\mathscr{F}f)(\zeta)\right]\|_{L^{p}(\mathbf{R}^{k})} \leq CC_{0}\|f\|_{L^{p}(\mathbf{R}^{k})}$$

See, e.g., [4] for the proof.

We will also use the following lemma concerning integral operator.

Lemma 5.2. Let $\Phi(x_n, y_n)$ be a measurable function on $(0, a) \times (0, a)$.

(i) Suppose that there exists a positive constant M such that $\int_0^a |\Phi(x_n, y_n)| dy_n \le M$ for a.e. x_n and $\int_0^a |\Phi(x_n, y_n)| dx_n \le M$ for a.e. y_n . Let $1 \le p \le \infty$. Then it holds that

$$\left|\int_0^a \Phi(\cdot, y_n) f(y_n) \, dy_n\right|_p \le M |f|_p.$$

(ii) Suppose that there exists a positive constant M such that $(\int_0^a |\Phi(x_n, y_n)|^2 dy_n)^{\frac{1}{2}} \leq M$ for a.e. x_n . Then it holds that

$$\left|\int_0^a \Phi(\cdot, y_n) f(y_n) \, dy_n\right|_{\infty} \le M |f|_2.$$

The proof of the above lemma is well known. We omit it.

To obtain the estimates for derivatives of the resolvent we still need some consideration. We proceed as in [3].

Lemma 5.3. There holds

$$\mathscr{F}_{\xi'}^{-1}\left[\int_0^a \frac{1}{2\mu_j} e^{-\mu_j |x_n - y_n|} \widehat{f}(\xi', y_n) \, dy_n\right](x', x_n) = \mathscr{F}_{\xi}^{-1}\left[\frac{1}{\mu_j^2 + \xi_n^2} \mathscr{F}Ef(\xi)\right](x', x_n)$$

for j = 1, 2, where $\xi = (\xi', \xi_n)$ and $Ef(x', x_n)$ is the zero-extension of $f(x', x_n)$ to \mathbf{R}^n .

Proof. Since $\mathscr{F}_{x_n}\left[\frac{1}{2\mu_j}e^{-\mu_j|x_n|}\right] = \frac{1}{\mu_j^2 + \xi_n^2}$, we obtain the desired relation. This completes the proof.

We will also use the following lemma ([3, Lemma 2.6]).

Lemma 5.4. ([3, Lemma 2.6]) Let 1 and define the operator T by

$$Tf(x_n) = \int_0^a \frac{f(y_n)}{x_n + y_n} \, dy_n$$

for $x_n \in (0, a)$ and $f \in L^p(0, a)$. Then there exists a positive constant C = C(p) such that

$$|Tf|_p \le C|f|_p.$$

This lemma follows from the fact that Tf can be written in a form of the Hilbert transformation on **R**. See [3, Lemma 2.6].

We are now in a position to Prove Theorem 2.1.

Proof of Theorem 2.1. Let η be any positive number. We will prove Theorem 2.1 with θ satisfying $\frac{\pi}{2} < \theta < \min \{\theta_1, \theta_3, \theta_4\}$, where θ_j (j = 1, 3, 4)are the numbers given in Lemmas 4.3 and 4.4. We note that $\Sigma(\eta, \theta) \subset$ $\Sigma(\eta, \theta_1) \cap \Sigma(-\eta_3, \theta_3)$ and that $\overline{\Sigma(\eta, \theta) - \Sigma(\eta_4, \theta_4)}$ is a compact set. Here η_3 and η_4 are the numbers given in Lemma 4.4.

We see from Theorem 3.8 that there exists a solution u of problem (1.1)–(1.2) which takes the form $u = R(\lambda)f$, where $\hat{R}(\lambda,\xi') = (\lambda + \hat{L}_{\xi'})^{-1} = \hat{G}(\lambda,\xi') + \hat{K}(\lambda,\xi')$ with $\hat{G}(\lambda,\xi')$ and $\hat{K}(\lambda,\xi')$ being the integral operators given in Theorem 3.8. We first prove that the estimates in Theorem 2.1 hold for $u = R(\lambda)f$. We then prove the uniqueness of solutions of problem (1.1)–(1.2).

Let us estimate $u = R(\lambda)f$. We first consider the case $\lambda \in \Sigma(\eta_4, \theta_4)$. We note that one can see from the form of $\widehat{R}(\lambda, \xi')$ that k-th order derivatives of $Q_0R(\lambda)f$ are estimated as (k + 1)-th order ones of $\widetilde{Q}R(\lambda)f$. So we here give the proof of the estimates for $\widetilde{Q}R(\lambda)f$ only.

Hereafter we set $G(\lambda) = \mathscr{F}_{\xi'}^{-1}[\widetilde{G}(\lambda,\xi')]$ and $K(\lambda) = \mathscr{F}_{\xi'}^{-1}[\widehat{K}(\lambda,\xi')]$. We also write $G(\lambda) = G^{(1)}(\lambda) + \cdots + G^{(4)}(\lambda)$, where $G^{(\ell)}(\lambda)$ is the matrix with $g_{\mu_j}^D$ and $g_{\mu_j}^N$ in $G(\lambda)$ replaced by $(-1)^{\ell+1}g_{\mu_j}^{(\ell)}$ and $g_{\mu_j}^{(\ell)}$, respectively. Here $g_{\mu_j}^{(\ell)}$ are the functions defined in (4.28).

Let us consider $\widetilde{Q}R(\lambda)\widetilde{Q}f$. We first estimate $\partial_x^k\widetilde{Q}G(\lambda)\widetilde{Q}f$ for $k \leq 1$. Since $\partial_{x_n}^2\left(g_{\mu_1,\mu_2}^M\widetilde{Q}f\right) = \left(\partial_{x_n}^2g_{\mu_1}^M - \partial_{x_n}^2g_{\mu_2}^M\right)\left[\widetilde{Q}\widehat{f}\right]$, we see from Lemma 4.6 that, for

 $\begin{aligned} |\beta'| + \ell &\leq 1 \text{ and } j = 1, 2, \\ \left| \partial_{\xi'}^{\alpha'} \left[(i\xi')^{\beta'} \partial_{x_n}^{\ell} \widetilde{Q} \widehat{G}^{(j)}(\lambda, \xi' x_n, y_n) \widetilde{Q} \right] \right| \\ &\leq C \sigma(\lambda, \xi')^{\frac{|\beta'| + \ell}{2} - \frac{1}{2} - \frac{|\alpha'|}{2}} \left\{ E^{(1)}(x_n, y_n) + E^{(2)}(x_n, y_n) \right\} \\ &\leq C(|\lambda| + 1)^{\frac{|\beta'| + \ell}{2} - \frac{1}{2}} \mathcal{E}(\lambda, x_n, y_n) |\xi'|^{-|\alpha'|}, \end{aligned}$

where

$$\mathcal{E}(\lambda, x_n, y_n) = e^{-c(|\lambda|+1)^{\frac{1}{2}}|x_n - y_n|} + e^{-c(|\lambda|+1)^{\frac{1}{2}}(x_n + y_n)} + e^{-c(|\lambda|+1)^{\frac{1}{2}}\{(a-x_n) + (a-y_n)\}}.$$

Since

$$\sup_{0 \le x_n \le a} \int_0^a \mathcal{E}(\lambda, x_n, y_n) \, dy_n + \sup_{0 \le y_n \le a} \int_0^a \mathcal{E}(\lambda, x_n, y_n) \, dx_n \le C(|\lambda| + 1)^{-\frac{1}{2}},$$

we see from Lemmas 5.1 and 5.2 that

$$\|\partial_x^k \tilde{Q} G^{(j)}(\lambda) \tilde{Q} f\|_p \le \frac{C}{(|\lambda|+1)^{1-\frac{k}{2}}} \|\tilde{Q} f\|_p, \quad j=1,2,$$

for $k \leq 1$. Similarly, one can estimate $\partial_x^k \tilde{Q} G^{(j)}(\lambda) \tilde{Q} f$ (j = 3, 4) and $\partial_x^k \tilde{Q} K(\lambda) \tilde{Q} f$ for $k \leq 1$.

We next consider $\partial_x^2 \tilde{Q} R(\lambda) \tilde{Q} f$. We first estimate $\partial_x^2 \tilde{Q} G^{(1)}(\lambda) \tilde{Q} f$. By Lemma 5.3, we have

$$\mathscr{F}_{\xi'}^{-1}\left[g_{\mu_j}^{(1)}\tilde{Q}\hat{f}\right](x',x_n) = \mathscr{F}_{\xi}^{-1}\left[\frac{1}{\mu_j^2 + \xi_n^2}\mathscr{F}_x[\tilde{Q}Ef]\right](x',x_n), \quad j = 1,2,$$

and, in particular,

$$\mathscr{F}_{\xi'}^{-1}\left[g_{\mu_1,\mu_2}^{(1)}\widetilde{Q}\widehat{f}\right](x',x_n) = \mathscr{F}_{\xi}^{-1}\left[M(\lambda,\xi)\mathscr{F}_x[\widetilde{Q}Ef]\right](x',x_n),$$

where Ef denotes the zero extension of f to \mathbf{R}^n and

$$M(\lambda,\xi) = \frac{1}{\mu_1^2 + \xi_n^2} - \frac{1}{\mu_2^2 + \xi_n^2}$$

with $\mu_j = \mu_j(\lambda, \xi'), \ j = 1, 2$. Since $\mu_1^2 + \xi_n^2 = \frac{\lambda + \nu |\xi|^2}{\nu}, \ \mu_2^2 + \xi_n^2 = \frac{\lambda^2 + \nu_1 |\xi|^2 \lambda + \gamma^2 |\xi|^2}{\nu_1 \lambda + \gamma^2}$ with $|\xi|^2 = |\xi'|^2 + \xi_n^2$, and $\nu_1 = \nu + \tilde{\nu}$, we have

$$M(\lambda,\xi) = -\frac{\lambda(\widetilde{\nu}\lambda + \gamma^2)}{(\lambda + \nu|\xi|^2)(\lambda^2 + \nu_1|\xi|^2\lambda + \gamma^2|\xi|^2)}.$$

Therefore, in view of the observation in Remarks 3.2 and 3.5, we obtain $\left|\partial_{\xi}^{\alpha}\left(\frac{\xi^{\beta}}{\mu_{j}^{2}+\xi_{n}^{2}}\right)\right| \leq C|\xi|^{-|\alpha|}$ for $|\beta| = 2$ and $\left|\partial_{\xi}^{\alpha}\left(\frac{\xi^{\beta}}{\lambda}M(\lambda,\xi)\right)\right| \leq C|\xi|^{-|\alpha|}$ for $|\beta| = 4$ uniformly in $\lambda \in \Sigma(\eta_{4}, \theta_{4})$. It then follows from Lemma 5.1 that

 $\|\partial_x^2 \widetilde{Q} G^{(1)}(\lambda) \widetilde{Q} f\|_p \le C \|\widetilde{Q} f\|_p,$

We next consider $\partial_x^2 \tilde{Q} G^{(2)}(\lambda) \tilde{Q} f$. For $|\beta'| + \ell = 2$, we see from Lemma 4.6 that

$$\begin{aligned} \left| \partial_{\xi'}^{\alpha} \left[(i\xi')^{\beta} \partial_{x_n}^{\epsilon} Q G^{(2)}(\lambda,\xi',x_n,y_n) Q \right] \right| \\ &\leq C \sigma(\lambda,\xi')^{\frac{1}{2} - \frac{|\alpha'|}{2}} E^{(2)}(x_n,y_n) \\ &\leq C \left\{ \frac{1}{x_n + y_n} + \frac{1}{(a - x_n) + (a - y_n)} \right\} |\xi'|^{-|\alpha'|}. \end{aligned}$$

It then follows from Lemma 5.1 that

$$\left\| \partial_x^2 \tilde{Q} G^{(2)}(\lambda) \tilde{Q} f(\cdot, y_n) \right\|_{L^p(\mathbf{R}^{n-1})}$$

 $\leq C \int_0^a \left\{ \frac{1}{x_n + y_n} + \frac{1}{(a - x_n) + (a - y_n)} \right\} \left\| \tilde{Q} f(\cdot, y_n) \right\|_{L^p(\mathbf{R}^{n-1})} dy_n$

and hence, by Lemma 5.4, we obtain

$$\left\|\partial_x^2 \widetilde{Q} G^{(2)}(\lambda) \widetilde{Q} f\right\|_p \le C \left\|\widetilde{Q} f\right\|_p.$$

In the same way one can also obtain the desired estimates for $\partial_x^2 \tilde{Q} G^{(j)}(\lambda) \tilde{Q} f$ (j = 3, 4) and $\partial_x^2 \tilde{Q} K(\lambda) \tilde{Q} f$.

We next consider $\tilde{Q}R(\lambda)Q_0f$. Similarly to above, one can prove $\left\|\partial_x^k \tilde{Q}R(\lambda)Q_0f\right\|_p \leq \frac{C}{|\lambda|+1} \|Q_0f\|_p$ for $k \leq 1$. As for $\partial_x^2 \tilde{Q}R(\lambda)Q_0f$, we first note that $\partial_{x_n}g_{\mu_2}^NQ_0\hat{f} = g_{\mu_2}^D[\partial_{x_n}Q_0\hat{f}]$, which is obtained by integration by parts. It then follows that

$$\tilde{Q}\hat{G}(\lambda,\xi')Q_0\hat{f} = -\frac{\gamma}{\nu_1\lambda + \gamma^2} \begin{pmatrix} 0\\ g^N_{\mu_2}[\nabla\widehat{Q_0}f]\\ g^D_{\mu_2}[\partial_{x_n}\widehat{Q_0}f] \end{pmatrix},$$

where $\nabla' = {}^{T}(\partial_{x_1}, \cdots, \partial_{x_{n-1}})$. Therefore, similarly to above, we obtain

$$\|\partial_x^2 \widetilde{Q} G(\lambda) Q_0 f\|_p \le \frac{C}{|\lambda|+1} \|Q_0 f\|_{W^{1,p}}.$$

As for $\partial_x^2 \tilde{Q} K(\lambda) Q_0 f$, we write

$$\widetilde{Q}\widehat{H}(\lambda,\xi')Q_0\widehat{f} = \begin{pmatrix} 0\\ \left(\frac{1}{\nu}h_{\mu_1}\beta_0 + \frac{|\xi'|^2}{\lambda}h_{\mu_1,\mu_2}\right)[\nabla\widehat{Q}_0f]\\ \frac{i\xi'}{\lambda}\partial_{x_n}h_{\mu_1,\mu_2}[\nabla\widehat{Q}_0f] \end{pmatrix}.$$

The desired estimate for $\partial_x^2 \tilde{Q} K(\lambda) Q_0 f$ now follows similarly to above.

As mentioned before, one can prove the estimates for $\partial_x^k Q_0 R(\lambda) f$ with $k \leq 1$ similarly to the estimates for $\partial_x^{k+1} \tilde{Q} R(\lambda) f$. We thus omit the details.

Let us prove the last inequality in Theorem 2.1. We assume that $\tilde{Q}f|_{x_n=0,a} = 0$. It suffices to prove

$$\|\partial_{x_n} \widetilde{Q}R(\lambda)\widetilde{Q}f\|_p \le \frac{C}{|\lambda|+1} \|\widetilde{Q}f\|_{W^{1,p}}.$$

Since $\tilde{Q}f\Big|_{x_n=0,a} = 0$, by integration by parts, we also have

$$\partial_{x_n} g^D_{\mu_j} \widetilde{Q} \widehat{f} = g^N_{\mu_j} \left[\partial_{x_n} \widetilde{Q} \widehat{f} \right], \quad j = 1, 2.$$

Therefore, we see from Lemmas 4.6, 5.1 and 5.2 that

$$\|\partial_{x_n} \widetilde{Q}G(\lambda)\widetilde{Q}f\|_p \le \frac{C}{|\lambda|+1} \|\partial_{x_n} \widetilde{Q}f\|_p.$$

As for $\tilde{Q}K(\lambda)\tilde{Q}f$, we note that an application ∂_{x_n} to $h_{\mu_j}(x_n)$ yields one of the factors μ_j , j = 1, 2. But, since $B'(\lambda, \xi', y_n)$ and $\boldsymbol{b}_n(\lambda, \xi', y_n)$ are written as linear combinations of $e^{\pm \mu_j y_n}$ (j = 1, 2), we see that if $\tilde{Q}f|_{x_n=0,a} = 0$, then

$$\int_0^a e^{\pm\mu_j y_n} \widetilde{Q}\widehat{f}(y_n) \, dy_n = \mp \frac{1}{\mu_j} \int_0^a e^{\pm\mu_j y_n} \partial_{y_n} \widetilde{Q}\widehat{f}(y_n) \, dy_n, \quad j = 1, 2.$$

Therefore, we can gain one of the factors μ_j^{-1} , j = 1, 2. It then follows that

$$\|\partial_{x_n} \widetilde{Q} K(\lambda) \widetilde{Q} f\|_p \le \frac{C}{|\lambda|+1} \|\partial_{x_n} \widetilde{Q} f\|_p.$$

We have thus obtained the desired estimates when $\lambda \in \Sigma(\eta_4, \theta_4)$.

We next consider the case $\lambda \in \Sigma(\eta, \theta) - \Sigma(\eta_4, \theta_4)$. We fix a positive number R_3 in such a way that Lemmas 4.5–4.8 hold for $\lambda \in \Sigma(-\eta_3, \theta_3)$ and $|\xi'| \geq R_3$.

We decompose $R(\lambda)f$ in the following way. Let $\tilde{\chi}(\xi')$ be a C^{∞} function on \mathbf{R}^{n-1} satisfying $0 \leq \tilde{\chi} \leq 1$ on \mathbf{R}^{n-1} , $\tilde{\chi}(\xi') = 0$ for $|\xi'| \leq R_3$ and $\tilde{\chi}(\xi') = 1$ for $|\xi'| \geq 2R_3$. We write $R(\lambda)f$ as

$$R(\lambda)f = \tilde{R}^{(0)}(\lambda)f + \tilde{R}^{(\infty)}(\lambda)f,$$

where

$$\widetilde{R}^{(0)}(\lambda)f = \mathscr{F}_{\xi'}^{-1}\left[(1 - \widetilde{\chi}(\xi'))\widehat{R}(\lambda,\xi')\widehat{f}\right]$$

and

$$\widetilde{R}^{(\infty)}(\lambda)f = \mathscr{F}_{\xi'}^{-1}\left[\widetilde{\chi}(\xi')\widehat{R}(\lambda,\xi')\widehat{f}\right].$$

Similarly to the case of $\lambda \in \Sigma(\eta_4, \theta_4)$, we can obtain the desired estimates for $\tilde{R}^{(\infty)}(\lambda)f$ since $\Sigma(\eta, \theta) \subset \Sigma(-\eta_3, \theta_3)$. The desired estimates also hold for $\tilde{R}^{(0)}(\lambda)f$. In fact, by Lemmas 3.4 and 4.3, we see that $D(\lambda, \xi') \neq 0$ on the compact set $\overline{\Sigma(\eta, \theta)} - \overline{\Sigma(\eta_4, \theta_4)} \times \{\xi'; |\xi'| \leq 2R_3\}$. Therefore, $\hat{R}(\lambda, \xi', x_n, y_n)$ is analytic, and so we can obtain the desired estimates for $\tilde{R}^{(0)}(\lambda)f$. We thus conclude that $u = R(\lambda)f$ satisfies the estimates in Theorem 2.1.

It remains to prove the uniqueness of solutions of problem (1.1)-(1.2). To prove the uniqueness, we consider the adjoint problem

(5.1)
$$(\overline{\lambda} + L^*)w = g, \quad \widetilde{Q}w\Big|_{x_n = 0,a} = 0,$$

where

$$L = \begin{pmatrix} 0 & -\gamma \operatorname{div} \\ \\ -\gamma \nabla & -\nu \Delta I_n - \tilde{\nu} \nabla \operatorname{div} \end{pmatrix}.$$

Similarly to the case of problem (1.1)–(1.2), we consider the Fourier transform of (5.1) in $x' \in \mathbf{R}^{n-1}$:

$$(\overline{\lambda} + \widehat{L}^*_{\xi'})\widehat{w} = \widehat{g}, \quad \widetilde{Q}\widehat{w}\Big|_{x_n = 0, a} = 0.$$

It is easily verified that $(\overline{\lambda} + \widehat{L}_{\xi'}^*)^{-1}$ has an integral representation $\widehat{R}^*(\overline{\lambda}, \xi')$ corresponding to Theorem 3.8, which is written in terms of the components of $\widehat{R}(\overline{\lambda}, \xi')$. Therefore, $R^*(\overline{\lambda})g = \mathscr{F}_{\xi'}^{-1} \left[\widehat{R}^*(\overline{\lambda}, \xi')\widehat{g}\right]$, which is a solution of (5.1), has the same estimates as those for $R(\lambda)f$. We thus conclude that for any $\lambda \in \Sigma(\eta, \theta)$ and $g \in W^{1,p} \times L^p$ problem (5.1) has a solution $w \in W^{1,p} \times \left(W^{2,p} \cap W_0^{1,p}\right)$.

Assume now that $u \in W^{1,p} \times (W^{2,p} \cap W_0^{1,p})$ and $(\lambda + L)u = 0$. Let $g \in C_0^{\infty}(\overline{\Omega}) \times C_0^{\infty}(\Omega)$ and let w be a solution of (5.1) satisfying $w \in W^{1,p'} \times (W^{2,p'} \cap W_0^{1,p'})$ with 1/p' = 1 - 1/p. It then follows that

$$0 = ((\lambda + L)u, w) = (u, (\overline{\lambda} + L^*)w) = (u, g).$$

This implies that u = 0, and the uniqueness of problem (1.1)–(1.2) holds for $\lambda \in \Sigma(\eta, \theta)$. This completes the proof.

We next give a proof of Theorem 2.2.

Proof of Theorem 2.2. We here prove the second inequality in Theorem 2.2 for $\partial_{x_n} Q_n G(\lambda) f$ only, since the other cases can be proved similarly by applying Lemmas 4.6–4.8 and 5.2.

We first consider the case $\lambda \in \Sigma(\eta_4, \theta_4)$. Let $E^{(j)}(x_n, y_n)$, j = 1, 2, be the functions defined in Lemma 4.6 and set $E(x_n, y_n) = E^{(1)}(x_n, y_n) + E^{(2)}(x_n, y_n)$. Then we have

$$\sup_{0 \le x_n \le a} \left(\int_0^a |E(x_n, y_n)|^2 \, dy_n \right)^{\frac{1}{2}} \le C\sigma(\lambda, \xi')^{-\frac{1}{4}}.$$

Since $\partial_{x_n}^2 \left(g_{\mu_2}^N Q_0 \hat{f} \right) = \left(\partial_{x_n}^2 g_{\mu_2}^N \right) \left[Q_0 \hat{f} \right] - Q_0 \hat{f}$, we see from Lemmas 4.6 and 5.2 that

$$\begin{aligned} \|\partial_{x_n} Q_n G(\lambda) Q_0 f\|_{\infty} &\leq \frac{C}{|\lambda|+1} \left\{ \|Q_0 f\|_{\infty} + \left\| \left(\partial_{x_n}^2 g_{\mu_2}^N \right) \left[Q_0 \hat{f} \right] \right\|_{L^1_{\xi'} L^\infty_{x_n}} \right\} \\ &\leq \frac{C}{|\lambda|+1} \left\{ \|Q_0 f\|_{\infty} + \left\| (|\lambda|+1+|\xi'|^2)^{\frac{1}{4}} \left| Q_0 \hat{f}(\xi') \right|_2 \right\|_{L^1_{\xi'}} \right\} \\ &\leq C \left\{ \frac{1}{|\lambda|+1} \|Q_0 f\|_{\infty} + \frac{1}{(|\lambda|+1)^{\frac{3}{4}}} \|Q_0 f\|_{H^{[\frac{n}{2}]+1}} \right\} \\ &\leq \frac{C}{(|\lambda|+1)^{\frac{3}{4}}} \|Q_0 f\|_{H^{[\frac{n}{2}]+1}} . \end{aligned}$$

We next consider $\partial_{x_n} Q_n G(\lambda) Q' f$. Similarly to above, we see from Lemmas 4.6 and 5.2 that, for $0 < \varepsilon < 1$,

$$\begin{aligned} \|\partial_{x_n} Q_n G(\lambda) Q' f\|_{\infty} &\leq C \left\| \partial_{x_n} Q_n \widehat{G}(\lambda, \xi') Q' \widehat{f} \right\|_{L^1_{\xi'} L^\infty_{x_n}} \\ &\leq C \left\| (|\lambda| + 1 + |\xi'|^2)^{-\frac{3}{4}} |\xi'| \left| Q' \widehat{f}(\xi') \right|_2 \right\|_{L^1_{\xi'}} \\ &\leq C \left\| (|\lambda| + 1)^{-\frac{\varepsilon}{4}} (|\lambda| + 1 + |\xi'|^2)^{-\frac{1}{4}(1-\varepsilon)} \left| Q' \widehat{f}(\xi') \right|_2 \right\|_{L^1_{\xi'}} \\ &\leq \frac{C}{(|\lambda|+1)^{\frac{\varepsilon}{4}}} \left\| Q' \widehat{f} \right\|_{H^s} \end{aligned}$$

with $s > \frac{n}{2} - \varepsilon$. Therefore, we can take $s = \left[\frac{n}{2}\right]$. Similarly, we can obtain

$$\|\partial_{x_n} Q_n G(\lambda) Q_n f\|_{\infty} \le \frac{C}{(|\lambda|+1)^{\frac{\varepsilon}{4}}} \|f'(\xi')\|_{H^{[\frac{n}{2}]}}$$

We next consider the case $\lambda \in \Sigma(\eta, \theta) - \Sigma(\eta_4, \theta_4)$. We decompose $(\lambda + L)^{-1}f$ into $(\lambda + L)^{-1}f = \tilde{R}^{(0)}(\lambda)f + \tilde{R}^{(\infty)}(\lambda)f$ as in the proof of Theorem 2.1. It then follows that $\tilde{R}^{(\infty)}(\lambda)f$ can be estimated as above. One can also see that $\tilde{R}^{(0)}(\lambda)f$ has the desired estimates since $D(\lambda, \xi') \neq 0$ on the compact set $\overline{\Sigma(\eta, \theta)} - \overline{\Sigma(\eta_4, \theta_4)} \times \{\xi'; |\xi'| \leq 2R_3\}$. This completes the proof.

We next prove the L^p estimates of the resolvent for $p = 1, \infty$. For this purpose we prepare the following lemma.

Lemma 5.5. Let $\ell = 0, 1$ and let $\widehat{\Phi}(\xi', x_n)$ be a function satisfying

$$|\partial_{\xi'}^{\alpha'}\widehat{\Phi}(\xi',\cdot)|_1 \le C\sigma(\lambda,\xi')^{\frac{\ell}{2}-1-\frac{|\alpha'|}{2}}$$

for all α' with $|\alpha'| \leq n$. Then $\Phi(x', x_n) = (\mathscr{F}_{\xi'}^{-1}\widehat{\Phi})(x', x_n)$ satisfies

$$\|\Phi\|_1 \le C(|\lambda|+1)^{\frac{\ell}{2}-1}.$$

Proof. The proof is based on the Riemann-Lebesgue lemma as in the estimates of solutions to the Cauchy problem given in [8].

Since $2 - \ell + n > n - 1$, by integration by parts, we see that

$$\begin{aligned} |\Phi(x',\cdot)|_{1} &= \left| (2\pi)^{n-1} \int_{\mathbf{R}^{n-1}} \widehat{\Phi}(\xi',x_{n}) \left(\sum_{j=1}^{n-1} \frac{x_{j}}{i|x'|^{2}} \partial_{\xi_{j}} \right)^{n} e^{ix'\cdot\xi'} d\xi' \right|_{L^{1}_{x_{n}}} \\ &\leq C|x'|^{-n} \int_{\mathbf{R}^{n-1}} \sum_{|\alpha'|=n} \left| \partial_{\xi_{j}}^{\alpha'} \widehat{\Phi}(\xi',x_{n}) \right|_{L^{1}_{x_{n}}} d\xi' \\ &\leq C|x'|^{-n} \int_{\mathbf{R}^{n-1}} \left(|\lambda| + 1 + |\xi'|^{2} \right)^{\frac{\ell}{2} - 1 - \frac{n}{2}} d\xi' \\ &\leq C|x'|^{-n} (|\lambda| + 1)^{-\frac{3}{2} + \frac{\ell}{2}}. \end{aligned}$$

This implies that

$$\begin{aligned} \||\Phi(x',\cdot)|_1\|_{L^1(|x'| \ge (|\lambda|+1)^{-\frac{1}{2}})} &\leq C(|\lambda|+1)^{-\frac{3}{2}+\frac{\ell}{2}} \||x'|^{-n}\|_{L^1(|x'| \ge (|\lambda|+1)^{-\frac{1}{2}})} \\ &\leq C(|\lambda|+1)^{-1+\frac{\ell}{2}}. \end{aligned}$$

We next show that

$$\||\Phi(x',\cdot)|_1\|_{L^1(|x'| \le (|\lambda|+1)^{-\frac{1}{2}})} \le C(|\lambda|+1)^{-1+\frac{\ell}{2}}.$$

In case $\ell = 0$, similarly to above, we have

$$\begin{split} |\Phi(x',\cdot)|_{1} &= \left| (2\pi)^{n-1} \int_{\mathbf{R}^{n-1}} \widehat{\Phi}(\xi',x_{n}) \left(\sum_{j=1}^{n-1} \frac{x_{j}}{i|x'|^{2}} \partial_{\xi_{j}} \right)^{n-2} e^{ix'\cdot\xi'} d\xi' \right|_{L^{1}_{x_{n}}} \\ &\leq C|x'|^{-(n-2)} \int_{\mathbf{R}^{n-1}} \sum_{|\alpha'|=n-2} \left| \partial_{\xi_{j}}^{\alpha'} \widehat{\Phi}(\xi',x_{n}) \right|_{L^{1}_{x_{n}}} d\xi' \\ &\leq C|x'|^{-(n-2)} \int_{\mathbf{R}^{n-1}} \left(|\lambda|+1+|\xi'|^{2} \right)^{\frac{\ell}{2}-1-\frac{n-2}{2}} d\xi' \\ &\leq C|x'|^{-(n-2)} (|\lambda|+1)^{-\frac{1}{2}}. \end{split}$$

This implies

$$\begin{aligned} \||\Phi(x',\cdot)|_1\|_{L^1(|x'| \le (|\lambda|+1)^{-\frac{1}{2}})} &\le C(|\lambda|+1)^{-\frac{1}{2}} \left\| |x'|^{-(n-2)} \right\|_{L^1(|x'| \le (|\lambda|+1)^{-\frac{1}{2}})} \\ &\le C(|\lambda|+1)^{-1}. \end{aligned}$$

The proof of the lemma for $\ell = 0$ is thus complete.

Let us consider the case $\ell = 1$. We first observe that

(5.2)
$$\int_{\mathbf{R}^{n-1}} \partial_{\xi'}^{\alpha'} \widehat{\Phi}(\xi', x_n) e^{ix' \cdot \xi'} d\xi' = -\int_{\mathbf{R}^{n-1}} \partial_{\xi'}^{\alpha'} \widehat{\Phi}(\xi', x_n) e^{ix' \cdot (\xi' + \frac{x'}{|x'|^2} \pi)} d\xi' = -\int_{\mathbf{R}^{n-1}} \partial_{\xi'}^{\alpha'} \widehat{\Phi}(\xi' - z', x_n) e^{ix' \cdot \xi'} d\xi'.$$

Here and in what follows we write $z' = \frac{x'}{|x'|^2}\pi$. From (5.2) we see that

(5.3)
$$\leq C|x'|^{-(n-2)} \int_{\mathbf{R}^{n-1}} \sum_{|\alpha'|=n-2} \left| \partial_{\xi'}^{\alpha'} \left(\widehat{\Phi}(\xi'-z',x_n) - \widehat{\Phi}(\xi',x_n) \right) \right|_{L^1_{x_n}} d\xi'.$$

By assumption, we have

 $|\Phi(x')|_1$

(5.4)
$$\left| \partial_{\xi'}^{\alpha'} \left(\widehat{\Phi}(\xi' - z', x_n) - \widehat{\Phi}(\xi', x_n) \right) \right|_{L^1_{x_n}} \le C \Phi^{(1)}(\xi', x'),$$

where

$$\Phi^{(1)}(\xi', x') = \left(|\lambda| + 1 + |\xi'|^2\right)^{-\frac{n-1}{2}} + \left(|\lambda| + 1 + |\xi' - z'|^2\right)^{-\frac{n-1}{2}}.$$

We also have

(5.5)

$$\begin{aligned} \left| \partial_{\xi'}^{\alpha'} \left(\widehat{\Phi}(\xi' - z', x_n) - \widehat{\Phi}(\xi', x_n) \right) \right|_{L^1_{x_n}} \\ &\leq C |z'| \sum_{\substack{|\beta'|=n-1 \\ \beta' \mid = n-1}} \int_0^1 \left| \partial_{\xi'}^{\beta'} \widehat{\Phi}(\xi' - \theta z', x_n) \right|_{L^1_{x_n}} d\theta \\ &\leq C |x'|^{-1} \Phi^{(2)}(\xi', x'), \end{aligned}$$

where

$$\Phi^{(2)}(\xi', x') = \int_0^1 \left(|\lambda| + 1 + |\xi' - \theta z'|^2 \right)^{-\frac{n}{2}} d\theta.$$

Let τ be a number satisfying $1 - \frac{1}{n} < \tau < 1$. It then follows from (5.4) and (5.5) that, for $|\alpha'| = n - 2$,

(5.6)
$$\begin{aligned} \int_{\mathbf{R}^{n-1}} \left| \partial_{\xi'}^{\alpha'} \left(\widehat{\Phi}(\xi' - z', x_n) - \widehat{\Phi}(\xi', x_n) \right) \right|_{L^{1}_{x_n}} d\xi' \\ &\leq C |x'|^{-(1-\tau)} \int_{\mathbf{R}^{n-1}} \widehat{\Phi}^{(1)}(\xi', x')^{\tau} \widehat{\Phi}^{(2)}(\xi', x')^{1-\tau} d\xi' \\ &= C |x'|^{-(1-\tau)} \int_{\Omega_1} + \int_{\Omega_2} + \int_{\Omega_3} \widehat{\Phi}^{(1)}(\xi', x')^{\tau} \widehat{\Phi}^{(2)}(\xi', x')^{1-\tau} d\xi', \end{aligned}$$

where

$$\Omega_{1} = \{\xi'; |\xi'| \ge 2|z'|\},
\Omega_{2} = \{\xi'; |\xi'| \le 2|z'|\} \cap \{\xi'; |\xi' - z'| \ge 2|z'|\},
\Omega_{3} = \{\xi'; |\xi'| \le 2|z'|\} \cap \{\xi'; |\xi' - z'| \le 2|z'|\}.$$

On Ω_1 we have $|\xi' - z'| \ge \frac{1}{2}|\xi'|$ and $|\xi' - \theta z'| \ge \frac{1}{2}|\xi'|$ for any $\theta \in [0, 1]$. It follows that (5.7)

$$\int_{\Omega_1} \widehat{\Phi}^{(1)}(\xi', x')^{\tau} \widehat{\Phi}^{(2)}(\xi', x')^{1-\tau} d\xi' \leq C \int_{\mathbf{R}^{n-1}} \left(|\lambda| + 1 + |\xi'|^2 \right)^{-\frac{n}{2} + \frac{\tau}{2}} d\xi' \\ \leq C(|\lambda| + 1)^{-\frac{1}{2} + \frac{\tau}{2}}.$$

On Ω_2 we have $|\xi' - z'| \ge |\xi'|$ and $|\xi' - \theta z'| \ge |\xi' - z'| - (1 - \theta)|z'| \ge \frac{1}{2}|\xi'|$ for any $\theta \in [0, 1]$. Therefore, as above, we obtain

(5.8)
$$\int_{\Omega_2} \widehat{\varPhi}^{(1)}(\xi', x')^{\tau} \widehat{\varPhi}^{(2)}(\xi', x')^{1-\tau} d\xi' \le C(|\lambda|+1)^{-\frac{1}{2}+\frac{\tau}{2}}.$$

As for the integral on Ω_3 , we have

(5.9)

$$\int_{\Omega_{3}} \widehat{\Phi}^{(1)}(\xi', x')^{\tau} \widehat{\Phi}^{(2)}(\xi', x')^{1-\tau} d\xi' \\
\leq C \int_{\{|\xi'| \le 2|z'|\}} |\xi'|^{-(n-1)\tau} (|\lambda|+1)^{-\frac{n}{2}(1-\tau)} d\xi' \\
+ C \int_{\{|\xi'-z'| \le 2|z'|\}} |\xi' - z'|^{-(n-1)\tau} (|\lambda|+1)^{-\frac{n}{2}(1-\tau)} d\xi' \\
\leq C(|\lambda|+1)^{-\frac{n}{2}(1-\tau)} |x'|^{-(1-\tau)(n-1)}.$$

Consequently we see from (5.6)–(5.9) that

$$\int_{\mathbf{R}^{n-1}} \left| \partial_{\xi'}^{\alpha'} \left(\widehat{\Phi}(\xi' - z', x_n) - \widehat{\Phi}(\xi', x_n) \right) \right|_{L^1_{x_n}} d\xi'$$

$$\leq C \left\{ (|\lambda| + 1)^{-\frac{1}{2} + \frac{\tau}{2}} |x'|^{-(1-\tau)} + (|\lambda| + 1)^{-\frac{n}{2}(1-\tau)} |x'|^{-(1-\tau)n} \right\}.$$

This, together with (5.3), implies that

$$\begin{aligned} |\Phi(x')|_1 &\leq C \left\{ (|\lambda|+1)^{-\frac{1}{2}+\frac{\tau}{2}} |x'|^{-(n-1)+\tau} \\ &+ (|\lambda|+1)^{-\frac{n}{2}(1-\tau)} |x'|^{-(n-2)-(1-\tau)n} \right\} \end{aligned}$$

Since $1 - \frac{1}{n} < \tau$, integrating this over $\{x'; |x'| \le (|\lambda| + 1)^{-\frac{1}{2}}\}$, we obtain

$$\||\Phi(x',\cdot)|_1\|_{L^1(|x'| \le (|\lambda|+1)^{-\frac{1}{2}})} \le C(|\lambda|+1)^{-\frac{1}{2}}.$$

This completes the proof.

We now prove the L^p estimates of the resolvent for $p = 1, \infty$.

Proof of Theorem 2.3. We here prove the estimates only for $\lambda \in \Sigma(\eta_4, \theta_4)$. The other case can be treated by using the decomposition $(\lambda + L)^{-1}f = \widetilde{R}^{(0)}(\lambda)f + \widetilde{R}^{(\infty)}(\lambda)f$ as in the proof of Theorem 2.1.

Let us denote the integral kernels of $(\lambda + L)^{-1}$, $G(\lambda)f$ and $K(\lambda)f$ by $R(\lambda, x' - y', x_n, y_n)$, $G(\lambda, x' - y', x_n, y_n)$ and $K(\lambda, x' - y', x_n, y_n)$, respectively. Let $E(x_n, y_n) = E^{(1)}(x_n, y_n) + E^{(2)}(x_n, y_n)$ be defined as in the proof of

Theorem 2.2. It holds that

$$\sup_{0 \le y_n \le a} \int_0^a E(x_n, y_n) \, dx_n + \sup_{0 \le x_n \le a} \int_0^a E(x_n, y_n) \, dy_n \le C\sigma(\lambda, \xi')^{-\frac{1}{2}}.$$

Therefore, by Lemmas 4.6-4.8 and 5.5, we have

$$\begin{aligned} \|\partial_x^k \widetilde{Q}(\lambda+L)^{-1} \widetilde{Q}f\|_1 &\leq \sup_{0 \leq y_n \leq a} \|\partial_x^k \widetilde{Q}R(\lambda,\cdot,\cdot,y_n)\|_1 \|\widetilde{Q}f\|_1 \\ &\leq \frac{C}{(|\lambda|+1)^{1-\frac{k}{2}}} \|\widetilde{Q}f\|_1, \quad k = 0, 1, \end{aligned}$$

and

$$\begin{aligned} \|\partial_x^k \widetilde{Q}(\lambda+L)^{-1} \widetilde{Q}f\|_{\infty} &\leq \sup_{0 \leq x_n \leq a} \|\partial_x^k \widetilde{Q}R(\lambda, \cdot, x_n, \cdot)\|_1 \|\widetilde{Q}f\|_{\infty} \\ &\leq \frac{C}{(|\lambda|+1)^{1-\frac{k}{2}}} \|\widetilde{Q}f\|_{\infty}, \quad k = 0, 1. \end{aligned}$$

Similarly one can estimate $Q_0(\lambda + L)^{-1}f$ to obtain the desired estimate.

We next consider $\partial_x Q_0(\lambda + L)^{-1} \widetilde{Q} f$. We here estimate only $\partial_x Q_0 G(\lambda) Q_n f$. Since $\partial_{x'}^{\alpha'} G(\lambda) f = G(\lambda) \partial_{x'}^{\alpha'} f$, we see from Lemmas 4.6 and 5.5 that

$$\|\partial_{x'}^{\alpha'}Q_0G(\lambda)Q_nf\|_p \le \frac{C}{|\lambda|+1}\|\partial_{x'}^{\alpha'}Q_nf\|_p, \quad p=1,\infty, \ |\alpha'|=1.$$

Since $\partial_{x_n}^2(g_{\mu_2}^{(D)}\hat{f}) = \mu_2^2 g_{\mu_2}^{(D)}\hat{f} - \hat{f}$ and $\mu_2^2 = \frac{\lambda^2}{\nu_1 \lambda + \gamma^2} + |\xi'|^2$, we have

$$\partial_{x_n} Q_0 G(\lambda) Q_n f$$

= $-\frac{\gamma}{\nu_1 \lambda + \gamma^2} \left\{ \frac{\lambda^2}{\nu_1 \lambda + \gamma^2} Q_0 G(\lambda) Q_n f + \nabla' Q_0 G(\lambda) [\nabla' Q_n f] - Q_n f \right\}.$

It then follows from Lemmas 4.6 and 5.5 that

$$\|\partial_{x_n} Q_0 G(\lambda) Q_n f\|_p \le \frac{C}{|\lambda|+1} \{ \|Q_n f\|_p + \|\nabla' Q_n f\|_p \}, \quad p = 1, \infty.$$

The remaining part of $\partial_x Q_0(\lambda + L)^{-1} \tilde{Q} f$ can be estimated similarly. One can also estimate $\partial_x \tilde{Q}(\lambda + L)^{-1} Q_0 f$ in a similar manner.

We next estimate $\partial_x Q_0(\lambda + L)^{-1} Q_0 f$. Consider first $Q_0 K(\lambda) Q_0 f$. We set

$$\widetilde{k}(\lambda, x', x_n, y_n) = \mathscr{F}^{-1} \left[h_{\mu_2}(\lambda, \xi', x_n) \beta_0(\lambda, \xi', y_n) \right] (x')$$

$$\widetilde{K}(\lambda) f = \int_{\mathbf{R}^{n-1}} \widetilde{k}(x' - y', x_n, y_n) Q_0 f(y', y_n) \, dy' dy_n.$$

Then

$$Q_0 K(\lambda) Q_0 f = -\frac{\gamma}{\nu_1 \lambda + \gamma^2} \nabla' \widetilde{K}(\lambda) \left[\nabla' Q_0 f \right].$$

It then follows from Lemmas 4.7, 4.8 and 5.5 that

$$\|\partial_{x'}Q_0K(\lambda)Q_0f\|_p \le \frac{C}{|\lambda|+1} \|\partial_{x'}\nabla'Q_0f\|_p, \quad p=1,\infty.$$

Also, since

$$Q_0 K(\lambda) Q_0 f = -\frac{\gamma}{\nu_1 \lambda + \gamma^2} \widetilde{K}(\lambda) \left[\Delta' Q_0 f \right],$$

we similarly obtain, by Lemmas 4.7, 4.8 and 5.5,

$$\|\partial_{x_n}Q_0K(\lambda)Q_0f\|_p \le \frac{C}{|\lambda|+1} \|\Delta'Q_0f\|_p.$$

As for $\partial_x Q_0 G(\lambda) Q_0 f$, one can treat it as in the case of $\partial_x \tilde{Q}(\lambda + L)^{-1} \tilde{Q} f$.

Let us finally consider $\partial_x \tilde{Q}(\lambda + L)^{-1} f$, assuming that $\tilde{Q}f\Big|_{x_n=0,a} = 0$. As for $\partial_x \tilde{Q}(\lambda + L)^{-1} \tilde{Q}f$, we rewrite it as in the proof of the last inequality of Theorem 2.1. One can then estimate it similarly to the case of $\partial_x^k \tilde{Q}(\lambda + L)^{-1} \tilde{Q}f$ with $k \leq 1$ to obtain the desired inequality. The estimate for $\partial_x \tilde{Q}(\lambda + L)^{-1} Q_0 f$ has been already obtained. This completes the proof.

We finally prove Theorems 2.5–2.7.

Proof of Theorems 2.5–2.7. We prove Theorems 2.5–2.7 for $\tilde{\eta} = \min \{\eta_2, \eta_3\}$ and $\frac{\pi}{2} < \tilde{\theta} < \min \{\theta_2, \theta_3, \theta_4\}$, where η_j (j = 2, 3) and θ_j (j = 2, 3, 4) are the numbers given in Lemmas 4.3 and 4.4. For this $\{\tilde{\eta}, \tilde{\theta}\}$ we see from Lemma 4.3 that $\Sigma(-\tilde{\eta}, \tilde{\theta}) \subset \rho(-\hat{L}_{\xi'})$ for $|\xi'| \ge r$. Furthermore, we deduce that $\Sigma(-\tilde{\eta}, \tilde{\theta}) \subset \Sigma(-\eta_2, \theta_2) \cap \Sigma(-\eta_3, \theta_3)$ and that $\overline{\Sigma(-\tilde{\eta}, \tilde{\theta})} - \Sigma(\eta_4, \theta_4)$ is a compact set. Here η_4 and θ_4 are the numbers given in Lemma 4.4.

In view of the proof of Theorems 2.1–2.3, the desired estimates for $R^{(1)}(\lambda)f$ hold for $\lambda \in \Sigma(\eta_4, \theta_4)$. In case $\lambda \in \Sigma(-\tilde{\eta}, \tilde{\theta}) - \Sigma(\eta_4, \theta_4)$, we decompose $R^{(1)}(\lambda)f$ in the following way. Let $\chi(\xi')$ and $\tilde{\chi}(\xi')$ be the cut-off functions given in the definition of $R^{(1)}(\lambda)f$ and in the proof of Theorem 2.1, respectively. We write $R^{(1)}(\lambda)f$ as

$$R^{(1)}(\lambda)f = \tilde{R}^{(2)}(\lambda)f + \tilde{R}^{(\infty)}(\lambda)f,$$

where

$$\widetilde{R}^{(2)}(\lambda)f = \mathscr{F}_{\xi'}^{-1}\left[\chi(\xi')(1-\widetilde{\chi}(\xi'))\widehat{R}(\lambda,\xi')\widehat{f}\right]$$

and

$$\widetilde{R}^{(\infty)}(\lambda)f = \mathscr{F}_{\xi'}^{-1}\left[\chi(\xi')\widetilde{\chi}(\xi')\widehat{R}(\lambda,\xi')\widehat{f}\right].$$

By Lemma 3.4 and Lemma 4.3 (ii) we see that $D(\lambda, \xi') \neq 0$ on the compact set $\overline{\Sigma(-\tilde{\eta}, \tilde{\theta})} - \Sigma(\eta_4, \theta_4)$. Therefore, as above, one can see that $R^{(1)}(\lambda)f$ has the desired estimates for $\lambda \in \Sigma(-\tilde{\eta}, \tilde{\theta}) - \Sigma(\eta_4, \theta_4)$ as in the proof of Theorems 2.1–2.3. This completes the proof.

References

- T. Abe and Y. Shibata, On a resolvent estimate of the Stokes equation on an infinite layer, J. Math. Soc. Japan 55 (2003), pp. 469–497.
- [2] T. Abe and Y. Shibata, On a resolvent estimate of the Stokes equation on an infinite layer part 2, $\lambda = 0$ case, J. Math. Fluid Mech. 5 (2003), pp. 245–274.
- [3] H. Abels and M. Wiegner, Resolvent estimates for the Stokes operator on an infinite layer, Differential and Integral Equations 18 (2005), pp. 1081–1110.
- [4] J. Bergh and J. Löfström, Interpolation Spaces, An Introduction, Springer-Verlag, Berlin, Heidelberg, New York (1976).
- [5] Y. Kagei, Asymptotic behavior of the semigroup associated with the linearized compressible Navier-Stokes equation in an infinite layer, preprint.
- [6] Y. Kagei and T. Kobayashi, On large time behavior of solutions to the Compressible Navier-Stokes Equations in the half space in R³, Arch. Rational Mech. Anal. 165 (2002), pp. 89–159.
- [7] Y. Kagei, and T. Kobayashi, Asymptotic behavior of solutions to the compressible Navier-Stokes equations on the half space, Arch. Rational Mech. Anal. 177 (2005), pp. 231–330.

[8] T. Kobayashi and Y. Shibata, Remarks on the rate of decay of solutions to linearized compressible Navier-Stokes equations, Pacific J. Math. 207 (2002), pp. 199–234.

List of MHF Preprint Series, Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

- MHF2005-1 Hideki KOSAKI Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA Simple estimators for non-linear Markovian trend from sampled data: I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA Estimation for a discretely observed small diffusion process with a linear drift
- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA AIC for ergodic diffusion processes from discrete observations

- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI Nonlinear regression modeling via regularized wavelets and smoothing parameter selection
- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO, Yoshihiro MIZOGUCHI & Yasuo KAWAHARA On reversible cellular automata with finite cell array

- MHF2005-28 Toru KOMATSU Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI An explicit formula of the Shapley value for a cooperative game induced from the conjugate point
- MHF2005-34 Hideki MURAKAWA A regularization of a reaction-diffusion system approximation to the two-phase Stefan problem
- MHF2006-1 Masahisa TABATA Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme
- MHF2006-2 Ken-ichi MARUNO & G R W QUISPEL Construction of integrals of higher-order mappings
- MHF2006-3 Setsuo TANIGUCHI On the Jacobi field approach to stochastic oscillatory integrals with quadratic phase function
- MHF2006-4 Kouji HASHIMOTO, Kaori NAGATOU & Mitsuhiro T. NAKAO A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains
- MHF2006-5 Hidefumi KAWASAKI A duality theory based on triangular cylinders separating three convex sets in \mathbb{R}^n
- MHF2006-6 Raimundas VIDŪNAS Uniform convergence of hypergeometric series
- MHF2006-7 Yuji KODAMA & Ken-ichi MARUNO N-Soliton solutions to the DKP equation and Weyl group actions

- MHF2006-8 Toru KOMATSU Potentially generic polynomial
- MHF2006-9 Toru KOMATSU Generic sextic polynomial related to the subfield problem of a cubic polynomial
- MHF2006-10 Shu TEZUKA & Anargyros PAPAGEORGIOU Exact cubature for a class of functions of maximum effective dimension
- MHF2006-11 Shu TEZUKA On high-discrepancy sequences
- MHF2006-12 Raimundas VIDŪNAS Detecting persistent regimes in the North Atlantic Oscillation time series
- MHF2006-13 Toru KOMATSU Tamely Eisenstein field with prime power discriminant
- MHF2006-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO Generating function associated with the Hankel determinant formula for the solutions of the Painlevé IV equation
- MHF2006-15 Raimundas VIDŪNAS Darboux evaluations of algebraic Gauss hypergeometric functions
- MHF2006-16 Masato KIMURA & Isao WAKANO New mathematical approach to the energy release rate in crack extension
- MHF2006-17 Toru KOMATSU Arithmetic of the splitting field of Alexander polynomial
- MHF2006-18 Hiroki MASUDA Likelihood estimation of stable Lévy processes from discrete data
- MHF2006-19 Hiroshi KAWABI & Michael RÖCKNER Essential self-adjointness of Dirichlet operators on a path space with Gibbs measures via an SPDE approach
- MHF2006-20 Masahisa TABATA Energy stable finite element schemes and their applications to two-fluid flow problems
- MHF2006-21 Yuzuru INAHAMA & Hiroshi KAWABI Asymptotic expansions for the Laplace approximations for Itô functionals of Brownian rough paths
- MHF2006-22 Yoshiyuki KAGEI Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer