

## アスコルビン酸大量投与の血漿カルニチンに及ぼす効果

伊規須, 英輝  
九州大学健康科学センター

藤野, 武彦  
九州大学健康科学センター

銅直, 孝子  
九州大学健康科学センター

<https://doi.org/10.15017/397>

---

出版情報 : 健康科学. 5, pp.53-54, 1983-03-20. 九州大学健康科学センター  
バージョン :  
権利関係 :

## High Ascorbic Acid Intake and Plasma Carnitine

Hideki IGISU, Takehiko FUJINO, Takako DONAO

アスコルビン酸大量投与の血漿カルニチンに及ぼす効果

伊規須 英輝, 藤野 武彦, 銅直孝子

Carnitine<sup>1, 2</sup> as well as ascorbic acid<sup>3</sup> were reported to ameliorate some type of hyperlipidemia. On the other hand, a recent study<sup>4</sup> showed that ascorbic acid is a co-factor in the biosynthesis of carnitine. We therefore studied whether or not high ascorbic acid intake affects plasma level of carnitine.

Nineteen male university students (18-21 years old) who gave their informed consent participated in the study. Eleven students ("subjects") took ascorbic acid (3.0 grams per day) and 8 students ("controls") took placebo (composed mainly of tartaric acid) for three months. We obtained heparinized blood after overnight fast and measured carnitine<sup>5</sup> and ascorbic acid<sup>6</sup> in plasma at 1, 2 and 3 months after the study was started.

The ascorbic acid level was significantly higher in "subjects" than in "controls". However, no difference was found in the carnitine level (see accompanying table). Thus, high ascorbic acid intake does not appear to change plasma carnitine level.

Table. Plasma ascorbic acid and carnitine after high ascorbic acid intake (mean±SD).

	1 month	2 months	3 months
Ascorbic acid (mg/100ml)			
subjects	1.62*±0.32 (n=11)	1.63*±0.25 (n=11)	1.42*±0.52 (n=11)
controls	1.08 ±0.19 (n=8)	1.14 ±0.08 (n=7)	0.72 ±0.20 (n=8)
Carnitine (μmoles/100ml)			
subjects	4.75 ±0.91 (n=11)	4.83 ±0.70 (n=11)	4.58 ±0.78 (n=11)
controls	5.08 ±0.85 (n=8)	5.20 ±0.69 (n=7)	4.86 ±0.88 (n=8)

\*p<0.005

### References

1. Maebashi M, Kawamura N, Sato M, Imamura A, Yoshinaga K. Lipid-lowering effect of carnitine in patients with type-IV hyperlipoproteinemia. *Lancet* 1978 ; ii : 805-07.
2. Lacour B, DiGiulio S, Chanard J, Ciancioni C, Haguët M, Lebkiiri B, Basile C, Druëke T, Assan R, Funck-Brentano JL. Carnitine improves lipid anomalies in haemodialysis patients. *Lancet* 1980 ; ii ; 763-65.
3. Geoly KL, Diamond LH. Ascorbic acid and hypertriglyceridemia. *Ann Intern Med*

- 1980 ; 93 : 511.
4. Hulse JD, Ellis SR, Henderson LM. Carnitine biosynthesis. *J Biol Chem* 1978 ; 253 : 1654-59.
  5. Pearson DJ, Chase JFA, Tubbs PK. The assay of (-)-carnitine and its o-acyl derivatives. The DTNB method. In: *Methods in enzymology*. ed. Lowenstein JM, New York; Academic Press, 1969 ; 14 : p.615-17.
  6. Okamura M. An improved method for determination of 1-ascorbic acid and 1-dehydroascorbic acid in blood plasma. *Clin Chim Acta* 1980 ; 103 : 259-68.