

A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions

Kobayashi, Kenta
Faculty of Mathematics, Kyushu University

<https://hdl.handle.net/2324/4023>

出版情報 : 2007-02-14. 九州大学大学院数理学研究院
バージョン :
権利関係 :



MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

**A constructive a priori error
estimation for finite element
discretizations in a non-convex
domain using singular functions**

K. Kobayashi

MHF 2007-9

(Received February 14, 2007)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions

Kenta Kobayashi¹

Abstract

In solving elliptic problems by the finite element method in a bounded domain with a re-entrant corner, the rate of convergence could be improved by adding a singular function to the usual C^0 approximating basis. When the domain is enclosed by line segments which forms a corner of $\pi/2$ or $3\pi/2$, we have obtained an explicit a priori H^1 error estimation of $O(h)$ for such a finite element solution of the Poisson equation. Particularly, we emphasize that all constants in our error estimates are numerically determined, which plays an essential role in the numerical verification of solutions for non-linear elliptic problems.

Key Words: Finite element method, A priori error estimation, Poisson equation

Mathematics Subject Classification (2000): 65N30, 65N15, 35J05

1 Introduction

In this paper, we consider the elliptic problem on polygonal domain Ω which is enclosed by line segments and right angles. Ω is assumed to be connected but it is not necessarily simply connected. First of all, we assume $\Omega = \Omega_0$, where Ω_0 is an inside of the L-shape domain shown in Fig.1. The general case is described in Section 3.




Figure 1: The shape of Ω_0

¹Faculty of Mathematics, Kyushu University, Fukuoka 812-8581 Japan.

This research is partly supported by the 21st Century COE program of Faculty of Mathematics, Kyushu University

For $f \in L^2(\Omega)$, we consider the weak solution of the following partial differential equation

$$(1.1) \quad \begin{cases} -\Delta u = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

It is known that u has a singular function representation [9][10]

$$(1.2) \quad u(x, y) = w(x, y) + \lambda\sigma(x, y)$$

where $w(x, y) \in H^2(\Omega) \cap H_0^1(\Omega)$, λ is a constant, $\sigma(x, y) \in H_0^1(\Omega)$ and

$$\sigma(x, y) \sim r^{2/3} \sin\left(\frac{2}{3}\theta\right)$$

in a neighborhood of the origin. Here, (r, θ) is the polar coordinates of (x, y) where θ satisfies $0 \leq \theta < 2\pi$.

There is arbitrariness in the choice of σ . To simplify the calculation of H_0^1 inner product, we take

$$\sigma(x, y) = (1 - x^2)(1 - y^2)r^{2/3} \sin\left(\frac{2}{3}\theta\right),$$

in this paper.

Solving this problem by the finite element method, we use the square mesh with mesh size h . The mesh of $h = 1/8$ is shown in Fig.2.




Figure 2: The square mesh when $h = 1/8$

We use the piecewise bilinear function,

$$\phi_{i,j}(x, y) \equiv \max\left(1 - \left|\frac{x}{h} - i\right|, 0\right) \cdot \max\left(1 - \left|\frac{y}{h} - j\right|, 0\right),$$

as the finite element basis.

We define Φ_h by the set of functions $\phi_{i,j}$ in $H_0^1(\Omega)$,

$$\Phi_h = \left\{ \phi_{i,j} \mid (ih, jh) \in \Omega \right\}.$$

Since u does not generally have H^2 regularity, we can not obtain $O(h)$ error estimates with H_0^1 norm by using this approximating basis Φ_h .

Therefore, we adopt $\Phi_h \cup \{\sigma\}$ as the finite element basis. In this case, it is known that the following error estimation holds [8][9][10],

$$\|u - u_h\|_{H_0^1(\Omega)} \leq Ch\|f\|_{L^2(\Omega)},$$

where u_h is the finite element solution. The following $O(h^2)$ estimation for the L^2 -error is also obtained by the Aubin-Nitsche trick [6],

$$\|u - u_h\|_{L^2(\Omega)} \leq C^2 h\|f\|_{L^2(\Omega)}.$$

The main purpose of this paper is to obtain this constant C .

The coefficient λ in (1.2) is often called the stress intensity factor in a context of mechanics. In our error estimation, the explicit evaluation of the coefficient λ is essential (Lemma 4.2 and Lemma 4.6 in section 4). For the coefficient λ , the following extraction formula holds [9][10]

$$\lambda = \frac{1}{\pi} \left\{ \iint_{\Omega} f \eta s_- dx dy + \iint_{\Omega} u \Delta(\eta s_-) dx dy \right\},$$

where

$$s_- = r^{-2/3} \sin\left(\frac{2}{3}\theta\right),$$

and η is a cut-off function which equals one at the origin and zero on $\{(x, y) \mid \max(|x|, |y|) = 1\}$. Poincaré-Friedrichs inequality is needed to evaluate λ by this extraction formula. However, since Poincaré-Friedrichs inequality is reduced to a kind of a problem of eigenvalue bounds, it is not easy to obtain good estimation, except for the case that Ω is simple domain such as rectangle. In this paper, instead of using cut-off function, we use the maximum principle for the super harmonic functions to evaluate λ directly.

There are several approaches to deal with the lack of regularity at the re-entrant corner. The method in this paper is based on [8]. This method is a simple finite element method but is enough to obtain H_0^1 and L^2 error bounds. The dual singular function method (DSFM) [3][7] is presented to obtain better approximation for the coefficient λ . DSFM consists of a system of w and λ which is derived from extraction formula, and is often implemented as an iterative procedure. A multigrid version of this method appears in [4]. An efficient method using improved extraction formula was presented in [5]. Another useful method is based on the local mesh refinement [2]. The advantage of using mesh refinement is that calculation of the element matrix is easy, because the information about the singular function is not needed.

In many applications, it is useful to obtain explicit error estimation. For example, numerical verification method for nonlinear problems are based on explicit error bounds for linear equations [12][16].

We should mention that in [15] an explicit error estimation for non-convex domain is proposed. However, the error estimation in [15] needs specific information about the finite element solution of (1.1), and furthermore, the order of their error estimation is

about $O(h^{0.6927})$ with H_0^1 norm. What we are going to present is much higher order error estimation which can be a priori calculated.

The present paper is organized as follows. In section 2, we present a priori error estimation when the case of Ω is simple L-shape domain. The general case is explained in section 3. Section 4 contains proof of lemmas which appear in section 2 and 3. We show numerical results in section 5 and conclude this paper by section 6.

Throughout this paper, we take the angle of polar coordinates in $[0, 2\pi)$. Let 1_A denotes the function which takes value 1 if condition A holds, and takes value 0 otherwise.

2 A priori error estimation

The main purpose of this section is to prove the following theorem. Lemmas appear in this section will be proved in section 4.

Theorem 2.1 *When $\Omega = \Omega_0$, as to the finite element solution u_h by using $\Phi_h \cup \{\sigma\}$ as the basis, the following error estimation holds,*

$$\begin{aligned}\|u - u_h\|_{H_0^1(\Omega)} &\leq 1.156h\|f\|_{L^2(\Omega)}, \\ \|u - u_h\|_{L^2(\Omega)} &\leq 1.335h^2\|f\|_{L^2(\Omega)}.\end{aligned}$$

Proof. We represent the exact solution as,

$$(2.1) \quad u = w + \lambda\sigma.$$

Where λ is a constant which depends on Ω and f , and w is a function which belongs to $H^2(\Omega) \cap H_0^1(\Omega)$.

Define w_h as the bilinear interpolation of w . Then, since $w \in H^2(\Omega) \cap H_0^1(\Omega)$,

$$\|w - w_h\|_{H_0^1(\Omega)} \leq \frac{h}{\pi}|w|_{H^2(\Omega)}$$

holds [13].

Since Ω is a polygonal domain, the following equality holds [11].

$$|w|_{H^2(\Omega)} = \|\Delta w\|_{L^2(\Omega)}.$$

We immediately have,

$$\|w - w_h\|_{H_0^1(\Omega)} \leq \frac{h}{\pi}\|\Delta w\|_{L^2(\Omega)}.$$

Let

$$\tilde{u}_h = w_h + \lambda\sigma$$

then, we have

$$(2.2) \quad \|u - \tilde{u}_h\|_{H_0^1(\Omega)} \leq \frac{h}{\pi}\|\Delta w\|_{L^2(\Omega)}.$$

Form (2.1),

$$(2.3) \quad \|\Delta w\|_{L^2(\Omega)} \leq \|f\|_{L^2(\Omega)} + |\lambda| \|\Delta \sigma\|_{L^2(\Omega)}.$$

From Lemma 4.1,

$$(2.4) \quad \left\| \Delta \left\{ (1-x^2)(1-y^2)r^{2/3} \sin\left(\frac{2}{3}\theta\right) \right\} \right\|_{L^2(\Omega)} \leq \sqrt{\frac{4000}{81} - \frac{11713}{1782}}\pi.$$

For the coefficient λ , Lemma 4.2 implies

$$(2.5) \quad |\lambda| \leq \frac{1}{\pi} \left\| \left(r^{-2/3} - 2^{-2/3}r^{2/3} \right) \sin\left(\frac{2}{3}\theta\right) \right\|_{L^2(\Omega)} \|f(x, y)\|_{L^2(\Omega)}$$

and, from Lemma 4.3,

$$\left\| \left(r^{-2/3} - 2^{-2/3}r^{2/3} \right) \sin\left(\frac{2}{3}\theta\right) \right\|_{L^2(\Omega)} \leq \sqrt{\frac{3 \cdot 2^{1/3}}{5}}\pi.$$

Consequently, from (2.2),(2.3),(2.4) and (2.5),

$$\begin{aligned} |u - \tilde{u}_h|_{H_0^1(\Omega)} &\leq \frac{h}{\pi} \left(1 + \sqrt{\frac{4000}{81} - \frac{11713}{1782}}\pi \cdot \frac{1}{\pi} \sqrt{\frac{3 \cdot 2^{1/3}}{5}}\pi \right) \|f\|_{L^2(\Omega)} \\ &= 1.1552884253 \cdots h \|f\|_{L^2(\Omega)} \leq 1.156h \|f\|_{L^2(\Omega)}. \end{aligned}$$

Since the finite element solution u_h is the best approximation in H_0^1 space, we have

$$\|u - u_h\|_{H_0^1(\Omega)} \leq \|u - \tilde{u}_h\|_{H_0^1(\Omega)} \leq 1.156h \|f\|_{L^2(\Omega)}.$$

$$\|u - u_h\|_{L^2(\Omega)} \leq 1.1553^2 h^2 \|f\|_{L^2(\Omega)} \leq 1.335h^2 \|f\|_{L^2(\Omega)}$$

is also obtained by the Aubin-Nitsche trick. \square

3 Generalization

We consider general cases in this section. We suppose that Ω is enclosed by line segments which forms a corner of $\pi/2$ or $3\pi/2$. Ω is assumed to be connected but it is not necessarily simply connected (Fig.3).

The following singular bases are used together with usual interpolating basis

$$T_k \sigma \left(\frac{x}{l_k}, \frac{y}{l_k} \right), \quad (k = 1, \dots, n),$$

where n is a number of re-entrant corner,

$$\sigma(x, y) = \begin{cases} (1-x^2)(1-y^2)r^{2/3} \sin\left(\frac{2}{3}\theta\right), & ((x, y) \in \Omega_0), \\ 0, & (\text{otherwise}), \end{cases}$$




Figure 3: Admissible pattern




Figure 4: Admissible pattern

and T_k is a combination of parallel translation and rotation.

l_k denotes sizes of the singular bases. Different sizes of the singular bases are admissible. It is also admissible even if some part of the support of the singular bases are overlapped (Fig.4).

There are some restrictions on defining the singular bases. Let us now suppose that Γ is the support of a singular basis, and $\partial\Gamma$ consists of line segments $\gamma_1 \sim \gamma_6$ where γ_1 and γ_6 form the re-entrant corner (Fig.5). In this case, γ_1 and γ_6 must be contained in $\partial\Omega$ and $\gamma_2 \sim \gamma_5$ must coincide with the grid line of the mesh. Therefore, both Fig.6 and Fig.7 are not admissible.

In this situation, we have the following Theorem.

Theorem 3.1 *For the finite element solution u_h with the basis $\Phi_h \cup \left\{ T_k \sigma(x/l_k, y/l_k) \mid k = 1, \dots, n \right\}$, the following error estimation holds,*

$$\begin{aligned} \|u - u_h\|_{H_0^1(\Omega)} &\leq \left(0.319 + \sum_{k=1}^n \sqrt{0.971 + 1.469 \frac{|\Omega|}{l_k^2}} \right) h \|f\|_{L^2(\Omega)}, \\ \|u - u_h\|_{L^2(\Omega)} &\leq \left(0.319 + \sum_{k=1}^n \sqrt{0.971 + 1.469 \frac{|\Omega|}{l_k^2}} \right)^2 h^2 \|f\|_{L^2(\Omega)}, \end{aligned}$$




Figure 5: The support of a singular basis




Figure 6: Nonadmissible pattern




Figure 7: Nonadmissible pattern

where $|\Omega|$ denotes the area of Ω .

Proof. Let

$$u(x, y) = w(x, y) + \sum_{k=1}^n \lambda_k T_k \sigma \left(\frac{x}{l_k}, \frac{y}{l_k} \right), \quad w(x, y) \in H^2(\Omega) \cap H_0^1(\Omega).$$

be the exact solution.

We define \tilde{u}_h as follows,

$$\tilde{u}_h(x, y) = w_h(x, y) + \sum_{k=1}^n \lambda_k T_k \sigma \left(\frac{x}{l_k}, \frac{y}{l_k} \right)$$

where w_h denotes the bilinear interpolation of w .

We also define

$$\bar{\sigma}(x, y) = \begin{cases} (1 - r)^2 r^{2/3} \sin \left(\frac{2}{3} \theta \right), & (r < 1), \\ 0, & (r \geq 1). \end{cases}$$

Now, we exclude the grid of the square mesh from Ω and define it as Ω_* . Then,

$$\begin{aligned}
\|u - \tilde{u}_h\|_{H_0^1(\Omega)} &= \|w - w_h\|_{H_0^1(\Omega_*)} \leq \frac{h}{\pi} |w|_{H^2(\Omega_*)} \\
&= \frac{h}{\pi} \left\| u - \sum_{k=1}^n \lambda_k T_k \sigma \left(\frac{x}{l_k}, \frac{y}{l_k} \right) \right\|_{H^2(\Omega_*)} \\
&\leq \frac{h}{\pi} \left\| u - \sum_{k=1}^n \lambda_k T_k \bar{\sigma} \left(\frac{x}{l_k}, \frac{y}{l_k} \right) \right\|_{H^2(\Omega)} \\
&\quad + \frac{h}{\pi} \left\| \sum_{k=1}^n \lambda_k T_k \left(\sigma \left(\frac{x}{l_k}, \frac{y}{l_k} \right) - \bar{\sigma} \left(\frac{x}{l_k}, \frac{y}{l_k} \right) \right) \right\|_{H^2(\Omega_*)} \\
&= \frac{h}{\pi} \left\| \Delta \left(u - \sum_{k=1}^n \lambda_k T_k \bar{\sigma} \left(\frac{x}{l_k}, \frac{y}{l_k} \right) \right) \right\|_{L^2(\Omega)} \\
&\quad + \frac{h}{\pi} \left\| \sum_{k=1}^n \lambda_k \Delta T_k \left(\sigma \left(\frac{x}{l_k}, \frac{y}{l_k} \right) - \bar{\sigma} \left(\frac{x}{l_k}, \frac{y}{l_k} \right) \right) \right\|_{L^2(\Omega_*)} \\
&\leq \frac{h}{\pi} \|f(x, y)\|_{L^2(\Omega)} \\
&\quad + \frac{h}{\pi} \sum_{k=1}^n \frac{|\lambda_k|}{l_k} \left(\|\Delta \bar{\sigma}(x, y)\|_{L^2(\Omega_0)} + \left\| \Delta \left(\sigma(x, y) - \bar{\sigma}(x, y) \right) \right\|_{L^2(\Omega_0)} \right).
\end{aligned}$$

From Lemma 4.4,

$$\|\Delta \bar{\sigma}(x, y)\|_{L^2(\Omega_0)} = \frac{3\sqrt{\pi}}{2}.$$

Lemma 4.5 implies that

$$\left\| \Delta \left(\sigma(x, y) - \bar{\sigma}(x, y) \right) \right\|_{L^2(\Omega_0)} \leq \sqrt{\frac{4000}{81} - \frac{232367}{46332} \pi}.$$

Then, we have

$$\|u - \tilde{u}_h\|_{H_0^1(\Omega)} \leq \frac{h}{\pi} \|f\|_{L^2(\Omega)} + \frac{h}{\pi} \left(\frac{3\sqrt{\pi}}{2} + \sqrt{\frac{4000}{81} - \frac{232367}{46332} \pi} \right) \sum_{k=1}^n \frac{|a_k|}{l_k}.$$

From Lemma 4.6, we have the following evaluation

$$\begin{aligned}
|\lambda_k| &\leq \left\{ \iint_{\Omega} 1_{0 \leq \theta < 3\pi/2} \cdot \left| G_{l_k} \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right) \right|^2 dx dy \right. \\
&\quad \left. + \iint_{\Omega} 1_{3\pi/2 \leq \theta < 2\pi} \cdot \left| G_{l_k} \left(r, \frac{1}{9} (8 + \cos(4\theta)), 1 \right) \right|^2 dx dy \right\}^{1/2} \|f(x, y)\|_{L^2(\Omega)},
\end{aligned}$$

where

$$G_l(r, X, Y) = \frac{l^{2/3}}{\sqrt{2\pi}} \sqrt{\sqrt{r^{-8/3} + l^{-8/3} - 2r^{-4/3}l^{-4/3}X} + l^{-4/3} - r^{-4/3}Y}.$$

It follows from Lemma 4.7 that

$$|\lambda_k| \leq \frac{1}{\pi} \sqrt{\left(\frac{5}{2} - \frac{3\pi}{8}\right) l_k^2 + 2|\Omega|} \cdot \|f\|_{L^2(\Omega)}.$$

Then,

$$\begin{aligned} & \|u - \tilde{u}_h\|_{H_0^1(\Omega)} \\ & \leq \frac{h}{\pi} \left(1 + \left(\frac{3\sqrt{\pi}}{2} + \sqrt{\frac{4000}{81} - \frac{232367}{46332}\pi} \right) \sum_{k=1}^n \frac{1}{\pi} \sqrt{\left(\frac{5}{2} - \frac{3\pi}{8}\right) + \frac{2|\Omega|}{l_k^2}} \right) \\ & \quad \times \|f\|_{L^2(\Omega)} \\ & = \left(0.31830988 \cdots + \sum_{k=1}^n \sqrt{0.97070784 \cdots + 1.46865243 \cdots \frac{|\Omega|}{l_k^2}} \right) h \|f\|_{L^2(\Omega)} \\ & \leq \left(0.319 + \sum_{k=1}^n \sqrt{0.971 + 1.469 \frac{|\Omega|}{l_k^2}} \right) h \|f\|_{L^2(\Omega)}. \end{aligned}$$

Since the finite element solution u_h is the best approximation in H_0^1 space,

$$\begin{aligned} \|u - u_h\|_{H_0^1(\Omega)} & \leq \|u - \tilde{u}_h\|_{H_0^1(\Omega)} \\ & \leq \left(0.319 + \sum_{k=1}^n \sqrt{0.971 + 1.469 \frac{|\Omega|}{l_k^2}} \right) h \|f\|_{L^2(\Omega)}. \end{aligned}$$

We also obtain

$$\|u - u_h\|_{L^2(\Omega)} \leq \left(0.319 + \sum_{k=1}^n \sqrt{0.971 + 1.469 \frac{|\Omega|}{l_k^2}} \right)^2 h^2 \|f\|_{L^2(\Omega)}$$

by the Aubin-Nitsche trick. □

4 Lemmas

Lemma 4.1

$$\left\| \Delta \left\{ (1 - x^2)(1 - y^2)r^{2/3} \sin\left(\frac{2}{3}\theta\right) \right\} \right\|_{L^2(\Omega_0)} \leq \sqrt{\frac{4000}{81} - \frac{11713}{1782}\pi}.$$

Proof.

$$\begin{aligned} & \Delta \left\{ (1 - x^2)(1 - y^2)r^{2/3} \sin\left(\frac{2}{3}\theta\right) \right\} \\ & = \frac{2}{3}r^{2/3} \left\{ (4r^2 - 10) \sin\left(\frac{2}{3}\theta\right) + r^2 \sin\left(\frac{10}{3}\theta\right) \right\} \\ & = \frac{2}{3}r^{2/3} \sin\left(\frac{2}{3}\theta\right) \left\{ 9r^2 - 10 - 20r^2 \sin^2\left(\frac{2}{3}\theta\right) + 16r^2 \sin^4\left(\frac{2}{3}\theta\right) \right\}. \end{aligned}$$

When $1 \leq r \leq \sqrt{2}$,

$$\begin{aligned}
& \Delta \left\{ (1-x^2)(1-y^2)r^{2/3} \sin \left(\frac{2}{3}\theta \right) \right\} \\
&= \frac{2}{3}r^{2/3} \sin \left(\frac{2}{3}\theta \right) \left\{ 9r^2 - 10 - 4r^2 \sin^2 \left(\frac{2}{3}\theta \right) \right. \\
&\quad \left. - 16r^2 \sin^2 \left(\frac{2}{3}\theta \right) \left(1 - \sin^2 \left(\frac{2}{3}\theta \right) \right) \right\} \\
&\leq \frac{2}{3}r \sin \left(\frac{2}{3}\theta \right) \left\{ 8 - 4r^2 \sin^2 \left(\frac{2}{3}\theta \right) \right\},
\end{aligned}$$

and

$$\begin{aligned}
& \Delta \left\{ (1-x^2)(1-y^2)r^{2/3} \sin \left(\frac{2}{3}\theta \right) \right\} \\
&= -\frac{2}{3}r^{2/3} \sin \left(\frac{2}{3}\theta \right) \left\{ 10 - 4r^2 \sin^2 \left(\frac{2}{3}\theta \right) - r^2 \left(3 - 4 \sin^2 \left(\frac{2}{3}\theta \right) \right)^2 \right\} \\
&\geq -\frac{2}{3}r \sin \left(\frac{2}{3}\theta \right) \left\{ 10 - 4r^2 \sin^2 \left(\frac{2}{3}\theta \right) \right\},
\end{aligned}$$

which implies

$$\begin{aligned}
& \left| \Delta \left\{ (1-x^2)(1-y^2)r^{2/3} \sin \left(\frac{2}{3}\theta \right) \right\} \right| \\
&\leq \frac{2}{3}r \sin \left(\frac{2}{3}\theta \right) \left\{ 10 - 4r^2 \sin^2 \left(\frac{2}{3}\theta \right) \right\} \\
&= \frac{20\sqrt{10}}{9\sqrt{3}} - \frac{8}{3} \left(r \sin \left(\frac{2}{3}\theta \right) - \sqrt{\frac{5}{6}} \right)^2 \left(r \sin \left(\frac{2}{3}\theta \right) + \sqrt{\frac{10}{3}} \right) \leq \frac{20\sqrt{10}}{9\sqrt{3}}.
\end{aligned}$$

Then

$$\begin{aligned}
& \left\| \Delta \left\{ (1-x^2)(1-y^2)r^{2/3} \sin \left(\frac{2}{3}\theta \right) \right\} \right\|_{L^2(\Omega_0)}^2 \\
&\leq \frac{4}{9} \int_0^1 \int_0^{3\pi/2} r^{4/3} \left\{ (4r^2 - 10) \sin \left(\frac{2}{3}\theta \right) + r^2 \sin \left(\frac{10}{3}\theta \right) \right\}^2 r d\theta dr \\
&\quad + \left(|\Omega_0| - \frac{3}{4}\pi \right) \left(\frac{20\sqrt{10}}{9\sqrt{3}} \right)^2 \\
&= \frac{127}{22}\pi + \left(3 - \frac{3}{4}\pi \right) \frac{4000}{243} = \frac{4000}{81} - \frac{11713}{1782}\pi.
\end{aligned}$$

□

Lemma 4.2 When $\Omega = \Omega_0$,

$$|\lambda| \leq \frac{1}{\pi} \left\| \left(r^{-2/3} - 2^{-2/3}r^{2/3} \right) \sin \left(\frac{2}{3}\theta \right) \right\|_{L^2(\Omega)} \|f(x, y)\|_{L^2(\Omega)}.$$

Proof. For any $0 < \varepsilon < 1$, let g_ε be a weak solution of the following equation,

$$(4.1) \quad \begin{cases} -\Delta g_\varepsilon = -\Delta \tilde{g}_\varepsilon & \text{in } \Omega, \\ g_\varepsilon = 0 & \text{on } \partial\Omega, \end{cases}$$

where,

$$\tilde{g}_\varepsilon(x, y) = \begin{cases} \frac{1}{\pi} \left((2\varepsilon^{-4/3} - 2^{-2/3})r^{2/3} - \varepsilon^{-8/3}r^2 \right) \sin\left(\frac{2}{3}\theta\right), & (r < \varepsilon), \\ \frac{1}{\pi} \left(r^{-2/3} - 2^{-2/3}r^{2/3} \right) \sin\left(\frac{2}{3}\theta\right), & (\varepsilon \leq r). \end{cases}$$

From the fact that $\tilde{g}_\varepsilon \in C^1(\Omega)$ and $-\Delta \tilde{g}_\varepsilon \geq 0$, both g_ε and $\tilde{g}_\varepsilon - g_\varepsilon$ are superharmonic function. Since g_ε and $\tilde{g}_\varepsilon - g_\varepsilon$ takes non-negative value on $\partial\Omega$,

$$0 \leq g_\varepsilon(x, y) \leq \tilde{g}_\varepsilon(x, y).$$

Consequently,

$$|g_\varepsilon(x, y)| \leq |\tilde{g}_\varepsilon(x, y)|.$$

From (1.1) and (4.1), taking g_ε and u as test functions,

$$\begin{aligned} \iint_{\Omega} f g_\varepsilon dx dy &= \iint_{\Omega} \nabla u \cdot \nabla g_\varepsilon dx dy = - \iint_{\Omega} u \Delta \tilde{g}_\varepsilon dx dy \\ &= \frac{32\varepsilon^{-8/3}}{9\pi} \iint_{\Omega} 1_{r < \varepsilon} \cdot u(x, y) \sin\left(\frac{2}{3}\theta\right) dx dy \\ &= \frac{32\varepsilon^{-8/3}\lambda}{9\pi} \int_0^\varepsilon \int_0^{3\pi/2} (1-x^2)(1-y^2)r^{2/3} \sin^2\left(\frac{2}{3}\theta\right) r d\theta dr \\ &\quad + \frac{32\varepsilon^{-8/3}}{9\pi} \int_0^\varepsilon \int_0^{3\pi/2} w(x, y) \sin\left(\frac{2}{3}\theta\right) r d\theta dr \\ &= \left(1 - \frac{4}{7}\varepsilon^2 + \frac{\varepsilon^4}{20}\right) \lambda \\ &\quad + \frac{16\varepsilon^{-8/3}}{9\pi} \int_0^\varepsilon \int_0^{3\pi/2} (\varepsilon^2 - r^2) \frac{\partial}{\partial r} w(x, y) \sin\left(\frac{2}{3}\theta\right) d\theta dr. \end{aligned}$$

Then

$$\begin{aligned}
& \left(1 - \frac{4}{7}\varepsilon^2 + \frac{\varepsilon^4}{20}\right) |\lambda| \\
& \leq \|f\|_{L^2(\Omega)} \|g_\varepsilon\|_{L^2(\Omega)} \\
& \quad + \frac{16\varepsilon^{-8/3}}{9\pi} \left(\int_0^\varepsilon \int_0^{3\pi/2} \left| (\varepsilon^2 - r^2) \sin\left(\frac{2}{3}\theta\right) \right|^{7/6} r^{-1/6} d\theta dr \right)^{6/7} \\
& \quad \times \left(\int_0^\varepsilon \int_0^{3\pi/2} \left| \frac{\partial}{\partial r} w(x, y) \right|^7 r d\theta dr \right)^{1/7} \\
& \leq \|f\|_{L^2(\Omega)} \|\tilde{g}_\varepsilon\|_{L^2(\Omega)} \\
& \quad + \frac{16\varepsilon^{1/21}}{9\pi} \left(\int_0^1 \int_0^{3\pi/2} \left| (1 - r^2) \sin\left(\frac{2}{3}\theta\right) \right|^{7/6} r^{-1/6} d\theta dr \right)^{6/7} \\
& \quad \times \left(\iint_\Omega \left(\left| \frac{\partial}{\partial x} w(x, y) \right| + \left| \frac{\partial}{\partial y} w(x, y) \right| \right)^7 dx dy \right)^{1/7}.
\end{aligned}$$

From $w \in H^2$ and the Sobolev embedding theorem [1],

$$\frac{\partial w}{\partial x} \in L^7(\Omega), \quad \frac{\partial w}{\partial y} \in L^7(\Omega).$$

Then, we have the conclusion when $\varepsilon \rightarrow 0$. □

Lemma 4.3

$$\left\| \left(r^{-2/3} - 2^{-2/3} r^{2/3} \right) \sin\left(\frac{2}{3}\theta\right) \right\|_{L^2(\Omega_0)} \leq \sqrt{\frac{3 \cdot 2^{1/3}}{5}} \pi.$$

Proof.

$$\begin{aligned}
& \left\| \left(r^{-2/3} - 2^{-2/3} r^{2/3} \right) \sin\left(\frac{2}{3}\theta\right) \right\|_{L^2(\Omega_0)}^2 \\
& \leq \int_0^{\sqrt{2}} \int_0^{3\pi/2} \left(r^{-2/3} - 2^{-2/3} r^{2/3} \right)^2 \sin^2\left(\frac{2}{3}\theta\right) r d\theta dr = \frac{3 \cdot 2^{1/3}}{5} \pi.
\end{aligned}$$
□

Lemma 4.4

$$\left\| \Delta \left\{ 1_{r < 1} \cdot (1 - r)^2 r^{2/3} \sin\left(\frac{2}{3}\theta\right) \right\} \right\|_{L^2(\Omega_0)} = \frac{3\sqrt{\pi}}{2}.$$

Proof. We have

$$\begin{aligned}
& \left\| \Delta \left\{ 1_{r<1} \cdot (1-r)^2 r^{2/3} \sin \left(\frac{2}{3} \theta \right) \right\} \right\|_{L^2(\Omega_0)}^2 \\
&= \left\| 1_{r<1} \cdot \frac{2}{3} (10r-7) r^{-1/3} \sin \left(\frac{2}{3} \theta \right) \right\|_{L^2(\Omega_0)}^2 \\
&= \frac{4}{9} \int_0^1 \int_0^{3\pi/2} (10r-7)^2 r^{-2/3} \sin^2 \left(\frac{2}{3} \theta \right) r d\theta dr = \frac{9}{4} \pi,
\end{aligned}$$

by calculating directly. □

Lemma 4.5

$$\begin{aligned}
& \left\| \Delta \left\{ \left((1-x^2)(1-y^2) - 1_{r<1} \cdot (1-r)^2 \right) r^{2/3} \sin \left(\frac{2}{3} \theta \right) \right\} \right\|_{L^2(\Omega_0)} \\
& \leq \sqrt{\frac{4000}{81} - \frac{232367}{46332}} \pi.
\end{aligned}$$

Proof.

$$\begin{aligned}
& \Delta \left\{ \left((1-x^2)(1-y^2) - (1-r)^2 \right) r^{2/3} \sin \left(\frac{2}{3} \theta \right) \right\} \\
&= \frac{2}{3} r^{2/3} \left\{ \left(4r^2 - 20 + \frac{7}{r} \right) \sin \left(\frac{2}{3} \theta \right) + r^2 \sin \left(\frac{10}{3} \theta \right) \right\}
\end{aligned}$$

When $r \geq 1$, same as proof of Lemma 4.1,

$$\left| \Delta \left\{ (1-x^2)(1-y^2) r^{2/3} \sin \left(\frac{2}{3} \theta \right) \right\} \right| \leq \frac{20\sqrt{10}}{9\sqrt{3}},$$

Then

$$\begin{aligned}
& \left\| \Delta \left\{ \left((1-x^2)(1-y^2) - 1_{r<1} \cdot (1-r)^2 \right) r^{2/3} \sin \left(\frac{2}{3} \theta \right) \right\} \right\|_{L^2(\Omega_0)}^2 \\
& \leq \frac{4}{9} \int_0^1 \int_0^{3\pi/2} r^{4/3} \left\{ \left(4r^2 - 20 + \frac{7}{r} \right) \sin \left(\frac{2}{3} \theta \right) + r^2 \sin \left(\frac{10}{3} \theta \right) \right\}^2 r d\theta dr \\
& \quad + \left(|\Omega_0| - \frac{3}{4} \pi \right) \left(\frac{20\sqrt{10}}{9\sqrt{3}} \right)^2 \\
&= \frac{4193}{572} \pi + \left(3 - \frac{3}{4} \pi \right) \frac{4000}{243} = \frac{4000}{81} - \frac{232367}{46332} \pi.
\end{aligned}$$

□

Lemma 4.6

$$|\lambda_k| \leq \left\{ \int \int_{\Omega} 1_{0 \leq \theta < 3\pi/2} \cdot \left| G_{l_k} \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right) \right|^2 dx dy \right. \\ \left. + \int \int_{\Omega} 1_{3\pi/2 \leq \theta < 2\pi} \cdot \left| G_{l_k} \left(r, \frac{1}{9} (8 + \cos(4\theta)), 1 \right) \right|^2 dx dy \right\}^{1/2} \|f(x, y)\|_{L^2(\Omega)}.$$

Proof. Using the parallel translation and the rotation, we move re-entrant corner to the origin and re-entrant angle to $[0, 3\pi/2]$.

For $\varepsilon < l_k$, let g_ε be a weak solution of following equation,

$$\begin{cases} -\Delta g_\varepsilon = -1_{0 < \theta < 3\pi/2} \cdot 1_{r < \varepsilon} \cdot \Delta \tilde{g}_\varepsilon & \text{in } \Omega, \\ g_\varepsilon = 0, & \text{on } \partial\Omega, \end{cases}$$

where,

$$\tilde{g}_\varepsilon(x, y) = \begin{cases} \left(2\varepsilon^{-4/3} r^{4/3} - \varepsilon^{-8/3} r^{8/3} \right) G_{l_k} \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right), & (r < \varepsilon, 0 \leq \theta \leq 3\pi/2), \\ G_{l_k} \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right), & (r \geq \varepsilon, 0 \leq \theta \leq 3\pi/2), \\ G_{l_k} \left(r, \frac{1}{9} (8 + \cos(4\theta)), 1 \right), & (3\pi/2 < \theta < 2\pi). \end{cases}$$

From the fact that $\tilde{g}_\varepsilon \in C^1(\Omega)$ and Lemma 4.8, 4.9 and 4.10, both g_ε and $\tilde{g}_\varepsilon - g_\varepsilon$ are superharmonic function. Since g_ε and $\tilde{g}_\varepsilon - g_\varepsilon$ takes non-negative value on $\partial\Omega$,

$$0 \leq g_\varepsilon(x, y) \leq \tilde{g}_\varepsilon(x, y).$$

Consequently,

$$|g_\varepsilon(x, y)| \leq |\tilde{g}_\varepsilon(x, y)|.$$

Then, same as in Lemma 4.2,

$$\begin{aligned} \int \int_{\Omega} f g_\varepsilon dx dy &= \int \int_{\Omega} \nabla u \cdot \nabla g_\varepsilon dx dy = - \int \int_{\Omega} 1_{0 < \theta < 3\pi/2} \cdot 1_{r < \varepsilon} \cdot u \Delta \tilde{g}_\varepsilon dx dy \\ &= - \int \int_{\Omega} 1_{0 < \theta < 3\pi/2} \cdot 1_{r < \varepsilon} \cdot u(x, y) \\ &\quad \times \Delta \left\{ \left(2\varepsilon^{-4/3} r^{4/3} - \varepsilon^{-8/3} r^{8/3} \right) G_{l_k} \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right) \right\} dx dy. \end{aligned}$$

From Lemma 4.11,

$$\begin{aligned}
& \iint_{\Omega} f g_{\varepsilon} dx dy \\
&= \frac{32\varepsilon^{-8/3} l_k^{2/3}}{9\pi} (1 + O(\varepsilon^{4/3})) \iint_{\Omega} 1_{r < \varepsilon} \cdot u(x, y) \sin\left(\frac{2}{3}\theta\right) dx dy \\
&= \frac{32\varepsilon^{-8/3} l_k^{2/3} \lambda_k}{9\pi} (1 + O(\varepsilon^{4/3})) \\
&\quad \times \int_0^{\varepsilon} \int_0^{3\pi/2} \left(1 - \frac{x^2}{l_k^2}\right) \left(1 - \frac{y^2}{l_k^2}\right) \left(\frac{r}{l_k}\right)^{2/3} \sin^2\left(\frac{2}{3}\theta\right) r d\theta dr \\
&\quad + \frac{32\varepsilon^{-8/3} l_k^{2/3}}{9\pi} (1 + O(\varepsilon^{4/3})) \int_0^{\varepsilon} \int_0^{3\pi/2} w(x, y) \sin\left(\frac{2}{3}\theta\right) r d\theta dr \\
&= (1 + O(\varepsilon^{4/3})) \lambda_k \\
&\quad + \frac{32\varepsilon^{-8/3} l_k^{2/3}}{9\pi} (1 + O(\varepsilon^{4/3})) \int_0^{\varepsilon} \int_0^{3\pi/2} w(x, y) \sin\left(\frac{2}{3}\theta\right) r d\theta dr.
\end{aligned}$$

Then, same as in Lemma 4.2,

$$\begin{aligned}
(1 + O(\varepsilon^{4/3})) |\lambda_k| &\leq \|f\|_{L^2(\Omega)} \|g_{\varepsilon}\|_{L^2(\Omega)} + O(\varepsilon^{1/21}) \\
&\leq \|f\|_{L^2(\Omega)} \|\tilde{g}_{\varepsilon}\|_{L^2(\Omega)} + O(\varepsilon^{1/21}).
\end{aligned}$$

We have the conclusion when $\varepsilon \rightarrow 0$. □

Lemma 4.7

$$\begin{aligned}
& \left\{ \iint_{\Omega} 1_{0 \leq \theta < 3\pi/2} \cdot \left| G_l \left(r, \cos\left(\frac{4}{3}\theta\right), \cos\left(\frac{4}{3}\theta\right) \right) \right|^2 dx dy \right. \\
& \quad \left. + \iint_{\Omega} 1_{3\pi/2 \leq \theta < 2\pi} \cdot \left| G_l \left(r, \frac{1}{9}(8 + \cos(4\theta)), 1 \right) \right|^2 dx dy \right\}^{1/2} \\
& \leq \frac{1}{\pi} \sqrt{\left(\frac{5}{2} - \frac{3\pi}{8}\right) l^2 + 2|\Omega|}
\end{aligned}$$

Proof.

$$\begin{aligned}
& \left| G_l \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right) \right|^2 \\
&= \frac{l^{4/3}}{2\pi^2} \left\{ \sqrt{r^{-8/3} + l^{-8/3} - 2r^{-4/3}l^{-4/3} \cos \left(\frac{4}{3} \theta \right)} \right. \\
&\quad \left. + l^{-4/3} - r^{-4/3} \cos \left(\frac{4}{3} \theta \right) \right\} \\
&\leq \frac{l^{4/3}}{2\pi^2} \left\{ |l^{-4/3} - r^{-4/3}| + \sqrt{2r^{-4/3}l^{-4/3} \left(1 - \cos \left(\frac{4}{3} \theta \right) \right)} \right. \\
&\quad \left. + l^{-4/3} - r^{-4/3} \cos \left(\frac{4}{3} \theta \right) \right\} \\
&= \frac{l^{4/3}}{\pi^2} \left\{ \max(l^{-4/3} - r^{-4/3}, 0) \right. \\
&\quad \left. + r^{-2/3}l^{-2/3} \sin \left(\frac{2}{3} \theta \right) + r^{-4/3} \sin^2 \left(\frac{2}{3} \theta \right) \right\},
\end{aligned}$$

$$\begin{aligned}
& \left| G_l \left(r, \frac{1}{9} (8 + \cos(4\theta)), 1 \right) \right|^2 \\
&= \frac{l^{4/3}}{2\pi^2} \left\{ \sqrt{r^{-8/3} + l^{-8/3} - \frac{2}{9} r^{-4/3} l^{-4/3} (8 + \cos(4\theta))} + l^{-4/3} - r^{-4/3} \right\} \\
&\leq \frac{l^{4/3}}{2\pi^2} \left\{ |l^{-4/3} - r^{-4/3}| + \sqrt{\frac{2}{9} r^{-4/3} l^{-4/3} (1 - \cos(4\theta))} + l^{-4/3} - r^{-4/3} \right\} \\
&= \frac{l^{4/3}}{\pi^2} \left\{ \max(l^{-4/3} - r^{-4/3}, 0) + \frac{1}{3} r^{-2/3} l^{-2/3} |\sin(2\theta)| \right\}.
\end{aligned}$$

It follows from these inequalities that,

$$\begin{aligned}
& \iint_{\Omega} 1_{0 \leq \theta < 3\pi/2} \cdot \left| G_l \left(r, \cos \left(\frac{4}{3} \theta \right), \cos \left(\frac{4}{3} \theta \right) \right) \right|^2 dx dy \\
&\quad + \iint_{\Omega} 1_{3\pi/2 \leq \theta < 2\pi} \cdot \left| G_l \left(r, \frac{1}{9} (8 + \cos(4\theta)), 1 \right) \right|^2 dx dy \\
&\leq \frac{l^{4/3}}{\pi^2} \int_0^l \int_0^{3\pi/2} \left\{ r^{-2/3} l^{-2/3} \sin \left(\frac{2}{3} \theta \right) + r^{-4/3} \sin^2 \left(\frac{2}{3} \theta \right) \right\} r d\theta dr \\
&\quad + \frac{l^{4/3}}{\pi^2} \int_0^l \int_{3\pi/2}^{2\pi} \frac{1}{3} r^{-2/3} l^{-2/3} |\sin(2\theta)| r d\theta dr \\
&\quad + \left(|\Omega| - \frac{3\pi l^2}{4} \right) \frac{l^{4/3}}{\pi^2} \cdot 2l^{-4/3}
\end{aligned}$$

$$= \frac{l^2}{\pi^2} \left(\frac{5}{2} - \frac{3\pi}{8} \right) + \frac{2}{\pi^2} |\Omega|.$$

□

Lemma 4.8

$$-\Delta G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right) = 0 \quad \text{in } \Omega.$$

Proof.

$$G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right)$$

is a constant times imaginary part of

$$\left(z^{-4/3} - l^{-4/3} \right)^{1/2}, \quad z = re^{i\theta}.$$

Therefore, this function is harmonic in Ω .

□

Lemma 4.9 For $r < \varepsilon < l$ and $0 < \theta < 3\pi/2$,

$$-\Delta \left(\left(2\varepsilon^{-4/3}r^{4/3} - \varepsilon^{-8/3}r^{8/3} \right) G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right) \right) \geq 0 \quad \text{in } \Omega.$$

Proof.

$$\begin{aligned} & -\Delta \left\{ \left(2\varepsilon^{-4/3}r^{4/3} - \varepsilon^{-8/3}r^{8/3} \right) G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right) \right\} \\ &= \frac{4\sqrt{2}}{9a^2b^2cl^2\pi} \cdot \frac{(a+c-1)^2(a+c+1)^2(a-b+c)}{\left(a+c-\cos \left(\frac{4}{3}\theta \right) \right)^{3/2}} \end{aligned}$$

where

$$a = r^{4/3}l^{-4/3}, \quad b = \varepsilon^{4/3}l^{-4/3}, \quad c = \sqrt{a^2 + 1 - 2a \cos \left(\frac{4}{3}\theta \right)}.$$

Since

$$a - b + c = a - b + \sqrt{a^2 + 1 - 2a \cos \left(\frac{4}{3}\theta \right)} \geq a - b + |a - 1| \geq 1 - b \geq 0,$$

this lemma is proved.

□

Lemma 4.10 For $3\pi/2 < \theta < 2\pi$,

$$-\Delta G_l \left(r, \frac{1}{9} \left(8 + \cos(4\theta) \right), 1 \right) \geq 0 \quad \text{in } \Omega.$$

Proof.

$$\begin{aligned} -\Delta G_l \left(r, \frac{1}{9} (8 + \cos(4\theta)), 1 \right) \\ = \frac{2\sqrt{2}\sqrt{a+d-1}}{9a^2d^3l^2\pi} (2d^3 + 2(a-1)^2d - 4(a-1)^3 + d^2) \end{aligned}$$

where

$$a = r^{4/3}l^{-4/3}, \quad d = \sqrt{a^2 + 1 - \frac{2}{9}a(8 + \cos(4t))}.$$

Since

$$d \geq \sqrt{a^2 + 1 - 2a} = |a - 1|$$

holds,

$$2d^3 + 2(a-1)^2d - 4(a-1)^3 + d^2 \geq 4|a-1|^3 - 4(a-1)^3 + 2(a-1)^2d + d^2 \geq 0.$$

Then, this lemma is proved. \square

Lemma 4.11 When $r < \varepsilon$,

$$\begin{aligned} -\Delta \left\{ \left(2\varepsilon^{-4/3}r^{4/3} - \varepsilon^{-8/3}r^{8/3} \right) G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right) \right\} \\ = \frac{32\varepsilon^{-8/3}l^{2/3}}{9\pi} \sin \left(\frac{2}{3}\theta \right) \cdot (1 + O(\varepsilon^{4/3})). \end{aligned}$$

Proof. Define

$$a = r^{4/3}l^{-4/3}, \quad b = \varepsilon^{4/3}l^{-4/3}, \quad c = \sqrt{a^2 + 1 - 2a \cos \left(\frac{4}{3}\theta \right)},$$

then we have,

$$\begin{aligned} -\Delta \left\{ \left(2\varepsilon^{-4/3}r^{4/3} - \varepsilon^{-8/3}r^{8/3} \right) G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right) \right\} \\ = \frac{4\sqrt{2}}{9a^2b^2cl^2\pi} \cdot \frac{(a+c-1)^2(a+c+1)^2(a-b+c)}{\left(a+c-\cos \left(\frac{4}{3}\theta \right) \right)^{3/2}}. \end{aligned}$$

We can easily confirm the following expressions

$$\begin{aligned} c &= 1 + O(\varepsilon^{4/3}), \\ a+c-1 &= \frac{2a}{c-a+1} \left(1 - \cos \left(\frac{4}{3}\theta \right) \right) = 2a \sin^2 \left(\frac{2}{3}\theta \right) \cdot (1 + O(\varepsilon^{4/3})), \\ a+c+1 &= 2 + O(\varepsilon^{4/3}), \\ a-b+c &= 1 + O(\varepsilon^{4/3}), \\ a+c-\cos \left(\frac{4}{3}\theta \right) &= \frac{c+a+1}{c-a+1} \left(1 - \cos \left(\frac{4}{3}\theta \right) \right) \\ &= 2 \sin^2 \left(\frac{2}{3}\theta \right) \cdot (1 + O(\varepsilon^{4/3})). \end{aligned}$$

Then,

$$\begin{aligned} & -\Delta \left\{ \left(2\varepsilon^{-4/3}r^{4/3} - \varepsilon^{-8/3}r^{8/3} \right) G_l \left(r, \cos \left(\frac{4}{3}\theta \right), \cos \left(\frac{4}{3}\theta \right) \right) \right\} \\ & = \frac{32\varepsilon^{-8/3}l^{2/3}}{9\pi} \sin \left(\frac{2}{3}\theta \right) \cdot (1 + O(\varepsilon^{4/3})). \end{aligned}$$

□

5 Numerical result

In this section, numerical results are shown to confirm the validity of the error estimation. All calculations were carried out on a Pentium IV PC at 2.2GHz with Borland C++ compiler. There are some difficulties in calculating H_0^1 inner product between singular basis and bilinear basis because gradient of singular basis diverges at the origin. To deal with this difficulty, the following Double Exponential transformation (DE transformation)[14] is used to calculate the integral at each of the square elements,

$$\begin{aligned} & \int_{y_l}^{y_l+h} \int_{x_k}^{x_k+h} p(x, y) dx dy \\ (5.1) \quad & = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(x_k + \varphi(x), y_l + \varphi(y)) \varphi'(x) \varphi'(y) dx dy, \end{aligned}$$

where $p(x, y)$ is an integrand and

$$\varphi(t) = \frac{h}{2} \left(\tanh \left(\frac{\pi}{2} \sinh t \right) + 1 \right).$$

(5.1) is approximated by the trapezoidal rule as follows,

$$\begin{aligned} & \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} p(x_k + \varphi(x), y_l + \varphi(y)) \varphi'(x) \varphi'(y) dx dy \\ & \approx \frac{L^2}{N^2} \sum_{j=-N}^N \sum_{i=-N}^N \left(x_k + \varphi \left(\frac{kL}{N} \right), y_l + \varphi \left(\frac{kL}{N} \right) \right) \varphi' \left(\frac{kL}{N} \right) \varphi' \left(\frac{kL}{N} \right). \end{aligned}$$

We took $L = 4$ and $N = 100$ which is sufficient to obtain double floating point precision.

We consider following Poisson equation on two different shapes of Ω ,

$$(5.2) \quad \begin{cases} -\Delta u = 1 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

The first result is that Ω is L-shape domain which is shown in Fig.8. Fig.9 shows an arrangement of the singular bases and Fig.10 shows the shape of the numerical solution. The numerical results are presented in Table 5.1, where h is the mesh size and U_h denotes the numerical solution.




Figure 8: L-Shape domain




Figure 9: An arrangement of the singular bases




Figure 10: The numerical solution when $h = 1/10$

Table 5.1: Numerical results for the L-shape domain

h	$\ U_h - U_{h/2}\ _{H_0^1(\Omega)}$	$\ U_h - U_{h/2}\ _{L^2(\Omega)}$	Degree of freedom	Condition number
1/10	3.9052×10^{-2}	1.0684×10^{-3}	443	3.8210×10^3
1/20	1.9576×10^{-2}	2.6905×10^{-4}	1882	4.9653×10^4
1/30	1.3066×10^{-2}	1.2008×10^{-4}	4322	2.2077×10^5
1/40	9.8064×10^{-3}	6.7750×10^{-5}	7762	6.3104×10^5
1/50	7.8494×10^{-3}	4.3464×10^{-5}	12202	1.4173×10^6

Table 5.2: A priori error estimation for the L-shape domain

h	$\ u - u_h\ _{H_0^1(\Omega)}$	$\ u - u_h\ _{L^2(\Omega)}$
1/10	$7.161561 \dots \times 10^{-1}$	$2.293667 \dots \times 10^{-1}$
1/20	$3.580780 \dots \times 10^{-1}$	$5.734168 \dots \times 10^{-2}$
1/30	$2.387187 \dots \times 10^{-1}$	$2.548519 \dots \times 10^{-2}$
1/40	$1.790390 \dots \times 10^{-1}$	$1.433542 \dots \times 10^{-2}$
1/50	$1.432312 \dots \times 10^{-1}$	$9.174669 \dots \times 10^{-3}$

Table 5.3: Numerical results for the H-shape domain

h	$\ U_h - U_{h/2}\ _{H_0^1(\Omega)}$	$\ U_h - U_{h/2}\ _{L^2(\Omega)}$	Degree of freedom	Condition number
1/10	6.5080×10^{-2}	1.9355×10^{-3}	715	3.4622×10^2
1/20	3.4414×10^{-2}	5.4674×10^{-4}	3025	3.9510×10^3
1/30	2.3745×10^{-2}	2.6310×10^{-4}	6935	1.7397×10^4
1/40	1.8253×10^{-2}	1.5691×10^{-4}	12445	5.0101×10^4
1/50	1.4882×10^{-2}	1.0512×10^{-4}	19555	1.1370×10^5

Since the exact solution of (5.2) is not known, we calculated H_0^1 and L^2 norm of $U_h - U_{h/2}$ to obtain the rough estimation of the error. The true errors are expected to be several times larger than these value. To calculate error norm, we employed DE transformation again. The condition numbers of the element matrix were also calculated. The definition of the condition number is the ratio of the largest to the smallest eigenvalue.

From Theorem 3.1, H_0^1 and L^2 error estimations for this equation are obtained as follows,

$$\begin{aligned} \|u - u_h\|_{H_0^1(\Omega)} &\leq \left(0.319 + \sqrt{0.971 + 1.469 \cdot \frac{5}{1^2}} \right) \sqrt{5}h, \\ \|u - u_h\|_{L^2(\Omega)} &\leq \left(0.319 + \sqrt{0.971 + 1.469 \cdot \frac{5}{1^2}} \right)^2 \sqrt{5}h^2. \end{aligned}$$

The right-hand side of these inequalities are presented in Table 5.2. Since a priori error estimate can be applied to arbitrary f and Ω , in general, the actual error is often smaller than a priori error estimation.

The next result is a case that Ω is an H-shape domain which is shown in Fig.11. Fig.12 shows an arrangement of the singular bases and Fig.13 shows the shape of the numerical solution. The numerical results are presented in Table 5.3.




Figure 11: H-Shape domain




Figure 12: An arrangement of the singular bases

Table 5.4: A priori error estimation for the H-shape domain

h	$\ u - u_h\ _{H_0^1(\Omega)}$	$\ u - u_h\ _{L^2(\Omega)}$
1/10	7.926879...	22.21567...
1/20	3.963439...	5.553918...
1/30	2.642293...	2.468408...
1/40	1.981719...	1.388479...
1/50	1.585375...	0.888626...

From Theorem 3.1, H_0^1 and L^2 error estimation for this equation are obtained as follows,

$$\|u - u_h\|_{H_0^1(\Omega)} \leq \left(0.319 + 4\sqrt{0.971 + 1.469 \cdot \frac{8}{0.5^2}} \right) \sqrt{8}h,$$

$$\|u - u_h\|_{L^2(\Omega)} \leq \left(0.319 + 4\sqrt{0.971 + 1.469 \cdot \frac{8}{0.5^2}} \right)^2 \sqrt{8}h^2.$$

The right-hand side of these inequalities are presented in Table 5.4.

As we can see in Table 5.3 and Table 5.4, the value of a priori error estimation becomes larger when the number of the re-entrant corner increases and when the support of the singular function become smaller (n and l_k in Theorem 3.1).

6 Concluding remark

We presented a constructive a priori error estimation for finite element solution in a polygonal domain by using singular functions. The results are only when the domain is bounded by line segments which forms a right angle. However, it seems to be possible to extend this method to the general polygonal domain with triangular mesh. A priori error estimation for another important method such as DSFM or method of mesh refinement is further work.




Figure 13: The numerical solution when $h = 1/10$

Acknowledgment

The author is deeply grateful to Professor M. T. Nakao of Kyushu University for helpful discussions and suggestions. The author also wishes to thank Professor K. Ohtsuka of Hiroshima Kokusai Gakuin University and Professor M. Plum of Karlsruhe University for useful advice to improve the paper.

Reference

- [1] Adams R. A.: Sobolev Spaces. Academic Press, New York (1975).
- [2] Babuska, I., Kellog, R. B., and Pitkaranta, J.: Direct and inverse error estimates for finite elements with mesh refinement. *Numer. Math.* **33**, 447–471 (1979).
- [3] Blum, H., Dobrowolski, M.: On finite element methods for elliptic equations on domains with corners. *Computing* **28**, 53–63 (1982).
- [4] Brenner, S. C.: Multigrid methods for the computation of singular solutions and stress intensity factors. I: corner singularities. *Math. Comp.* **68**, 559–583 (1999).
- [5] Cai, Z., Kim, S.: A finite element method using singular function for the Poisson equation :corner singularities. *SIAM J. Numer. Anal.* **39**, 286–299 (2001).
- [6] Ciarlet, P. G.: The Finite Element Method for Elliptic Problems. North-Holland, Amsterdam (1978).
- [7] Dauge, M., Nicaise, S., Bourland, M., Lubuma, M. S.: Coefficients of the singularities for elliptic boundary value problems on domains with conical points. III: Finite element methods on polygonal domains. *SIAM J. Numer. Anal.* **29**, 136–155 (1992).
- [8] Fix, G., Gulati S., Wakoff G.I.: On the use of singular functions with the finite element method. *J. Comp. Phys.* **13**, 209–228 (1973).
- [9] Grisvard, P.: Elliptic Problems in Nonsmooth Domains. Pitman Publishing, Boston (1985).
- [10] Grisvard, P.: Singularities in Boundary Value Problems. *RMA* **22**, Masson, Paris (1992).
- [11] Ladyzhenskaya, OA.: The Mathematical Theory of Viscous Incompressible Flows. Gordon and Breach, (1969).
- [12] Nakao, M.T.: A numerical verification method for the existence of weak solutions for nonlinear boundary value problems. *J. Math. Anal. Appl.* **164**, 489–507 (1992).

- [13] Schultz, M.: Spline Analysis. Prentice-Hall, (1973).
- [14] Takahasi, H., Mori, M.: Double exponential formulas for numerical integration. Publ. Res. Inst. Math. Soc. **9**, 721–741 (1974).
- [15] Yamamoto, N., Nakao, M.T.: Numerical verifications of solutions for elliptic equations in nonconvex polygonal domains. Numer. Math. **65**, 503–521 (1993).
- [16] Watanabe, Y., Nakao, M.T.: Numerical verifications of solutions for nonlinear elliptic equations. Japan J. Indust. Appl. Math. **10**, 165–178 (1993).

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q -Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift
- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations

- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA
Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS
Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI
Nonlinear regression modeling via regularized wavelets and smoothing parameter selection
- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO, Yoshihiro MIZOGUCHI & Yasuo KAWAHARA
On reversible cellular automata with finite cell array

- MHF2005-28 Toru KOMATSU
Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU
A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO
Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI
A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI
A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI
An explicit formula of the Shapley value for a cooperative game induced from the conjugate point
- MHF2005-34 Hideki MURAKAWA
A regularization of a reaction-diffusion system approximation to the two-phase Stefan problem
- MHF2006-1 Masahisa TABATA
Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme
- MHF2006-2 Ken-ichi MARUNO & G R W QUISPEL
Construction of integrals of higher-order mappings
- MHF2006-3 Setsuo TANIGUCHI
On the Jacobi field approach to stochastic oscillatory integrals with quadratic phase function
- MHF2006-4 Kouji HASHIMOTO, Kaori NAGATOU & Mitsuhiro T. NAKAO
A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains
- MHF2006-5 Hidefumi KAWASAKI
A duality theory based on triangular cylinders separating three convex sets in R^n
- MHF2006-6 Raimundas VIDŪNAS
Uniform convergence of hypergeometric series
- MHF2006-7 Yuji KODAMA & Ken-ichi MARUNO
N-Soliton solutions to the DKP equation and Weyl group actions

- MHF2006-8 Toru KOMATSU
Potentially generic polynomial
- MHF2006-9 Toru KOMATSU
Generic sextic polynomial related to the subfield problem of a cubic polynomial
- MHF2006-10 Shu TEZUKA & Anargyros PAPAGEORGIOU
Exact cubature for a class of functions of maximum effective dimension
- MHF2006-11 Shu TEZUKA
On high-discrepancy sequences
- MHF2006-12 Raimundas VIDŪNAS
Detecting persistent regimes in the North Atlantic Oscillation time series
- MHF2006-13 Toru KOMATSU
Tamely Eisenstein field with prime power discriminant
- MHF2006-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the Hankel determinant formula for the solutions of the Painlevé IV equation
- MHF2006-15 Raimundas VIDŪNAS
Darboux evaluations of algebraic Gauss hypergeometric functions
- MHF2006-16 Masato KIMURA & Isao WAKANO
New mathematical approach to the energy release rate in crack extension
- MHF2006-17 Toru KOMATSU
Arithmetic of the splitting field of Alexander polynomial
- MHF2006-18 Hiroki MASUDA
Likelihood estimation of stable Lévy processes from discrete data
- MHF2006-19 Hiroshi KAWABI & Michael RÖCKNER
Essential self-adjointness of Dirichlet operators on a path space with Gibbs measures via an SPDE approach
- MHF2006-20 Masahisa TABATA
Energy stable finite element schemes and their applications to two-fluid flow problems
- MHF2006-21 Yuzuru INAHAMA & Hiroshi KAWABI
Asymptotic expansions for the Laplace approximations for Itô functionals of Brownian rough paths
- MHF2006-22 Yoshiyuki KAGEI
Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

- MHF2006-23 Yoshiyuki KAGEI
Asymptotic behavior of the semigroup associated with the linearized compressible Navier-Stokes equation in an infinite layer
- MHF2006-24 Akihiro MIKODA, Shuichi INOKUCHI, Yoshihiro MIZOGUCHI & Mitsuhiro FUJIO
The number of orbits of box-ball systems
- MHF2006-25 Toru FUJII & Sadanori KONISHI
Multi-class logistic discrimination via wavelet-based functionalization and model selection criteria
- MHF2006-26 Taro HAMAMOTO, Kenji KAJIWARA & Nicholas S. WITTE
Hypergeometric solutions to the q -Painlevé equation of type $(A_1 + A'_1)^{(1)}$
- MHF2006-27 Hiroshi KAWABI & Tomohiro MIYOKAWA
The Littlewood-Paley-Stein inequality for diffusion processes on general metric spaces
- MHF2006-28 Hiroki MASUDA
Notes on estimating inverse-Gaussian and gamma subordinators under high-frequency sampling
- MHF2006-29 Setsuo TANIGUCHI
The heat semigroup and kernel associated with certain non-commutative harmonic oscillators
- MHF2006-30 Setsuo TANIGUCHI
Stochastic analysis and the KdV equation
- MHF2006-31 Masato KIMURA, Hideki KOMURA, Masayasu MIMURA, Hidenori MIYOSHI, Takeshi TAKAISHI & Daishin UYEYAMA
Quantitative study of adaptive mesh FEM with localization index of pattern
- MHF2007-1 Taro HAMAMOTO & Kenji KAJIWARA
Hypergeometric solutions to the q -Painlevé equation of type $A_4^{(1)}$
- MHF2007-2 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Verified numerical computation of solutions for the stationary Navier-Stokes equation in nonconvex polygonal domains
- MHF2007-3 Kenji KAJIWARA, Marta MAZZOCCO & Yasuhiro OHTA
A remark on the Hankel determinant formula for solutions of the Toda equation
- MHF2007-4 Jun-ichi SATO & Hidefumi KAWASAKI
Discrete fixed point theorems and their application to Nash equilibrium
- MHF2007-5 Mitsuhiro T. NAKAO & Kouji HASHIMOTO
Constructive error estimates of finite element approximations for non-coercive elliptic problems and its applications

MHF2007-6 Kouji HASHIMOTO

A preconditioned method for saddle point problems

MHF2007-7 Christopher MALON, Seiichi UCHIDA & Masakazu SUZUKI

Mathematical symbol recognition with support vector machines

MHF2007-8 Kenta KOBAYASHI

On the global uniqueness of Stokes' wave of extreme form

MHF2007-9 Kenta KOBAYASHI

A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions