# A numerical verification method of bifurcating solutions for 3-dimensional Rayleigh-B´enard problems

Kim, Myoungnyoun Faculty of Mathematics, Kyushu University

Nakao, Mitsuhiro T. Faculty of Mathematics, Kyushu University

Watanabe, Yoshitaka Computing and Communications Center, Kyushu University

Nishida, Takaaki Faculty of Science and Engineering, Waseda University

https://hdl.handle.net/2324/4024

出版情報:2007-02-20. 九州大学大学院数理学研究院 バージョン: 権利関係:

## **MHF** Preprint Series

Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

A numerical verification method of bifurcating solutions for 3-dimensional Rayleigh-Bénard problems

# M. Kim, M. T. Nakao Y. Watanabe, T. Nishida

### MHF 2007-10

(Received February 20, 2007)

Faculty of Mathematics Kyushu University Fukuoka, JAPAN Myoungnyoun Kim · Mitsuhiro T. Nakao · Yoshitaka Watanabe · Takaaki Nishida

# A numerical verification method of bifurcating solutions for 3-dimensional Rayleigh-Bénard problems

**Abstract** This paper is the three dimensional extension of the two dimensional work [4] and [7] on a computer assisted proof of the existence of nontrivial steady state solutions for Rayleigh–Bénard convection based on the fixed point theorem using a Newton like operator. The differences are emerging of complicated types of bifurcation, direct attack on the problem without stream functions, and increased complexity of numerical computation. The last one makes it hard to proceed the verification of solutions corresponding to the points on bifurcation diagram for three dimensional case. Actually, this work should be the first result for the three dimensional Navier-Stokes problems which seems to be very difficult to solve by theoretical approaches.

#### **1** Introduction

The Rayleigh–Bénard convection describes the instability of fluid between two infinite solid plates with hot bottom and cool top. The motion of fluid is self-sustained as soon as gravitational energy release overcomes dissipation losses, which is called Rayleigh's mechanism by buoyancy (for detailed mechanism, see [1] and [3]). We will use the Oberbeck–Boussinesq equations as approximate equations for this convection problem after normalization of variables and parameters:

$$\frac{1}{\mathscr{P}}\left[\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla)\mathbf{u}\right] + \nabla p = \Delta \mathbf{u} - (\mathscr{G} - \mathscr{R}T)\mathbf{e}_z, \tag{1a}$$

$$\nabla \cdot \mathbf{u} = 0, \tag{1b}$$

$$\frac{\partial T}{\partial t} + (\mathbf{u} \cdot \nabla)T = \Delta T, \qquad (1c)$$

where  $\mathbf{u} = (u, v, w)$  is the velocity field, *p* the pressure, *T* the temperature,  $\mathscr{R}$  Rayleigh number,  $\mathscr{P}$  Prandtl number, and a parameter  $\mathscr{G}$  containing gravity factors. We use stress free boundary conditions on the velocity field and Dirichlet boundary conditions on the temperature ( $T_{z=0} = 0, T_{z=\pi} = \pi$ ). Under a reference pressure  $p_a$ , the equilibrium state comes from the pure heat conduction:

~

$$\mathbf{u} = \mathbf{0}, \quad T = \pi - z, \quad p = \mathscr{G}(\pi - z) - \frac{1}{2}\mathscr{R}(\pi - z)^2 + p_a.$$
 (2)

Myoungnyoun Kim

Mitsuhiro T. Nakao

Yoshitaka Watanabe

Computing and Communications Center, Kyushu University 33, Fukuoka 812-8581, Japan E-mail: watanabe@cc.kyushu-u.ac.jp

Takaaki Nishida

Faculty of Science and Engineering, Waseda University, 3-4-1 Okubo, Shinjuku-ku, Tokyo 169-8555, Japan E-mail: tkknish@waseda.jp

Faculty of Mathematics, Kyushu University 33, Fukuoka 812-8581, Japan E-mail: mnkim@math.kyushu-u.ac.jp

Faculty of Mathematics, Kyushu University 33, Fukuoka 812-8581, Japan E-mail: mtnakao@math.kyushu-u.ac.jp

Using the conduction solution (2) and eliminating time derivatives from (1), we obtain the steady state bifurcation equations for the perturbation  $(\mathbf{u}, \theta, p)$  to the equilibrium:

$$-\Delta \mathbf{u} + \frac{1}{\mathscr{P}} (\mathbf{u} \cdot \nabla) \mathbf{u} + \nabla p - \mathscr{R} \boldsymbol{\theta} \mathbf{e}_z = \mathbf{0}, \tag{3a}$$

$$\nabla \cdot \mathbf{u} = 0, \tag{3b}$$

$$-\Delta \theta + (\mathbf{u} \cdot \nabla) \theta - w = 0. \tag{3c}$$

Given positive wave numbers  $a, b \le 1$ , we assume that all fluid motions are essentially confined to

$$\Omega = \left\{ (x, y, z) \in \mathbb{R}^3 : 0 \le x \le \frac{2\pi}{a}, 0 \le y \le \frac{2\pi}{b}, 0 \le z \le \pi \right\}, \quad |\Omega| = \frac{4\pi^3}{ab},$$

and impose parity conditions on new boundaries as in [5] together with periodic boundary conditions in horizontal directions [4]. From these boundary conditions, the velocity field, the perturbations of temperature and pressure can be represented by the Fourier series [5]:

$$\mathbf{u} = \sum_{\alpha \neq \mathbf{0}} [u_{\alpha} \phi_1^{\alpha}, v_{\alpha} \phi_2^{\alpha}, w_{\alpha} \phi_3^{\alpha}], \quad \boldsymbol{\theta} = \sum_{\alpha_3 \neq \mathbf{0}} \theta_{\alpha} \phi_3^{\alpha}, \quad \boldsymbol{p} = \sum_{\alpha \neq \mathbf{0}} p_{\alpha} \phi_4^{\alpha}, \tag{4}$$

where  $\alpha \equiv (\alpha_1, \alpha_2, \alpha_3)$  is the three dimensional multi-index of non-negative integers  $\mathbb{Z}_0$ , and  $u_{\alpha}, v_{\alpha}, w_{\alpha}, \theta_{\alpha}$ ,  $p_{\alpha}$  are coefficients of **u**,  $\theta$ , *p* with respect to the base functions  $\phi_i^{\alpha}$  defined by,

$$\phi_1^{\alpha}(x, y, z) = K_{\alpha} \sin(a\alpha_1 x) \cos(b\alpha_2 y) \cos(\alpha_3 z), \quad \phi_2^{\alpha}(x, y, z) = K_{\alpha} \cos(a\alpha_1 x) \sin(b\alpha_2 y) \cos(\alpha_3 z), \\ \phi_3^{\alpha}(x, y, z) = K_{\alpha} \cos(a\alpha_1 x) \cos(b\alpha_2 y) \sin(\alpha_3 z), \quad \phi_4^{\alpha}(x, y, z) = K_{\alpha} \cos(a\alpha_1 x) \cos(b\alpha_2 y) \cos(\alpha_3 z),$$

where the normalization factor with respect to the usual  $L^2(\Omega)$  inner product  $\langle \cdot, \cdot \rangle$  is

$$K_{\alpha} = \sqrt{(2 - \delta_{0\alpha_1})(2 - \delta_{0\alpha_2})(2 - \delta_{0\alpha_3})/|\Omega|}, \quad \delta_{ij} = \text{Kronecker delta on } i, j$$

The various kinds of norms for  $\mathbf{u}$ ,  $\theta$  and p in (4) can be written as:

$$\begin{split} \|\mathbf{u}\|_{0}^{2} &= \sum_{\alpha \neq \mathbf{0}} \{u_{\alpha}^{2} + v_{\alpha}^{2} + w_{\alpha}^{2}\}, \|\nabla \mathbf{u}\|_{0}^{2} = \sum_{\alpha \neq \mathbf{0}} \{u_{\alpha}^{2} + v_{\alpha}^{2} + w_{\alpha}^{2}\}A_{\alpha}^{2}, \|\nabla^{2}\mathbf{u}\|_{0}^{2} = \sum_{\alpha \neq \mathbf{0}} \{u_{\alpha}^{2} + v_{\alpha}^{2} + w_{\alpha}^{2}\}A_{\alpha}^{4}, \\ \|\theta\|_{0}^{2} &= \sum_{\alpha_{3} \neq 0} \theta_{\alpha}^{2}, \qquad \|\nabla \theta\|_{0}^{2} = \sum_{\alpha_{3} \neq 0} \theta_{\alpha}^{2}A_{\alpha}^{2}, \qquad \|\nabla^{2}\theta\|_{0}^{2} = \sum_{\alpha_{3} \neq 0} \theta_{\alpha}^{2}A_{\alpha}^{4}, \\ \|p\|_{0}^{2} &= \sum_{\alpha \neq \mathbf{0}} p_{\alpha}^{2}, \qquad \|\nabla p\|_{0}^{2} = \sum_{\alpha \neq \mathbf{0}} p_{\alpha}^{2}A_{\alpha}^{2}, \qquad \|\nabla^{2}p\|_{0}^{2} = \sum_{\alpha \neq \mathbf{0}} p_{\alpha}^{2}A_{\alpha}^{4}, \end{split}$$

where  $A_{\alpha} \equiv \sqrt{(a\alpha_1)^2 + (b\alpha_2)^2 + \alpha_3^2}$  provided that the corresponding righthand sides converge. We now define the divergence free and orthogonal system by:

$$\boldsymbol{\Phi}^{\alpha} = \begin{bmatrix} -\frac{a\alpha_{1}\alpha_{3}}{A_{\alpha}B_{\alpha}}\phi_{1}^{\alpha}, -\frac{b\alpha_{2}\alpha_{3}}{A_{\alpha}B_{\alpha}}\phi_{2}^{\alpha}, \frac{B_{\alpha}}{A_{\alpha}}\phi_{3}^{\alpha} \end{bmatrix}, \alpha \in I_{1}, \quad \Psi^{\alpha} = \begin{bmatrix} \frac{b\alpha_{2}}{B_{\alpha}}\phi_{1}^{\alpha}, -\frac{a\alpha_{1}}{B_{\alpha}}\phi_{2}^{\alpha}, 0 \end{bmatrix}, \alpha \in I_{2},$$

where  $B_{\alpha} \equiv \sqrt{(a\alpha_1)^2 + (b\alpha_2)^2}$  and indices subsets are  $I_1 \equiv \{[1,0,1] + \mathbb{Z}_0^3\} \cup \{[0,1,1] + \mathbb{Z}_0^3\}$ ,  $I_2 \equiv [1,1,0] + \mathbb{Z}_0^3$ . Set  $I_0 = I_1 \cup I_2$  and then define the function spaces *V* and *W* with associated usual  $H^1$  – norm as follows:

$$V = \left\{ \mathbf{u} = \sum_{\alpha \in I_0} \left\{ \xi_{\alpha} \Phi^{\alpha} + \eta_{\alpha} \Psi^{\alpha} \right\} : \|\Delta \mathbf{u}\|_0 < \infty \right\} \subset H^2(\Omega)^3,$$
  
$$W = \left\{ \theta = \sum_{\alpha \in I_3} \theta_{\alpha} \phi_3^{\alpha} : \|\Delta \theta\|_0 < \infty \right\} \subset H^2(\Omega), \quad \text{where } I_3 \equiv [0, 0, 1] + \mathbb{Z}_0^3.$$

Note that  $\|\mathbf{u}\|_0^2 = \sum_{\alpha \in I_0} \{\xi_{\alpha}^2 + \eta_{\alpha}^2\}, \|\nabla \mathbf{u}\|_0^2 = \sum_{\alpha \in I_0} \{\xi_{\alpha}^2 + \eta_{\alpha}^2\} A_{\alpha}^2, \|\Delta \mathbf{u}\|_0^2 = \sum_{\alpha \in I_0} \{\xi_{\alpha}^2 + \eta_{\alpha}^2\} A_{\alpha}^4 \text{ for all } \mathbf{u} \in V,$ and  $\|\boldsymbol{\theta}\|_0^2 = \sum_{\alpha \in I_3} \theta_{\alpha}^2, \|\nabla \boldsymbol{\theta}\|_0^2 = \sum_{\alpha \in I_3} \theta_{\alpha}^2 A_{\alpha}^2, \|\Delta \boldsymbol{\theta}\|_0^2 = \sum_{\alpha \in I_3} \theta_{\alpha}^2 A_{\alpha}^4 \text{ for all } \boldsymbol{\theta} \in W.$ 

#### 2 A priori error estimates

For a fixed number  $N \ge 2$ , define the finite dimensional subspaces  $V_N$  and  $W_N$  of V and W by:

$$V_N \equiv \{ \mathbf{u} \in V : \xi_{\alpha} = \eta_{\alpha} = 0, \text{ if } |\alpha| \equiv \alpha_1 + \alpha_2 + \alpha_3 > N \}, \quad W_N \equiv \{ \theta \in W : \theta_{\alpha} = 0, \text{ if } |\alpha| > N \}.$$

Set  $X \equiv V \times W$  and  $X_N \equiv V_N \times W_N$ . Define the projections  $P_N : V \to V_N$  and  $Q_N : W \to W_N$  as in [7]:

$$\langle \nabla (\mathbf{u} - P_N \mathbf{u}), \nabla \mathbf{v} \rangle = 0, \,^{\forall} \mathbf{v} \in V_N, \quad \langle \nabla (\theta - Q_N \theta), \nabla \vartheta \rangle = 0, \,^{\forall} \vartheta \in W_N, \tag{5}$$

with respect to the inner product  $\langle \cdot, \cdot \rangle$  on  $L^2(\Omega)$  or properly extended spaces. Due to orthogonal relations of base functions in *X*, these projections  $P_N$  and  $Q_N$  are truncation operators:

$$P_{N}\mathbf{u}=\sum_{\alpha\in I_{0,N}\equiv I_{0}\cap I,N}\left\{\xi_{\alpha}\Phi^{\alpha}+\eta_{\alpha}\Psi^{\alpha}\right\},\quad Q_{N}\theta=\sum_{\alpha\in I_{3,N}\equiv I_{3}\cap I,N}\theta_{\alpha}\phi_{3}^{\alpha},$$

where  $I_{,N} \equiv \{ \alpha \in \mathbb{Z}_0^3 : |\alpha| \le N \}$ . The sum  $|\alpha|$  of multi-index  $\alpha \in \mathbb{Z}_0^3$  can be considered as the result of the inner product between two vectors  $[a^{-1}, b^{-1}, 1]$  and  $[a\alpha_1, b\alpha_2, \alpha_3]$ . This consideration with the Cauchy-Schwartz inequality gives us

$$|\alpha| \le \|[a^{-1}, b^{-1}, 1]\| \cdot \|[a\alpha_1, b\alpha_2, \alpha_3]\| = C_0 A_\alpha, \quad C_0 \equiv \sqrt{a^{-2} + b^{-2} + 1}.$$
(6)

Note that  $C_0$  depends only on the wave numbers *a* and *b*, so we can say that it depends only on  $\Omega$ . From these characterization of projections and special estimation (6), we have

**Theorem 1** For any  $(\mathbf{u}, \theta) \in X$  and  $(P_N \mathbf{u}, Q_N \theta) \in X_N$  in (5), the following holds:

$$\|\mathbf{u} - P_N \mathbf{u}\|_0 \le \frac{C_0^2}{(N+1)^2} \|\Delta \mathbf{u}\|_0, \quad \|\nabla (\mathbf{u} - P_N \mathbf{u})\|_0 \le \frac{C_0}{N+1} \|\Delta \mathbf{u}\|_0,$$
(7a)

$$\|\boldsymbol{\theta} - \boldsymbol{Q}_N \boldsymbol{\theta}\|_0 \le \frac{C_0^2}{(N+1)^2} \|\Delta \boldsymbol{\theta}\|_0, \ \|\nabla (\boldsymbol{\theta} - \boldsymbol{Q}_N \boldsymbol{\theta})\|_0 \le \frac{C_0}{N+1} \|\Delta \boldsymbol{\theta}\|_0.$$
(7b)

*Proof* Due to (6), we have  $1 \le \frac{C_0}{N+1} A_{\alpha}$  if  $|\alpha| > N$ . Hence we can establish the following estimates:

$$\begin{split} \|\mathbf{u} - P_{N}\mathbf{u}\|_{0}^{2} &= \sum_{\alpha \in I_{0} - I_{0,N}} \{\xi_{\alpha}^{2} + \eta_{\alpha}^{2}\} \leq \frac{C_{0}^{4}}{(N+1)^{4}} \sum_{\alpha \in I_{0} - I_{0,N}} \{\xi_{\alpha}^{2} + \eta_{\alpha}^{2}\} A_{\alpha}^{4} \leq \frac{C_{0}^{4}}{(N+1)^{4}} \|\Delta \mathbf{u}\|_{0}^{2}, \\ \|\nabla(\mathbf{u} - P_{N}\mathbf{u})\|_{0}^{2} &= \sum_{\alpha \in I_{0} - I_{0,N}} \{\xi_{\alpha}^{2} + \eta_{\alpha}^{2}\} A_{\alpha}^{2} \leq \frac{C_{0}^{2}}{(N+1)^{2}} \sum_{\alpha \in I_{0} - I_{0,N}} \{\xi_{\alpha}^{2} + \eta_{\alpha}^{2}\} A_{\alpha}^{4} \leq \frac{C_{0}^{2}}{(N+1)^{2}} \|\Delta \mathbf{u}\|_{0}^{2}, \\ \|\theta - P_{N}\theta\|_{0}^{2} &= \sum_{\alpha \in I_{3} - I_{3,N}} \theta_{\alpha}^{2} \leq \frac{C_{0}^{4}}{(N+1)^{4}} \sum_{\alpha \in I_{3} - I_{3,N}} \theta_{\alpha}^{2} A_{\alpha}^{4} \leq \frac{C_{0}^{4}}{(N+1)^{4}} \|\Delta \theta\|_{0}^{2}, \\ |\nabla(\theta - P_{N}\theta)\|_{0}^{2} &= \sum_{\alpha \in I_{3} - I_{3,N}} \theta_{\alpha}^{2} A_{\alpha}^{2} \leq \frac{C_{0}^{2}}{(N+1)^{2}} \sum_{\alpha \in I_{3} - I_{3,N}} \theta_{\alpha}^{2} A_{\alpha}^{4} \leq \frac{C_{0}^{2}}{(N+1)^{2}} \|\Delta \theta\|_{0}^{2}. \end{split}$$

These lead (7) after taking square root of them.  $\Box$ 

As usual, the  $L^{\infty}$  norms  $\|\mathbf{u}\|_{\infty}$  and  $\|\boldsymbol{\theta}\|_{\infty}$  of  $\mathbf{u} \in V$  and  $\boldsymbol{\theta} \in W$  are defined by

$$\|\mathbf{u}\|_{\infty} \equiv \sup_{\mathbf{x}\in\Omega} |\mathbf{u}(\mathbf{x})|, \quad \|\theta\|_{\infty} \equiv \sup_{\mathbf{x}\in\Omega} |\theta(\mathbf{x})|, \quad |\mathbf{u}| = \sqrt{u^2 + v^2 + w^2}, \quad |\theta| = \sqrt{\theta^2},$$
$$\mathbf{u} = [u, v, w], \quad u = \sum_{\alpha\in I_0} u_{\alpha}\phi_1^{\alpha}, \quad v = \sum_{\alpha\in I_0} v_{\alpha}\phi_2^{\alpha}, \quad w = \sum_{\alpha\in I_0} w_{\alpha}\phi_3^{\alpha}, \quad \theta = \sum_{\alpha\in I_3} \theta_{\alpha}\phi_3^{\alpha}.$$

For a fixed  $\alpha \in I_0$ , the vector  $\xi_{\alpha} \Phi^{\alpha} + \eta_{\alpha} \Psi^{\alpha}$  can be written of the form:  $[u_{\alpha} \phi_1^{\alpha}, v_{\alpha} \phi_2^{\alpha}, w_{\alpha} \phi_3^{\alpha}]$  with

$$u_{\alpha} \equiv -\frac{a\alpha_{1}\alpha_{3}}{A_{\alpha}B_{\alpha}}\xi_{\alpha} + \frac{b\alpha_{2}}{B_{\alpha}}\eta_{\alpha}, \quad v_{\alpha} \equiv -\frac{b\alpha_{2}\alpha_{3}}{A_{\alpha}B_{\alpha}}\xi_{\alpha} - \frac{a\alpha_{1}}{B_{\alpha}}\eta_{\alpha}, \quad w_{\alpha} \equiv \frac{B_{\alpha}}{A_{\alpha}}\xi_{\alpha}, \quad u_{\alpha}^{2} + v_{\alpha}^{2} + w_{\alpha}^{2} = \xi_{\alpha}^{2} + \eta_{\alpha}^{2}.$$

Now, the square sum  $f_{\alpha} \equiv (u_{\alpha}\phi_1^{\alpha})^2 + (v_{\alpha}\phi_2^{\alpha})^2 + (w_{\alpha}\phi_1^{\alpha})^2$  can be bounded on  $\Omega$  as follows, setting  $f_1 \equiv \cos^2(a\alpha_1 x), f_2 \equiv \cos^2(b\alpha_2 y), f_3 \equiv \cos^2(\alpha_3 z),$ 

$$f_{\alpha} = K_{\alpha}^{2}[u_{\alpha}^{2}(1-f_{1})f_{2}f_{3} + v_{\alpha}^{2}f_{1}(1-f_{2})f_{3} + w_{\alpha}^{2}f_{1}f_{2}(1-f_{3})]$$
  
$$\leq K_{\alpha}^{2}[u_{\alpha}^{2} + v_{\alpha}^{2} + w_{\alpha}^{2}][(1-f_{1})f_{2}f_{3} + f_{1}(1-f_{2})f_{3} + f_{1}f_{2}(1-f_{3})] \leq K_{\alpha}^{2}[\xi_{\alpha}^{2} + \eta_{\alpha}^{2}],$$

since, for each *i*,  $f_i$  on  $\Omega$  takes value in the interval [0,1]. Thus, we have  $\|\xi_{\alpha}\Phi^{\alpha} + \eta_{\alpha}\Psi^{\alpha}\|_{\infty} \leq K_{\alpha}\sqrt{\xi_{\alpha}^2 + \eta_{\alpha}^2}$  for any  $\alpha \in I_0$ .

**Lemma 2** For any  $(\mathbf{u}, \boldsymbol{\theta}) \in X$ , it holds that

$$\begin{aligned} \|\mathbf{u}\|_{\infty} &\leq \frac{\pi}{3}\sqrt{6 - \frac{2\pi^2}{5}}C_1 \|\Delta \mathbf{u}\|_0 < 1.50015C_1 \|\Delta \mathbf{u}\|_0, \\ \|\theta\|_{\infty} &\leq \frac{\pi}{3}\sqrt{6 - \frac{36\zeta(3)}{\pi^2} + \frac{\pi^2}{5}}C_1 \|\Delta \theta\|_0 < 1.98398C_1 \|\Delta \theta\|_0, \end{aligned}$$

where  $C_1 \equiv C_0^2 |\Omega|^{-\frac{1}{2}}$  depends only on  $\Omega$  and  $\zeta(s) \equiv \sum_{n=1}^{\infty} \frac{1}{n^s}$  is the Riemann zeta function for s > 1.

*Proof* From the above argument,  $\|\mathbf{u}\|_{\infty} \leq \sum_{\alpha \in I_0} \|\xi_{\alpha} \Phi^{\alpha} + \eta_{\alpha} \Psi^{\alpha}\|_{\infty} \leq \sum_{\alpha \in I_0} K_{\alpha} \sqrt{\xi_{\alpha}^2 + \eta_{\alpha}^2}$ . And the Cauchy-Schwartz inequality shows that

$$\|\mathbf{u}\|_{\infty} \leq \sum_{\alpha \in I_0} K_{\alpha} \sqrt{\xi_{\alpha}^2 + \eta_{\alpha}^2} \leq \sqrt{\sum_{\alpha \in I_0} K_{\alpha}^2 A_{\alpha}^{-4}} \sqrt{\sum_{\alpha \in I_0} \{\xi_{\alpha}^2 + \eta_{\alpha}^2\} A_{\alpha}^4} \leq \sqrt{C} \|\Delta \mathbf{u}\|_0, \quad C \equiv \sum_{\alpha \in I_0} K_{\alpha}^2 A_{\alpha}^{-4}.$$

The indices subset  $I_0$  can be decomposed into four mutually disjoint subsets:

$$\begin{split} I_0^{1,1,0} &\equiv [1,1,0] + \mathbb{Z}_0^2 \times \{0\}, \quad I_0^{1,0,1} \equiv [1,0,1] + \mathbb{Z}_0 \times \{0\} \times \mathbb{Z}_0, \\ I_0^{0,1,1} &\equiv [0,1,1] + \{0\} \times \mathbb{Z}_0^2, \quad I_0^{1,1,1} \equiv [1,1,1] + \mathbb{Z}_0^3. \end{split}$$

And the numbers n(k) of non-negative integer solutions of  $|\alpha| = k$  and the values  $K_{\alpha}^2$  on these subsets are:

$$n(k)\Big|_{I_0-I_0^{1,1,1}} = k-1, \quad n(k)\Big|_{I_0^{1,1,1}} = \frac{(k-1)(k-2)}{2}, \quad K_\alpha^2\Big|_{I_0-I_0^{1,1,1}} = \frac{4}{|\Omega|}, \quad K_\alpha^2\Big|_{I_0^{1,1,1}} = \frac{8}{|\Omega|}.$$

Due to (6) and the above relations, we can bound *C* as follows:

$$\begin{split} C &\leq C_0^4 \sum_{\alpha \in I_0} K_\alpha^2 |\alpha|^{-4} = \frac{4C_0^4}{|\Omega|} \left[ 3\sum_{k=2}^\infty \frac{k-1}{k^4} + 2\sum_{k=3}^\infty \frac{(k-1)(k-2)}{2k^4} \right] = 4C_1^2 \sum_{k=1}^\infty \frac{(k-1)(k+1)}{k^4} \\ &= 4C_1^2 \sum_{k=1}^\infty \left[ \frac{1}{k^2} - \frac{1}{k^4} \right] = 4C_1^2 \left[ \frac{\pi^2}{6} - \frac{\pi^4}{90} \right] = \frac{\pi^2}{9} \left[ 6 - \frac{2\pi^2}{5} \right] C_1^2, \end{split}$$

which proves the first part of the lemma.

Next, taking account that  $\|\theta\|_{\infty} \leq \sum_{\alpha \in I_3} \|\theta_{\alpha} \phi_3^{\alpha}\|_{\infty} = \sum_{\alpha \in I_3} K_{\alpha} |\theta_{\alpha}|$ , similar to the above, we have

$$\|\theta\|_{\infty} \leq \sum_{\alpha \in I_3} K_{\alpha} |\theta_{\alpha}| \leq \sqrt{\sum_{\alpha \in I_3} K_{\alpha}^2 A_{\alpha}^{-4}} \sqrt{\sum_{\alpha \in I_3} \theta_{\alpha}^2 A_{\alpha}^4} \leq \sqrt{\tilde{C}} \|\Delta \theta\|_0, \quad \text{with} \quad \tilde{C} \equiv \sum_{\alpha \in I_3} K_{\alpha}^2 A_{\alpha}^{-4}$$

The indices subset  $I_3$  can be decomposed into four mutually disjoint subsets:

$$I_3^{0,0,1} \equiv [0,0,1] + \{0\}^2 \times \mathbb{Z}_0, I_3^{1,0,1} \equiv I_0^{1,0,1}, I_3^{0,1,1} \equiv I_0^{0,1,1}, I_3^{1,1,1} \equiv I_0^{1,1,1}$$

And n(k) and  $K^2_{\alpha}$  on these subsets are:

$$\begin{split} n(k)\Big|_{I_3^{0,0,1}} &= 1, \quad n(k)\Big|_{I_3^{1,0,1} \cup I_3^{0,1,1}} = k-1, \quad n(k)\Big|_{I_3^{1,1,1}} = \frac{(k-1)(k-2)}{2}, \\ K_\alpha^2\Big|_{I_3^{0,0,1}} &= \frac{2}{|\Omega|}, \quad K_\alpha^2\Big|_{I_3^{1,0,1} \cup I_3^{0,1,1}} = \frac{4}{|\Omega|}, \quad K_\alpha^2\Big|_{I_3^{1,1,1}} = \frac{8}{|\Omega|}. \end{split}$$

Thus, the bound  $\tilde{C}$  can be obtained as follows:

$$\begin{split} \tilde{C} &\leq C_0^4 \sum_{\alpha \in I_3} K_\alpha^2 |\alpha|^{-4} = \frac{2C_0^4}{|\Omega|} \left[ \sum_{k=1}^\infty \frac{1}{k^4} + 4 \sum_{k=2}^\infty \frac{k-1}{k^4} + 4 \sum_{k=3}^\infty \frac{(k-1)(k-2)}{2k^4} \right] = 2C_1^2 \sum_{k=1}^\infty \frac{2k^2 - 2k + 1}{k^4} \\ &= 2C_1^2 \sum_{k=1}^\infty \left[ \frac{2}{k^2} - \frac{2}{k^3} + \frac{1}{k^4} \right] = 2C_1^2 \left[ \frac{\pi^2}{3} - 2\zeta(3) + \frac{\pi^4}{90} \right] = \frac{\pi^2}{9} \left[ 6 - \frac{36\zeta(3)}{\pi^2} + \frac{\pi^2}{5} \right] C_1^2, \end{split}$$

which completes the proof.  $\Box$ 

Since  $\Omega$  is a rectangle with side lengths  $\frac{2\pi}{a}$ ,  $\frac{2\pi}{b}$ , and  $\pi$ , we can also use the result in [6] as follows  $(0 < a, b \le 1)$ :

$$\begin{split} \|\theta\|_{\infty} &\leq |\Omega|^{-\frac{1}{2}} \left[ \gamma_{0} \|\theta\|_{0} + \frac{\gamma_{1}\pi c_{1}}{\sqrt{3}} \|\nabla\theta\|_{0} + \frac{\gamma_{2}\pi^{2}c_{2}}{3} \|\nabla^{2}\theta\|_{0} \right], \\ c_{1} &= \sqrt{4(a^{-2} + b^{-2}) + 1} \geq \sqrt{3}C_{0}, \\ c_{2} &= \sqrt{c_{1}^{4} + \frac{4}{5}} [16(a^{-4} + b^{-4}) + 1] \geq \sqrt{\frac{19}{15}}c_{1}^{2} \geq 3\sqrt{\frac{19}{15}}C_{0}^{2}, \\ \frac{\gamma_{2}\pi^{2}c_{2}}{3} \geq 5.59879 \,\gamma_{2}\frac{\pi}{3}\sqrt{6 - \frac{36\zeta(3)}{\pi^{2}} + \frac{\pi^{2}}{5}}C_{0}^{2} \geq 2.31862 \,\frac{\pi}{3}\sqrt{6 - \frac{36\zeta(3)}{\pi^{2}} + \frac{\pi^{2}}{5}}C_{0}^{2} \\ \end{split}$$

Here, we have used the fact taht  $\gamma_2$  is estimated as 0.41413 in [6]. Thus the coefficients in Lemma 2 are at least twice finer than those in [6].

**Corollary 3** Under the same assumptions of Theorem 1, the following holds for the same constant  $C_1$  in Lemma 2:

$$\|\mathbf{u} - P_N \mathbf{u}\|_{\infty} \le 2C_1 \sqrt{\frac{1}{N} - \frac{1}{3(N+1)^3}} \|\Delta \mathbf{u}\|_0 < \frac{2C_1}{\sqrt{N}} \|\Delta \mathbf{u}\|_0,$$
(8a)

$$\|\theta - Q_N \theta\|_{\infty} \le 2C_1 \sqrt{\frac{1}{N} - \frac{1}{2(N+1)^2} + \frac{1}{6N^3}} \|\Delta \theta\|_0 < \frac{2C_1}{\sqrt{N}} \|\Delta \theta\|_0.$$
(8b)

-

*Proof* We can proceed as that of Lemma 2 and need to change C for each case. For  $\|\mathbf{u} - P_N \mathbf{u}\|_{\infty}$ ,

$$\begin{split} C &\equiv \sum_{\alpha \in I_0 - I_{0,N}} K_{\alpha}^2 A_{\alpha}^{-4} \le C_0^4 \sum_{\alpha \in I_0 - I_{0,N}} K_{\alpha}^2 |\alpha|^{-4} = 4C_1^2 \left[ 3\sum_{k=N+1}^{\infty} \frac{k-1}{k^4} + 2\sum_{k=N+1}^{\infty} \frac{(k-1)(k-2)}{2k^4} \right] \\ &= 4C_1^2 \sum_{k=N+1}^{\infty} \frac{(k-1)(k+1)}{k^4} = 4C_1^2 \sum_{k=N+1}^{\infty} \left[ \frac{1}{k^2} - \frac{1}{k^4} \right] < 4C_1^2 \left[ \int_N^{\infty} \frac{1}{x^2} dx - \int_{N+1}^{\infty} \frac{1}{x^4} dx \right] \\ &= 4 \left[ \frac{1}{N} - \frac{1}{3(N+1)^3} \right] C_1^2 < \frac{4}{N} C_1^2. \end{split}$$

Similarly, we have the following estimates for  $\|\theta - Q_N \theta\|_{\infty}$ :

$$\begin{split} \tilde{C} &\equiv \sum_{\alpha \in I_3 - I_{3,N}} K_{\alpha}^2 A_{\alpha}^{-4} \leq C_0^4 \sum_{\alpha \in I_3 - I_{3,N}} K_{\alpha}^2 |\alpha|^{-4} = 2C_1^2 \left[ \sum_{k=N+1}^{\infty} \frac{1}{k^4} + 4 \sum_{k=N+1}^{\infty} \frac{k-1}{k^4} + 4 \sum_{k=N+1}^{\infty} \frac{(k-1)(k-2)}{2k^4} \right] \\ &= 2C_1^2 \sum_{k=N+1}^{\infty} \frac{2k^2 - 2k + 1}{k^4} = 2C_1^2 \sum_{k=N+1}^{\infty} \left[ \frac{2}{k^2} - \frac{2}{k^3} + \frac{1}{k^4} \right] < 2C_1^2 \left[ \int_N^{\infty} \frac{2}{x^2} dx - \int_{N+1}^{\infty} \frac{2}{x^3} dx + \int_N^{\infty} \frac{1}{x^4} dx \right] \\ &= 2 \left[ \frac{2}{N} - \frac{1}{(N+1)^2} + \frac{1}{3N^3} \right] C_1^2 = 4 \left[ \frac{1}{N} - \frac{1}{2(N+1)^2} + \frac{1}{6N^3} \right] C_1^2 < \frac{4}{N} C_1^2. \end{split}$$

The last inequality holds due to the assumption  $N \ge 2$ .  $\Box$ 

#### 3 A fixed point formulation

The steady state solution of (3) can be written of the form:

$$-\Delta \mathbf{u} + \nabla p = \mathbf{f}(\mathbf{u}, \boldsymbol{\theta}), \tag{9a}$$

$$\nabla \cdot \mathbf{u} = 0, \tag{9b}$$

$$-\Delta \theta = g(\mathbf{u}, \theta), \tag{9c}$$

where the right hand sides of (9) are defined by

$$\mathbf{f}(\mathbf{u},\boldsymbol{\theta}) = -\frac{1}{\mathscr{P}}(\mathbf{u}\cdot\nabla)\mathbf{u} + \mathscr{R}\boldsymbol{\theta}\mathbf{e}_z, \quad g(\mathbf{u},\boldsymbol{\theta}) = -(\mathbf{u}\cdot\nabla)\boldsymbol{\theta} + w.$$

If  $\mathbf{u} \in V$ , then  $P_N \mathbf{u}$  satisfies (9b) and  $-\Delta P_N \mathbf{u}$  converges to  $-\Delta \mathbf{u}$  in  $L^2$  sense. Taking inner product for both sides of (9a) with  $-\Delta P_N \mathbf{u}$ , we have

$$\langle -\Delta \mathbf{u}, -\Delta P_N \mathbf{u} \rangle = \langle -\Delta \mathbf{u}, -\Delta P_N \mathbf{u} \rangle + \langle p, \Delta \nabla P_N \mathbf{u} \rangle = \langle -\Delta \mathbf{u}, -\Delta P_N \mathbf{u} \rangle + \langle p, \nabla \Delta P_N \mathbf{u} \rangle$$
$$= \langle -\Delta \mathbf{u}, -\Delta P_N \mathbf{u} \rangle + \langle \nabla p, -\Delta P_N \mathbf{u} \rangle = \langle \mathbf{f}, -\Delta P_N \mathbf{u} \rangle.$$

This identity converges to  $\langle -\Delta \mathbf{u}, -\Delta \mathbf{u} \rangle = \langle \mathbf{f}, -\Delta \mathbf{u} \rangle$  which means  $\|\Delta \mathbf{u}\|_0 \le \|\mathbf{f}\|_0$ . And taking inner product for both sides of (9c) with  $-\Delta \theta$ , we have  $\langle -\Delta \theta, -\Delta \theta \rangle = \langle g, -\Delta \theta \rangle$  and  $\|\Delta \theta\|_0 \le \|g\|_0$ . Now, setting  $F(\mathbf{u}, \theta) \equiv (\mathbf{f}(\mathbf{u}, \theta), g(\mathbf{u}, \theta))$ , the weak form of (9) is written as :

 $\nabla \cdot \bar{\mathbf{u}} =$ 

$$\langle \nabla(\mathbf{u}, \boldsymbol{\theta}), \nabla(\mathbf{v}, \boldsymbol{\vartheta}) \rangle = \langle F(\mathbf{u}, \boldsymbol{\theta}), (\mathbf{v}, \boldsymbol{\vartheta}) \rangle, \quad \forall (\mathbf{v}, \boldsymbol{\vartheta}) \in X$$

We call the solution operator  $\mathscr{S}$  for (9) as *Stokes operator*. Thus  $(\mathbf{u}, \theta) = \mathscr{S}F(\mathbf{u}, \theta)$  means

$$\langle \nabla \mathscr{S}F(\mathbf{u}, \theta), \nabla(\mathbf{v}, \vartheta) \rangle = \langle F(\mathbf{u}, \theta), (\mathbf{v}, \vartheta) \rangle, \quad \forall (\mathbf{v}, \vartheta) \in X.$$
 (10)

Note that we always have  $\mathscr{S}^{-1}(\mathbf{u}, \boldsymbol{\theta}) = (-\Delta \mathbf{u} + \nabla p, -\Delta \boldsymbol{\theta})$  with an associated pressure  $p = p(\mathbf{u}, \boldsymbol{\theta})$ .

Usually, we use Newton's method (see [4]) to get an approximate solution  $(\mathbf{u}_N, \theta_N) \in X_N$  of (9) and define the approximate pressure  $p_N$  by

$$abla p_N \equiv \mathbf{f}_N(\mathbf{u}_N, \boldsymbol{\theta}_N) + \Delta \mathbf{u}_N$$

where  $\mathbf{f}_N$  is the truncation up to  $I_N$  of the expansion of  $\mathbf{f}$ . For the solution  $(\mathbf{u}, \theta)$  of (9) with its associated pressure p, let  $(\bar{\mathbf{u}}, \bar{\theta}) \equiv (\mathbf{u} - \mathbf{u}_N, \theta - \theta_N)$  and  $\bar{p} \equiv p - p_N$ . Then we have the following residual equations:

$$-\Delta \bar{\mathbf{u}} + \nabla \bar{p} = \mathbf{f}(\mathbf{u}_N + \bar{\mathbf{u}}, \theta_N + \bar{\theta}) + \Delta \mathbf{u}_N - \nabla p_N, \qquad (11a)$$

$$-\Delta\bar{\theta} = g(\mathbf{u}_N + \bar{\mathbf{u}}, \theta_N + \bar{\theta}) + \Delta\theta_N, \qquad (11c)$$

Set  $\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \equiv (\mathbf{f}(\mathbf{u}_N + \bar{\mathbf{u}},\theta_N + \bar{\theta}) + \Delta \mathbf{u}_N - \nabla p_N, g(\mathbf{u}_N + \bar{\mathbf{u}},\theta_N + \bar{\theta}) + \Delta \theta_N) \equiv (\bar{\mathbf{f}}(\bar{\mathbf{u}},\bar{\theta}),\bar{g}(\bar{\mathbf{u}},\bar{\theta}))$ , then the Stokes operator  $\mathscr{S}$  gives us a fixed point problem from (11):

$$(\bar{\mathbf{u}},\bar{\theta}) = \mathscr{S}\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \equiv \mathscr{K}(\bar{\mathbf{u}},\bar{\theta}).$$
(12)

Since  $X \subset H^1(\Omega)^4$ ,  $\mathcal{K}$  is a compact operator on X. Hence by Schauder's fixed point theorem, if we find a nonempty, closed, convex, and bounded set  $U \subset X$  satisfying  $\mathcal{K}U \subset U$ , then there exists a solution of (12) in U which is called a candidate set.

Define  $\mathbf{P}_N : X \to X_N$  by  $\mathbf{P}_N = (P_N, Q_N)$ , then (5) can be simplified as: for  $(\mathbf{u}, \theta) \in X$ 

$$\langle \nabla((\mathbf{u}, \boldsymbol{\theta}) - \mathbf{P}_N(\mathbf{u}, \boldsymbol{\theta})), \nabla(\mathbf{v}, \boldsymbol{\vartheta}) \rangle = 0, \quad \forall (\mathbf{v}, \boldsymbol{\vartheta}) \in X_N.$$
 (13)

Then (12) can be decomposed into two parts:

$$\mathbf{P}_{N}(\bar{\mathbf{u}},\bar{\theta}) = \mathbf{P}_{N}\mathscr{K}(\bar{\mathbf{u}},\bar{\theta}), \tag{14a}$$

$$(I - \mathbf{P}_N)(\bar{\mathbf{u}}, \bar{\theta}) = (I - \mathbf{P}_N) \mathscr{K}(\bar{\mathbf{u}}, \bar{\theta}).$$
(14b)

The Fréchet derivative  $F'(\mathbf{u}, \theta)$  of F at  $(\mathbf{u}, \theta)$  has the form: for any  $(\bar{\mathbf{u}}, \bar{\theta}) \in X$ ,

$$F'(\mathbf{u},\boldsymbol{\theta})(\bar{\mathbf{u}},\boldsymbol{\theta}) \equiv \left(\mathbf{f}'(\mathbf{u},\boldsymbol{\theta})(\bar{\mathbf{u}},\boldsymbol{\theta}), g'(\mathbf{u},\boldsymbol{\theta})(\bar{\mathbf{u}},\boldsymbol{\theta})\right),$$
  
$$\mathbf{f}'(\mathbf{u},\boldsymbol{\theta})(\bar{\mathbf{u}},\bar{\boldsymbol{\theta}}) \equiv -\frac{1}{\mathscr{P}}\left[(\mathbf{u}\cdot\nabla)\bar{\mathbf{u}} + (\bar{\mathbf{u}}\cdot\nabla)\mathbf{u}\right] + \mathscr{R}\bar{\boldsymbol{\theta}}\mathbf{e}_{z},$$
  
$$g'(\mathbf{u},\boldsymbol{\theta})(\bar{\mathbf{u}},\bar{\boldsymbol{\theta}}) \equiv -\left[(\mathbf{u}\cdot\nabla)\bar{\boldsymbol{\theta}} + (\bar{\mathbf{u}}\cdot\nabla)\boldsymbol{\theta}\right] + \bar{w}.$$

Now, define  $\mathscr{L}_N : X_N \to X_N$  by

$$\mathscr{L}_{N} \equiv \mathbf{P}_{N} \left[ I - \mathscr{S} F'(\mathbf{u}_{N}, \boldsymbol{\theta}_{N}) \right] \Big|_{X_{N}}$$

and assume  $\mathscr{L}_N$  is regular or one-to-one and onto. And we can express  $\mathscr{L}_N$  as:

$$\mathscr{L}_{N} = \mathbf{P}_{N}\mathscr{S}[\mathscr{S}^{-1} - F'(\mathbf{u}_{N}, \boldsymbol{\theta}_{N})]\Big|_{X_{n}} = \mathbf{P}_{N}\mathscr{S}\mathscr{L}_{0}, \quad \mathscr{L}_{0} \equiv [\mathscr{S}^{-1} - F'(\mathbf{u}_{N}, \boldsymbol{\theta}_{N})]\Big|_{X_{n}}$$

Define the Newton–like iteration operator  $\mathcal{N} : X \to X_N$  for (12) and the new map  $\mathcal{T}$  as follows:

$$\mathscr{N} \equiv \mathbf{P}_N - \mathscr{L}_N^{-1} \mathbf{P}_N (I - \mathscr{K}), \quad \mathscr{T} \equiv \mathscr{N} + (I - \mathbf{P}_N) \mathscr{K}.$$

The second part of  $\mathscr{T}$  is expected to be small or contractive if the truncation number N is sufficiently large. The operator  $\mathscr{N}$  is also compact since it maps X into the finite dimensional space  $X_N$ , and so is  $\mathscr{T}$ . **Lemma 4** The problem (14) is equivalent to the following fixed point problem:

$$(\bar{\mathbf{u}}, \bar{\boldsymbol{\theta}}) = \mathscr{T}(\bar{\mathbf{u}}, \bar{\boldsymbol{\theta}}). \tag{15}$$

*Proof* Assume  $(\bar{\mathbf{u}}, \bar{\theta}) \in X$  satisfies (14), then  $\mathscr{N}(\bar{\mathbf{u}}, \bar{\theta}) = \mathbf{P}_N(\bar{\mathbf{u}}, \bar{\theta})$  which means

$$\mathscr{T}(\bar{\mathbf{u}},\bar{\theta}) = \mathbf{P}_N(\bar{\mathbf{u}},\bar{\theta}) + (I - \mathbf{P}_N)\mathscr{K}(\bar{\mathbf{u}},\bar{\theta}) = \mathbf{P}_N(\bar{\mathbf{u}},\bar{\theta}) + (I - \mathbf{P}_N)(\bar{\mathbf{u}},\bar{\theta}) = (\bar{\mathbf{u}},\bar{\theta}).$$

Thus  $(\bar{\mathbf{u}}, \bar{\theta})$  satisfies (15). On the other hand, if  $(\bar{\mathbf{u}}, \bar{\theta})$  satisfies (15), then

$$\begin{aligned} \mathbf{P}_{N}(\bar{\mathbf{u}},\bar{\theta}) &= \mathbf{P}_{N}\mathscr{T}(\bar{\mathbf{u}},\bar{\theta}) = \mathbf{P}_{N}\mathscr{N}(\bar{\mathbf{u}},\bar{\theta}) = \mathbf{P}_{N}(\bar{\mathbf{u}},\bar{\theta}) - \mathscr{L}_{N}^{-1}\mathbf{P}_{N}(I-\mathscr{K})(\bar{\mathbf{u}},\bar{\theta}) \\ &\Rightarrow \mathscr{L}_{N}^{-1}\mathbf{P}_{N}(I-\mathscr{K})(\bar{\mathbf{u}},\bar{\theta}) = 0 \Rightarrow \mathbf{P}_{N}(I-\mathscr{K})(\bar{\mathbf{u}},\bar{\theta}) = 0 \Rightarrow \mathbf{P}_{N}(\bar{\mathbf{u}},\bar{\theta}) = \mathbf{P}_{N}\mathscr{K}(\bar{\mathbf{u}},\bar{\theta}), \\ (I-\mathbf{P}_{N})(\bar{\mathbf{u}},\bar{\theta}) &= (I-\mathbf{P}_{N})\mathscr{T}(\bar{\mathbf{u}},\bar{\theta}) = (I-\mathbf{P}_{N})\mathscr{K}(\bar{\mathbf{u}},\bar{\theta}). \end{aligned}$$

Here the second implication is due to the assumption on the regularity of  $\mathscr{L}_N$ .  $\Box$ 

From Lemma 4, we have an alternative verification condition of the form:  $\mathscr{T}U \subset U$  for a candidate set U which is nonempty, closed, convex, and bounded in X.

Now, for given real numbers  $\overline{\xi}_{\alpha}, \overline{\eta}_{\alpha}, \overline{\theta}_{\alpha} \ge 0$ , set real intervals as  $[\overline{\xi}_{\alpha}] \equiv [-\overline{\xi}_{\alpha}, \overline{\xi}_{\alpha}], [\overline{\eta}_{\alpha}] \equiv [-\overline{\eta}_{\alpha}, \overline{\eta}_{\alpha}], [\overline{\theta}_{\alpha}] \equiv [-\overline{\theta}_{\alpha}, \overline{\theta}_{\alpha}]$ , and define

$$U_{N} \equiv \left\{ \left( \sum_{\alpha \in I_{0,N}} \left\{ \xi_{\alpha} \Phi^{\alpha} + \eta_{\alpha} \Psi^{\alpha} \right\}, \sum_{\alpha \in I_{3,N}} \theta_{\alpha} \phi_{3}^{\alpha} \right) \in X_{N} : \xi_{\alpha} \in [\overline{\xi}_{\alpha}], \eta_{\alpha} \in [\overline{\eta}_{\alpha}], \theta_{\alpha} \in [\overline{\theta}_{\alpha}] \right\}.$$
(16)

And for given  $m_1, m_2 \ge 0$ , we define

$$U_{*} \equiv \left\{ (\mathbf{u}, \theta) \in X_{N}^{\perp} : \begin{array}{l} \|\mathbf{u}\|_{0} \leq \frac{C_{0}^{2}}{(N+1)^{2}} m_{1}, \|\nabla \mathbf{u}\|_{0} \leq \frac{C_{0}}{N+1} m_{1}, \|\mathbf{u}\|_{\infty} \leq \frac{2C_{1}}{\sqrt{N}} m_{1}, \\ \|\theta\|_{0} \leq \frac{C_{0}^{2}}{(N+1)^{2}} m_{2}, \|\nabla \theta\|_{0} \leq \frac{C_{0}}{N+1} m_{2} \end{array} \right\}.$$
(17)

Here  $X_N^{\perp}$  is the orthogonal complement of  $X_N$  in X with respect to the projection  $\mathbf{P}_N$  defined by (13). Now, set  $U \equiv U_N \oplus U_*$ , then we obtain:

**Theorem 5** Let  $U_N$ ,  $U_*$  and U be sets defined as above. If

$$\mathcal{N}U \subset U_N,$$
 (18a)

$$(I - \mathbf{P}_N) \mathscr{K} U \subset U_*. \tag{18b}$$

then there exists a fixed point of  $\mathcal{T}$  in U.

*Proof* Clearly,  $(0,0) \in U$  which means U is non-empty. Due to the definition, U is closed, convex and bounded in X. Under the condition (18), we have

$$\mathscr{T}U \subset \mathscr{N}U + (I - \mathbf{P}_N)\mathscr{K}U \subset U_N + U_* = U.$$

Since  $\mathscr{T}$  is compact, there exists a fixed point of  $\mathscr{T}$  in U by Schauder's fixed point theorem.  $\Box$ 

#### 4 Computable verification conditions

To construct the candidate set U in X satisfying (18), we use an algorithm based on iterative scheme as in [7].

First, set the initial values  $\overline{\xi}_{\alpha}^{(0)} = \overline{\eta}_{\alpha}^{(0)} = \overline{\theta}_{\alpha}^{(0)} = 0$ , and  $m_1^{(0)} = m_2^{(0)} = 0$ , which means  $U^{(0)}$  contains only one element  $(\mathbf{0}, 0)$ . For  $k \ge 0$ , with a fixed inflation factor  $0 < \delta \ll 1$ , set

$$\overline{\xi}_{\alpha}^{(k+\frac{1}{2})} = \overline{\xi}_{\alpha}^{(k)}(1+\delta), \quad \overline{\eta}_{\alpha}^{(k+\frac{1}{2})} = \overline{\eta}_{\alpha}^{(k)}(1+\delta), \quad \overline{\theta}_{\alpha}^{(k+\frac{1}{2})} = \overline{\theta}_{\alpha}^{(k)}(1+\delta), \quad m_{i}^{(k+\frac{1}{2})} = m_{i}^{(k)}(1+\delta), \quad i = 1, 2,$$

which define  $\delta$ -inflations  $U_N^{(k+\frac{1}{2})}$  and  $U_*^{(k+\frac{1}{2})}$  of  $U_N^{(k)}$  and  $U_*^{(k)}$  respectively. Set the  $\delta$ -inflation  $U^{(k+\frac{1}{2})}$  of  $U^{(k)}$  as the direct sum of  $U_N^{(k+\frac{1}{2})}$  and  $U_*^{(k+\frac{1}{2})}$ , i.e.,  $U^{(k+\frac{1}{2})} \equiv U_N^{(k+\frac{1}{2})} \oplus U_*^{(k+\frac{1}{2})}$ . Now,  $U^{(k+1)}$  can be constructed as the direct sum of  $U_N^{(k+1)}$  and  $U_*^{(k+1)}$  as follows:

$$U_N^{(k+1)} \equiv \mathcal{N}U^{(k+\frac{1}{2})}, \quad m_1^{(k+1)} \equiv \left\| \bar{\mathbf{f}}(U^{(k+\frac{1}{2})}) \right\|_0, \quad m_2^{(k+1)} \equiv \left\| \bar{g}(U^{(k+\frac{1}{2})}) \right\|_0, \tag{19}$$

where  $||f(U)||_0 \equiv \sup\{||f(\mathbf{u}, \theta)||_0 : (\mathbf{u}, \theta) \in U\}$  for any function f. Note that  $U^{(k+1)}$  cannot be calculated exactly, but its over–estimated enclosure can be obtained and will be set as  $U_N^{(k+1)}$  in the actual calculation on a computer. Thus the verification condition in a computer is:

**Theorem 6** For some k, if the following conditions

$$\overline{\xi}_{\alpha}^{(k+1)} < \overline{\xi}_{\alpha}^{(k+\frac{1}{2})}, \quad \overline{\eta}_{\alpha}^{(k+1)} < \overline{\eta}_{\alpha}^{(k+\frac{1}{2})}, \quad \overline{\theta}_{\alpha}^{(k+1)} < \overline{\theta}_{\alpha}^{(k+\frac{1}{2})}, \quad m_i^{(k+1)} < m_i^{(k+\frac{1}{2})}, i = 1, 2,$$
(20)

hold, then the set  $U^{(k+\frac{1}{2})}$  contains an element  $(\bar{\mathbf{u}}, \bar{\theta})$  satisfying  $(\bar{\mathbf{u}}, \bar{\theta}) = \mathscr{T}(\bar{\mathbf{u}}, \bar{\theta})$ .

*Proof* Due to Theorem 5, it is sufficient to check (18) holds for  $U^{(k+\frac{1}{2})}$ . By the condition (20) and the definition (19), we have  $\mathscr{N}U^{(k+\frac{1}{2})} = U_N^{(k+1)} \subset U_N^{(k+\frac{1}{2})}$ . And for any  $(\mathbf{u}, \theta) \in (I - \mathbf{P}_N) \mathscr{K}U^{(k+\frac{1}{2})}$ , there exists  $(\bar{\mathbf{u}}, \bar{\theta}) \in U^{(k+\frac{1}{2})}$  such that  $(\mathbf{u}, \theta) = (I - \mathbf{P}_N) \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})$ . Using Theorem 1, Corollary 3 and (19), we obtain

$$\begin{split} \|\mathbf{u}\|_{0} &= \left\| (I-P_{N})\Pi_{1}\mathscr{S}\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \right\|_{0} \leq \frac{C_{0}^{2}}{(N+1)^{2}} \|\bar{\mathbf{f}}(\bar{\mathbf{u}},\bar{\theta})\|_{0} \leq \frac{C_{0}^{2}}{(N+1)^{2}} m_{1}^{(k+1)} < \frac{C_{0}^{2}}{(N+1)^{2}} m_{1}^{(k+\frac{1}{2})}, \\ \|\nabla\mathbf{u}\|_{0} &= \left\| \nabla(I-P_{N})\Pi_{1}\mathscr{S}\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \right\|_{0} \leq \frac{C_{0}}{N+1} \|\bar{\mathbf{f}}(\bar{\mathbf{u}},\bar{\theta})\|_{0} \leq \frac{C_{0}}{N+1} m_{1}^{(k+1)} < \frac{C_{0}}{N+1} m_{1}^{(k+\frac{1}{2})}, \\ \|\mathbf{u}\|_{\infty} &= \left\| (I-P_{N})\Pi_{1}\mathscr{S}\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \right\|_{\infty} \leq \frac{2C_{1}}{\sqrt{N}} \|\bar{\mathbf{f}}(\bar{\mathbf{u}},\bar{\theta})\|_{0} \leq \frac{2C_{1}}{\sqrt{N}} m_{1}^{(k+1)} < \frac{2C_{1}}{\sqrt{N}} m_{1}^{(k+\frac{1}{2})}, \\ \|\theta\|_{0} &= \left\| (I-Q_{N})\Pi_{2}\mathscr{S}\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \right\|_{0} \leq \frac{C_{0}^{2}}{(N+1)^{2}} \|\bar{g}(\bar{\mathbf{u}},\bar{\theta})\|_{0} \leq \frac{C_{0}^{2}}{(N+1)^{2}} m_{2}^{(k+1)} < \frac{C_{0}^{2}}{(N+1)^{2}} m_{2}^{(k+\frac{1}{2})}, \\ \|\nabla\theta\|_{0} &= \left\| \nabla(I-Q_{N})\Pi_{2}\mathscr{S}\bar{F}(\bar{\mathbf{u}},\bar{\theta}) \right\|_{0} \leq \frac{C_{0}}{N+1} \|\bar{g}(\bar{\mathbf{u}},\bar{\theta})\|_{0} \leq \frac{C_{0}}{N+1} m_{2}^{(k+1)} < \frac{C_{0}}{N+1} m_{2}^{(k+\frac{1}{2})}, \end{split}$$

where  $\Pi_1: X \to V$  and  $\Pi_2: X \to W$  are the natural projections from X to V and W, respectively. These estimates mean that  $(\mathbf{u}, \boldsymbol{\theta}) \in U_*^{(k+\frac{1}{2})}$  and  $(I - \mathbf{P}_N) \mathscr{K} U^{(k+\frac{1}{2})} \subset U_*^{(k+\frac{1}{2})}$  holds.  $\Box$ 

To determine the finite dimensional set  $U_N^{(k+1)}$  in (19), we need to compute  $\mathcal{N}$  on U. At first, from definitions of  $\mathcal{L}_N$  and  $\mathcal{K}$ , we can rewrite  $\mathcal{N}$  as follows:

$$\mathscr{N} = \mathscr{L}_N^{-1} \left[ \mathscr{L}_N \mathbf{P}_N - \mathbf{P}_N + \mathbf{P}_N \mathscr{K} \right] = \mathscr{L}_N^{-1} \mathbf{P}_N \mathscr{S} F_0, \quad F_0 \equiv \bar{F} - F'(\mathbf{u}_N, \theta_N) \mathbf{P}_N.$$

For any fixed  $(\bar{\mathbf{u}}, \bar{\theta}) \in U$ , set  $(\mathbf{u}_h, \theta_h) \equiv \mathcal{N}(\bar{\mathbf{u}}, \bar{\theta})$  and operate  $\mathcal{L}_N$  on both sides, then

$$\mathbf{P}_{N}\mathscr{SL}_{0}(\mathbf{u}_{h},\boldsymbol{\theta}_{h})=\mathscr{L}_{N}(\mathbf{u}_{h},\boldsymbol{\theta}_{h})=\mathbf{P}_{N}\mathscr{SF}_{0}(\bar{\mathbf{u}},\bar{\boldsymbol{\theta}})$$

Using the projection property (13) of  $\mathbf{P}_N$ , we can derive

$$\langle \nabla \mathscr{SL}_0(\mathbf{u}_h, \mathbf{\theta}_h), \nabla(\mathbf{v}_N, \mathbf{\vartheta}_N) \rangle = \langle \nabla \mathscr{SF}_0(\bar{\mathbf{u}}, \bar{\mathbf{\theta}}), \nabla(\mathbf{v}_N, \mathbf{\vartheta}_N) \rangle, \quad \forall (\mathbf{v}_N, \mathbf{\vartheta}_N) \in X_N$$

Due to (10), this can be written as

$$\mathscr{L}_{0}(\mathbf{u}_{h},\boldsymbol{\theta}_{h}),(\mathbf{v}_{N},\vartheta_{N})\rangle = \langle F_{0}(\bar{\mathbf{u}},\bar{\boldsymbol{\theta}}),(\mathbf{v}_{N},\vartheta_{N})\rangle, \quad \forall (\mathbf{v}_{N},\vartheta_{N}) \in X_{N}.$$
(21)

The left hand side of (21) can be written as:

$$\langle \mathscr{L}_0(\mathbf{u}_h, \theta_h), (\mathbf{v}_N, \vartheta_N) \rangle = \langle \mathscr{S}^{-1}(\mathbf{u}_h, \theta_h) - F'(\mathbf{u}_N, \theta_N)(\mathbf{u}_h, \theta_h), (\mathbf{v}_N, \vartheta_N) \rangle = \langle \nabla(\mathbf{u}_h, \theta_h), \nabla(\mathbf{v}_N, \vartheta_N) \rangle - \langle F'(\mathbf{u}_N, \theta_N)(\mathbf{u}_h, \theta_h), (\mathbf{v}_N, \vartheta_N) \rangle.$$

This gives us the interval version of the Jacobian matrix in Newton's method with respect to the base functions  $\Phi^{\alpha}$ ,  $\Psi^{\alpha}$ , and  $\phi_{3}^{\alpha}$ . The right hand side of (21) forms an interval vector whose elements can be enclosed with upper and lower bounds. Thus the operator  $\mathscr{L}_{N}$  is regular when the solution  $(\mathbf{u}_{h}, \theta_{h}) \in X_{N}$  for (21) exists under the guaranteed computation with interval arithmetic.

In order to compute  $m_i^{(k+1)}$  in (19), we need to estimate  $\|\bar{\mathbf{f}}(\mathbf{u},\theta)\|_0$  and  $\|\bar{g}(\mathbf{u},\theta)\|_0$  for any  $(\mathbf{u},\theta) \in U$ . Pick up an element  $(\mathbf{u},\theta) \equiv (\mathbf{u}_h + \mathbf{u}_*, \theta_h + \theta_*) \in U$  with  $(\mathbf{u}_h, \theta_h) \in U_N$ ,  $(\mathbf{u}_*, \theta_*) \in U_*$ , then we have

$$\begin{aligned} \mathbf{f}(\mathbf{u},\boldsymbol{\theta}) &= \mathbf{f}(\mathbf{u}_N + \mathbf{u}_h + \mathbf{u}_*, \theta_N + \theta_h + \theta_*) + \Delta \mathbf{u}_N - \nabla p_N \\ &= \mathbf{f}(\mathbf{u}_N + \mathbf{u}_h, \theta_N + \theta_h) + \Delta \mathbf{u}_N - \nabla p_N \\ &- \frac{1}{\mathscr{P}} \left[ ((\mathbf{u}_N + \mathbf{u}_h) \cdot \nabla) \mathbf{u}_* + (\mathbf{u}_* \cdot \nabla) (\mathbf{u}_N + \mathbf{u}_h) + (\mathbf{u}_* \cdot \nabla) \mathbf{u}_* \right] + \mathscr{R} \theta_* \mathbf{e}_2 \\ \bar{g}(\mathbf{u},\boldsymbol{\theta}) &= g(\mathbf{u}_N + \mathbf{u}_h + \mathbf{u}_*, \theta_N + \theta_h + \theta_*) + \Delta \theta_N \\ &= g(\mathbf{u}_N + \mathbf{u}_h, \theta_N + \theta_h) + \Delta \theta_N \\ &- \left[ ((\mathbf{u}_N + \mathbf{u}_h) \cdot \nabla) \theta_* + (\mathbf{u}_* \cdot \nabla) (\theta_N + \theta_h) + (\mathbf{u}_* \cdot \nabla) \theta_* \right] + w_*. \end{aligned}$$

These forms enable us to estimate the desired norms as follows:

$$\begin{split} \left\| \mathbf{f}(\mathbf{u}, \theta) \right\|_{0} &\leq \left\| \mathbf{f}(\mathbf{u}_{N} + \mathbf{u}_{h}, \theta_{N} + \theta_{h}) + \Delta \mathbf{u}_{N} - \nabla p_{N} \right\|_{0} \\ &+ \frac{1}{\mathscr{P}} \left[ \left\| \mathbf{u}_{N} + \mathbf{u}_{h} \right\|_{\infty} \left\| \nabla \mathbf{u}_{*} \right\|_{0} + \left\| \mathbf{u}_{*} \right\|_{0} \| \nabla (\mathbf{u}_{N} + \mathbf{u}_{h}) \right\|_{\infty} + \left\| \mathbf{u}_{*} \right\|_{\infty} \| \nabla \mathbf{u}_{*} \|_{0} \right] + \mathscr{R} \| \theta_{*} \|_{0} \\ &\leq \left\| \mathbf{f}(\mathbf{u}_{N} + \mathbf{u}_{h}, \theta_{N} + \theta_{h}) + \Delta \mathbf{u}_{N} - \nabla p_{N} \right\|_{0} \\ &+ \frac{1}{\mathscr{P}} \left[ \frac{C_{0}}{N+1} \| \mathbf{u}_{N} + \mathbf{u}_{h} \|_{\infty} + \frac{C_{0}^{2}}{(N+1)^{2}} \| \nabla (\mathbf{u}_{N} + \mathbf{u}_{h}) \|_{\infty} \right] m_{1} + \frac{2C_{0}C_{1}}{(N+1)\sqrt{N}} m_{1}^{2} + \mathscr{R} \frac{C_{0}^{2}}{(N+1)^{2}} m_{2}, \\ \left\| \bar{g}(\mathbf{u}, \theta) \right\|_{0} &\leq \left\| g(\mathbf{u}_{N} + \mathbf{u}_{h}, \theta_{N} + \theta_{h}) + \Delta \theta_{N} \right\|_{0} \\ &+ \left[ \left\| \mathbf{u}_{N} + \mathbf{u}_{h} \right\|_{\infty} \| \nabla \theta_{*} \|_{0} + \left\| \mathbf{u}_{*} \right\|_{0} \| \nabla (\theta_{N} + \theta_{h}) \|_{\infty} + \left\| \mathbf{u}_{*} \right\|_{\infty} \| \nabla \theta_{*} \|_{0} \right] + \left\| w_{*} \right\|_{0} \\ &\leq \left\| g(\mathbf{u}_{N} + \mathbf{u}_{h}, \theta_{N} + \theta_{h}) + \Delta \theta_{N} \right\|_{0} \\ &+ \frac{C_{0}}{N+1} \left\| \mathbf{u}_{N} + \mathbf{u}_{h} \right\|_{\infty} m_{2} + \frac{C_{0}^{2}}{(N+1)^{2}} \| \nabla (\theta_{N} + \theta_{h}) \|_{\infty} m_{1} + \frac{2C_{0}C_{1}}{(N+1)\sqrt{N}} m_{1}m_{2} + \frac{C_{0}^{2}}{(N+1)^{2}} m_{1}. \end{split}$$

Note that upper bounds of  $L^2$  and  $L^{\infty}$  norms for  $(\mathbf{u}_N + \mathbf{u}_h, \theta_N + \theta_h) \in U_N \subset X_N$  can be computed by interval arithmetic, and these calculation may have additional inflations due to crude estimates. Thus we estimate  $\|(I - P_N)\Pi_1 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})\|_0$ ,  $\|\nabla(I - P_N)\Pi_1 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})\|_0$ ,  $\|(I - P_N)\Pi_1 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})\|_{\infty}$   $\|(I - Q_N)\Pi_2 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})\|_0$ , and  $\|\nabla(I - Q_N)\Pi_2 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})\|_0$  after decomposition of  $\Pi_1 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})$  and  $\Pi_2 \mathscr{S}\bar{F}(\bar{\mathbf{u}}, \bar{\theta})$  into finite and infinite parts, which gives us more accurate values of them and efficient estimates in real computations.

#### **5** Numerical results

For the interval arithmetic, we use the PROFIL package [2] on Linux Intel Pentium 4 (3.8 GHz) machine.

We set  $a^2 = \frac{1}{8}$ ,  $b^2 = \frac{3}{8}$  and  $\mathscr{P} = 10$  in the numerical experiments with 1% inflation factor. Then the critical Rayleigh number  $\mathscr{R}_c = 6.75$  can be attained at some special mode  $\alpha$  (see [5] for detail). We show several approximation results in figures 1-3. In these figures, the isothermal lines are drawn after adding the conduction solution (2) on the left, and contour lines of speed with streamlines are shown on the right. Note that streamlines for each type never change their shape during the change of relative Rayleigh numbers in short range. We present figures at the same relative Rayleigh number  $\mathscr{R}/\mathscr{R}_c = 1.1$  for easy comparison.



Fig. 1 Isothermal lines, and contour lines of speed with streamlines for roll type at  $\Re/\Re_c = 1.1$ .



Fig. 2 Isothermal lines, and contour lines of speed with streamlines for rectangular type at  $\Re/\Re_c = 1.1$ .



Fig. 3 Isothermal lines, and contour lines of speed with streamlines for hexagonal type at  $\Re/\Re_c = 1.1$ .

In Table 1, Table 2 and Table 3, we illustrate the verification results for each type of solutions with several relative Rayleigh numbers. In these tables, we show the relative Rayleigh number  $\mathscr{R}/\mathscr{R}_c$ , the truncation number N, the converged step k,  $L^{\infty}$  norms of approximate solutions  $(\mathbf{u}_N, \theta_N)$ ,  $L^{\infty}$  norms of finite parts  $(\mathbf{u}_h, \theta_h)$ , and the bounds  $m_1$ ,  $m_2$  of infinite parts. The converged step means the inflated candidate set at step  $k - \frac{1}{2}$  includes the new one at step k, namely, the verification was completed at the concerning iteration steps. In the roll type case (Table 1), the problem size becomes much smaller due to the elimination of one space variable which comes from the fact that the solutions are independent of that variable. For other types (Table 2, Table 3), we can find out the basic symmetry of solutions which make it possible to reduce the size of unknown coefficients.

From these tables, we can make a bifurcation diagram Fig. 4 with respect to the relative Rayleigh number  $\mathscr{R}/\mathscr{R}_c$  and sum  $\|\nabla \mathbf{u}_N\|_{\infty} + \|\nabla \theta_N\|_{\infty}$  of approximate solutions'  $L^{\infty}$  norms.

#### **6** Conclusion

We could verify several kinds of bifurcating solutions. This should be the first result on the fact that there actually exist exact solutions around approximate solutions drawn in the figures corresponding to the interesting

$\mathcal{R}/\mathcal{R}_c$	Ν	k	$\ \nabla \mathbf{u}_N\ _{\infty}$	$\  abla  heta_N\ _{\infty}$	$\ \nabla \mathbf{u}_h\ _{\infty}$	$\  abla  heta_h\ _{\infty}$	$m_1$	$m_2$
1.01	16	12	0.74	0.31	$5 \times 10^{-11}$	$2 \times 10^{-10}$	$1.44 \times 10^{-10}$	$1.92 \times 10^{-10}$
1.05	18	29	1.66	0.72	$2 \times 10^{-8}$	$2 \times 10^{-8}$	$4.06 \times 10^{-9}$	$3.57 imes10^{-9}$
1.5	44	19	5.58	2.47	$3 \times 10^{-11}$	$3 \times 10^{-10}$	$1.72 \times 10^{-11}$	$3.33 \times 10^{-11}$

Table 1 Verification results for roll type solutions.

$\mathscr{R}/\mathscr{R}_c$	N	k	$\ \nabla \mathbf{u}_N\ _{\infty}$	$\ \nabla \theta_N\ _{\infty}$	$\ \nabla \mathbf{u}_h\ _{\infty}$	$\  abla  heta_h\ _{\infty}$	$m_1$	$m_2$
1.01	16	7	0.93	0.38	$2 \times 10^{-11}$	$3 \times 10^{-11}$	$2.37 \times 10^{-12}$	$1.01 \times 10^{-11}$
1.05	16	10	2.20	0.96	$2 \times 10^{-8}$	$2 \times 10^{-7}$	$2.91 \times 10^{-9}$	$1.09  imes 10^{-8}$
1.1	24	11	3.26	1.47	$4 \times 10^{-10}$	$3 \times 10^{-9}$	$8.33 \times 10^{-13}$	$5.84  imes 10^{-12}$
1.2	28	33	4.99	2.27	$2 \times 10^{-10}$	$2 \times 10^{-9}$	$4.60 \times 10^{-12}$	$3.96  imes 10^{-11}$

 Table 2
 Verification results for rectangular type solutions.

$\mathcal{R}/\mathcal{R}_c$	N	k	$\ \nabla \mathbf{u}_N\ _{\infty}$	$\ \nabla \theta_N\ _{\infty}$	$\ \nabla \mathbf{u}_h\ _{\infty}$	$\  abla  heta_h\ _{\infty}$	$m_1$	<i>m</i> <sub>2</sub>
1.01	16	7	1.12	0.47	$2 \times 10^{-10}$	$6 \times 10^{-10}$	$4.53 \times 10^{-11}$	$1.93  imes 10^{-10}$
1.05	16	15	2.70	1.21	$2 \times 10^{-6}$	$2 \times 10^{-6}$	$3.22  imes 10^{-8}$	$1.08  imes 10^{-7}$
1.1	24	19	4.09	1.88	$4 \times 10^{-9}$	$3 \times 10^{-8}$	$2.50 \times 10^{-11}$	$1.68  imes 10^{-10}$

Table 3 Verification results for hexagonal type solutions.



Fig. 4 Bifurcation diagram. × for hexagonal, o for rectangular, and \* for roll type cases.

natural phenomena. Due to the limit of our computational power up to now, we proved only small part (Fig. 4) of bifurcation diagram except for the roll type solutions. This could be enhanced using parallel computation after update of PROFIL package into parallel version (not available now) or increasing physical memories for computation on some large scale computers.

More interesting problem would be the verification of the bifurcation point such that suggested in [5], which is more complicated problem than the usual. We believe that these interesting and important problems could be resolved in the near future.

#### References

- Fabrice Caini and Philippe Gelene, Rayleigh-benard convection, http://www.enseeiht.fr/hmf/travaux/CD9598/ travaux/optmfn/BES\_PHOENICS/96-97/Caini-Gelene/RB\_convectio\_tur.html; accessed Dec 2, 2006.
- Siegfried M. Rump (Head), Profil. http://www.ti3.tu-harburg.de/Software/PROFILEnglisch.html; accessed Dec 4, 2006, Institute for Reliable Computing.
- 3. Paul Manneville, Rayleigh-Bénard convection, thirty years of experimental, theoretical, and modeling work, http://www.ladhyx.polytechnique.fr/people/pops/benard.pdf; accessed Feb 16, 2005.

- 4. Mitsuhiro T. Nakao, Yoshitaka Watanabe, Nobito Yamamoto, and Takaaki Nishida, Some computer assisted proofs for solutions of the heat convection problems, Reliable Computing 9 (2003), no. 5, 359–372.
  Takaaki Nishida, Tsutomu Ikeda, and Hideaki Yoshihara, *Mathematical modeling and numerical simulation in continuum*
- mechanics, Lecture Notes in Computational Sciences and Engineering, vol. 19, ch. Pattern Formation of Heat Convection Problems, pp. 209–218, Springer, 2002.
  Michael Plum, *Explicit H<sub>2</sub>-estimates and pointwise bounds for solutions of second-order elliptic boundary value problems*,
- Journal of Mathematical Analysis and Applications 165 (1992), 36-61.
- 7. Yoshitaka Watanabe, Nobito Yamamoto, Mistuhiro T. Nakao, and Takaaki Nishida, *A numerical verification of nontrivial solutions for the heat convection problem*, Journal of Mathematical Fluid Mechanics **6** (2004), 1–20.

# List of MHF Preprint Series, Kyushu University 21st Century COE Program Development of Dynamic Mathematics with High Functionality

- MHF2005-1 Hideki KOSAKI Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA Simple estimators for non-linear Markovian trend from sampled data: I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA Estimation for a discretely observed small diffusion process with a linear drift
- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA AIC for ergodic diffusion processes from discrete observations

- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA Integration by parts formulae for the Wiener measure restricted to subsets in  $\mathbb{R}^d$
- MHF2005-21 Yuu HARIYA A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI Nonlinear regression modeling via regularized wavelets and smoothing parameter selection
- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO, Yoshihiro MIZOGUCHI & Yasuo KAWAHARA On reversible cellular automata with finite cell array

- MHF2005-28 Toru KOMATSU Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI An explicit formula of the Shapley value for a cooperative game induced from the conjugate point
- MHF2005-34 Hideki MURAKAWA A regularization of a reaction-diffusion system approximation to the two-phase Stefan problem
- MHF2006-1 Masahisa TABATA Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme
- MHF2006-2 Ken-ichi MARUNO & G R W QUISPEL Construction of integrals of higher-order mappings
- MHF2006-3 Setsuo TANIGUCHI On the Jacobi field approach to stochastic oscillatory integrals with quadratic phase function
- MHF2006-4 Kouji HASHIMOTO, Kaori NAGATOU & Mitsuhiro T. NAKAO A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains
- MHF2006-5 Hidefumi KAWASAKI A duality theory based on triangular cylinders separating three convex sets in  $\mathbb{R}^n$
- MHF2006-6 Raimundas VIDŪNAS Uniform convergence of hypergeometric series
- MHF2006-7 Yuji KODAMA & Ken-ichi MARUNO N-Soliton solutions to the DKP equation and Weyl group actions

- MHF2006-8 Toru KOMATSU Potentially generic polynomial
- MHF2006-9 Toru KOMATSU Generic sextic polynomial related to the subfield problem of a cubic polynomial
- MHF2006-10 Shu TEZUKA & Anargyros PAPAGEORGIOU Exact cubature for a class of functions of maximum effective dimension
- MHF2006-11 Shu TEZUKA On high-discrepancy sequences
- MHF2006-12 Raimundas VIDŪNAS Detecting persistent regimes in the North Atlantic Oscillation time series
- MHF2006-13 Toru KOMATSU Tamely Eisenstein field with prime power discriminant
- MHF2006-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO Generating function associated with the Hankel determinant formula for the solutions of the Painlevé IV equation
- MHF2006-15 Raimundas VIDŪNAS Darboux evaluations of algebraic Gauss hypergeometric functions
- MHF2006-16 Masato KIMURA & Isao WAKANO New mathematical approach to the energy release rate in crack extension
- MHF2006-17 Toru KOMATSU Arithmetic of the splitting field of Alexander polynomial
- MHF2006-18 Hiroki MASUDA Likelihood estimation of stable Lévy processes from discrete data
- MHF2006-19 Hiroshi KAWABI & Michael RÖCKNER Essential self-adjointness of Dirichlet operators on a path space with Gibbs measures via an SPDE approach
- MHF2006-20 Masahisa TABATA Energy stable finite element schemes and their applications to two-fluid flow problems
- MHF2006-21 Yuzuru INAHAMA & Hiroshi KAWABI Asymptotic expansions for the Laplace approximations for Itô functionals of Brownian rough paths
- MHF2006-22 Yoshiyuki KAGEI Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

MHF2006-23 Yoshiyuki KAGEI Asymptotic behavior of the semigroup associated with the linearized compressible Navier-Stokes equation in an infinite layer

- MHF2006-24 Akihiro MIKODA, Shuichi INOKUCHI, Yoshihiro MIZOGUCHI & Mitsuhiko FUJIO The number of orbits of box-ball systems
- MHF2006-25 Toru FUJII & Sadanori KONISHI Multi-class logistic discrimination via wavelet-based functionalization and model selection criteria
- MHF2006-26 Taro HAMAMOTO, Kenji KAJIWARA & Nicholas S. WITTE Hypergeometric solutions to the q-Painlevé equation of type  $(A_1 + A'_1)^{(1)}$
- MHF2006-27 Hiroshi KAWABI & Tomohiro MIYOKAWA The Littlewood-Paley-Stein inequality for diffusion processes on general metric spaces
- MHF2006-28 Hiroki MASUDA Notes on estimating inverse-Gaussian and gamma subordinators under highfrequency sampling
- MHF2006-29 Setsuo TANIGUCHI The heat semigroup and kernel associated with certain non-commutative harmonic oscillators
- MHF2006-30 Setsuo TANIGUCHI Stochastic analysis and the KdV equation
- MHF2006-31 Masato KIMURA, Hideki KOMURA, Masayasu MIMURA, Hidenori MIYOSHI, Takeshi TAKAISHI & Daishin UEYAMA Quantitative study of adaptive mesh FEM with localization index of pattern
- MHF2007-1 Taro HAMAMOTO & Kenji KAJIWARA Hypergeometric solutions to the q-Painlevé equation of type  $A_4^{(1)}$
- MHF2007-2 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO Verified numerical computation of solutions for the stationary Navier-Stokes equation in nonconvex polygonal domains
- MHF2007-3 Kenji KAJIWARA, Marta MAZZOCCO & Yasuhiro OHTA A remark on the Hankel determinant formula for solutions of the Toda equation
- MHF2007-4 Jun-ichi SATO & Hidefumi KAWASAKI Discrete fixed point theorems and their application to Nash equilibrium
- MHF2007-5 Mitsuhiro T. NAKAO & Kouji HASHIMOTO Constructive error estimates of finite element approximations for non-coercive elliptic problems and its applications

- MHF2007-6 Kouji HASHIMOTO A preconditioned method for saddle point problems
- MHF2007-7 Christopher MALON, Seiichi UCHIDA & Masakazu SUZUKI Mathematical symbol recognition with support vector machines
- MHF2007-8 Kenta KOBAYASHI On the global uniqueness of Stokes' wave of extreme form
- MHF2007-9 Kenta KOBAYASHI A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions
- MHF2007-10 Myoungnyoun KIM, Mitsuhiro T. NAKAO, Yoshitaka WATANABE & Takaaki NISHIDA

A numerical verification method of bifurcating solutions for 3-dimensional Rayleigh-Bénard problems