

Measurement of prostaglandin metabolites is useful in diagnosis of small bowel ulcerations

松野, 雄一

<https://hdl.handle.net/2324/4060049>

出版情報 : Kyushu University, 2019, 博士 (医学), 課程博士

バージョン :

権利関係 : ©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

World Journal of *Gastroenterology*

World J Gastroenterol 2019 April 14; 25(14): 1640-1782



REVIEW

- 1640 Precision surgical approach with lymph-node dissection in early gastric cancer
Kinami S, Nakamura N, Tomita Y, Miyata T, Fujita H, Ueda N, Kosaka T

MINIREVIEWS

- 1653 Update on hepatocellular carcinoma: Pathologists' review
El Jabbour T, Lagana SM, Lee H
- 1666 Application of artificial intelligence in gastroenterology
Yang YJ, Bang CS

ORIGINAL ARTICLE**Basic Study**

- 1684 Clinical significance of programmed death 1/programmed death ligand 1 pathway in gastric neuroendocrine carcinomas
Yang MW, Fu XL, Jiang YS, Chen XJ, Tao LY, Yang JY, Huo YM, Liu W, Zhang JF, Liu PF, Liu Q, Hua R, Zhang ZG, Sun YW, Liu DJ
- 1697 Functional role of long non-coding RNA CASC19/miR-140-5p/CEMIP axis in colorectal cancer progression *in vitro*
Wang XD, Lu J, Lin YS, Gao C, Qi F
- 1715 Seven-senescence-associated gene signature predicts overall survival for Asian patients with hepatocellular carcinoma
Xiang XH, Yang L, Zhang X, Ma XH, Miao RC, Gu JX, Fu YN, Yao Q, Zhang JY, Liu C, Lin T, Qu K

Case Control Study

- 1729 Chronic functional constipation is strongly linked to vitamin D deficiency
Panarese A, Pesce F, Porcelli P, Riezzo G, Iacovazzi PA, Leone CM, De Carne M, Rinaldi CM, Shahini E

Retrospective Study

- 1741 Clinical characteristics of non-alcoholic fatty liver disease in Chinese adult hypopituitary patients
Yuan XX, Zhu HJ, Pan H, Chen S, Liu ZY, Li Y, Wang LJ, Lu L, Yang HB, Gong FY

Observational Study

- 1753 Measurement of prostaglandin metabolites is useful in diagnosis of small bowel ulcerations
Matsuno Y, Umeno J, Esaki M, Hirakawa Y, Fuyuno Y, Okamoto Y, Hirano A, Yasukawa S, Hirai F, Matsui T, Hosomi S, Watanabe K, Hosoe N, Ogata H, Hisamatsu T, Yanai S, Kochi S, Kurahara K, Yao T, Torisu T, Kitazono T, Matsumoto T

Prospective Study

- 1764** Endoscopic response to tumor necrosis factor inhibitors predicts long term benefits in Crohn's disease
Alfaro I, Masamunt MC, Planell N, López-García A, Castro J, Gallego M, Barastegui R, Giner A, Vara A, Salas A, Ricart E, Panés J, Ordás I

CASE REPORT

- 1775** Application of a 3D-printed "fistula stent" in plugging enteroatmospheric fistula with open abdomen: A case report
Xu ZY, Ren HJ, Huang JJ, Li ZA, Ren JA

ABOUT COVER

Editorial board member of *World Journal of Gastroenterology*, Jun-Te Hsu, MD, Professor, Surgeon, Surgical Oncologist, Department of Surgery, Chang Gung Memorial Hospital, Taoyuan County and Chang Gung Medicine Colledge, Tao Yuan 333, Taiwan

AIMS AND SCOPE

World Journal of Gastroenterology (*World J Gastroenterol*, *WJG*, print ISSN 1007-9327, online ISSN 2219-2840, DOI: 10.3748) is a peer-reviewed open access journal. The *WJG* Editorial Board consists of 642 experts in gastroenterology and hepatology from 59 countries.

The primary task of *WJG* is to rapidly publish high-quality original articles, reviews, and commentaries in the fields of gastroenterology, hepatology, gastrointestinal endoscopy, gastrointestinal surgery, hepatobiliary surgery, gastrointestinal oncology, gastrointestinal radiation oncology, etc. The *WJG* is dedicated to become an influential and prestigious journal in gastroenterology and hepatology, to promote the development of above disciplines, and to improve the diagnostic and therapeutic skill and expertise of clinicians.

INDEXING/ABSTRACTING

The *WJG* is now indexed in Current Contents®/Clinical Medicine, Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports®, Index Medicus, MEDLINE, PubMed, PubMed Central, Scopus and Directory of Open Access Journals. The 2018 edition of Journal Citation Report® cites the 2017 impact factor for *WJG* as 3.300 (5-year impact factor: 3.387), ranking *WJG* as 35th among 80 journals in gastroenterology and hepatology (quartile in category Q2).

RESPONSIBLE EDITORS FOR THIS ISSUE

Responsible Electronic Editor: *Han Song*

Proofing Editorial Office Director: *Ze-Mao Gong*

NAME OF JOURNAL

World Journal of Gastroenterology

ISSN

ISSN 1007-9327 (print) ISSN 2219-2840 (online)

LAUNCH DATE

October 1, 1995

FREQUENCY

Weekly

EDITORS-IN-CHIEF

Subrata Ghosh, Andrzej S Tarnawski

EDITORIAL BOARD MEMBERS

<http://www.wjgnet.com/1007-9327/editorialboard.htm>

EDITORIAL OFFICE

Ze-Mao Gong, Director

PUBLICATION DATE

April 14, 2019

COPYRIGHT

© 2019 Baishideng Publishing Group Inc

INSTRUCTIONS TO AUTHORS

<https://www.wjgnet.com/bpg/gerinfo/204>

GUIDELINES FOR ETHICS DOCUMENTS

<https://www.wjgnet.com/bpg/GerInfo/287>

GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH

<https://www.wjgnet.com/bpg/gerinfo/240>

PUBLICATION MISCONDUCT

<https://www.wjgnet.com/bpg/gerinfo/208>

ARTICLE PROCESSING CHARGE

<https://www.wjgnet.com/bpg/gerinfo/242>

STEPS FOR SUBMITTING MANUSCRIPTS

<https://www.wjgnet.com/bpg/GerInfo/239>

ONLINE SUBMISSION

<https://www.f6publishing.com>

Observational Study

Measurement of prostaglandin metabolites is useful in diagnosis of small bowel ulcerations

Yuichi Matsuno, Junji Umeno, Motohiro Esaki, Yoichiro Hirakawa, Yuta Fuyuno, Yasuharu Okamoto, Atsushi Hirano, Shigeyoshi Yasukawa, Fumihito Hirai, Toshiyuki Matsui, Shuhei Hosomi, Kenji Watanabe, Naoki Hosoe, Haruhiko Ogata, Tadakazu Hisamatsu, Shunichi Yanai, Shuji Kochi, Koichi Kurahara, Tsuneyoshi Yao, Takehiro Torisu, Takanari Kitazono, Takayuki Matsumoto

ORCID number: Yuichi Matsuno (0000-0002-1973-2573); Junji Umeno (0000-0002-5253-5600); Motohiro Esaki (0000-0002-1195-0074); Yoichiro Hirakawa (0000-0002-4405-4399); Yuta Fuyuno (0000-0003-3527-5648); Yasuharu Okamoto (0000-0001-5693-8565); Atsushi Hirano (0000-0002-4791-4550); Shigeyoshi Yasukawa (0000-0002-7421-2706); Fumihito Hirai (0000-0002-5493-5675); Toshiyuki Matsui (0000-0002-1817-8064); Shuhei Hosomi (0000-0002-8808-5672); Kenji Watanabe (0000-0002-3781-4724); Naoki Hosoe (0000-0002-4516-5648); Haruhiko Ogata (0000-0002-3304-3635); Tadakazu Hisamatsu (0000-0002-1178-3536); Shunichi Yanai (0000-0003-1871-2412); Shuji Kochi (0000-0002-1783-9545); Koichi Kurahara (0000-0002-6115-341X); Tsuneyoshi Yao (0000-0002-7381-4641); Takehiro Torisu (0000-0001-8256-2549); Takanari Kitazono (0000-0002-9006-5937); Takayuki Matsumoto (0000-0001-9786-3854).

Author contributions: All authors helped to perform the research; Umeno J, Esaki M and Matsumoto T conceived the study; Matsuno Y and Hirakawa Y contributed to statistical analysis and data interpretation; Matsuno Y, Umeno J, Torisu T and Esaki M drafted the manuscript; Matsuno Y, Umeno J, Esaki M, Fuyuno Y, Okamoto Y, Hirano A, Yasukawa S, Hirai F,

Yuichi Matsuno, Junji Umeno, Yoichiro Hirakawa, Yuta Fuyuno, Yasuharu Okamoto, Atsushi Hirano, Takehiro Torisu, Takanari Kitazono, Department of Medicine and Clinical Science, Graduate School of Medical Sciences, Kyushu University, Fukuoka 812-8582, Japan

Motohiro Esaki, Department of Endoscopic Diagnostics and Therapeutic, Saga University Hospital, Saga 849-8501, Japan

Yoichiro Hirakawa, Department of Epidemiology and Public Health, Graduate School of Medical Sciences, Kyushu University, Fukuoka 812-8582, Japan

Shigeyoshi Yasukawa, Fumihito Hirai, Toshiyuki Matsui, Department of Gastroenterology, Fukuoka University Chikushi Hospital, Chikushino 818-8502, Japan

Shuhei Hosomi, Department of Gastroenterology, Osaka City University Graduate School of Medicine, Osaka 545-8586, Japan

Kenji Watanabe, Department of Intestinal Inflammation Research, Hyogo College of Medicine, Nishinomiya 663-8501, Japan

Naoki Hosoe, Haruhiko Ogata, Center for Diagnostic and Therapeutic Endoscopy, School of Medicine, Keio University, Tokyo 160-0016, Japan

Tadakazu Hisamatsu, the Third Department of Internal Medicine, Kyorin University School of Medicine, Mitaka 181-8611, Japan

Shunichi Yanai, Takayuki Matsumoto, Division of Gastroenterology, Department of Internal Medicine, Iwate Medical University, Morioka 020-8505, Japan

Shuji Kochi, Koichi Kurahara, Division of Gastroenterology, Matsuyama Red Cross Hospital, Matsuyama 790-8524, Japan

Tsuneyoshi Yao, Department of Gastroenterology, Sada Hospital, Fukuoka 810-0004, Japan

Corresponding author: Takayuki Matsumoto, MD, PhD, Professor, Division of Gastroenterology, Department of Internal Medicine, Iwate Medical University, 19-1 Uchimaru, Morioka 020-8505, Japan. tmatsumo@iwate-med.ac.jp

Telephone: +81-196515111 (ext. 2314)

Fax: +81-196526664

Matsui T, Hosomi S, Watanabe K, Hosoe N, Ogata H, Hisamatsu T, Yanai S, Kochi S, Kurahara K, Yao T, and Torisu T were responsible for recruitment and phenotyping of the patients; Kitazono T and Matsumoto T revised the manuscript; all authors approved the final version of the manuscript.

Supported by the Practical Research Project for Rare/Intractable Diseases from the Japan Agency for Medical Research and Development (AMED), No. 15ek0109053h0002 to Matsumoto T, and by grants from the Japan Society for the Promotion of Science (JSPS) KAKENHI, No. 25460953, to Umeno J, Esaki M, and Matsumoto T.

Institutional review board

statement: This study was approved by the institutional review board at each collecting site in accordance with the Declaration of Helsinki Principles.

Informed consent statement: All urine and blood samples were collected after obtaining written informed consent.

Conflict-of-interest statement: All authors declare no conflicts-of-interest related to this article.

Data sharing statement: No additional data are available.

STROBE statement: The authors have read the STROBE Statement checklist of items, and the manuscript was prepared according to the STROBE Statement-checklist of items.

Open-Access: This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Received: February 3, 2019

Peer-review started: February 6, 2019

First decision: February 21, 2019

Revised: February 27, 2019

Accepted: March 11, 2019

Article in press: March 12, 2019

Abstract

BACKGROUND

We recently reported on a hereditary enteropathy associated with a gene encoding a prostaglandin transporter and referred to as chronic enteropathy associated with *SLCO2A1* gene (CEAS). Crohn's disease (CD) is a major differential diagnosis of CEAS, because these diseases share some clinical features. Therefore, there is a need to develop a convenient screening test to distinguish CEAS from CD.

AIM

To examine whether prostaglandin E major urinary metabolites (PGE-MUM) can serve as a biomarker to distinguish CEAS from CD.

METHODS

This was a transactional study of 20 patients with CEAS and 98 patients with CD. CEAS was diagnosed by the confirmation of homozygous or compound heterozygous mutation of *SLCO2A1*. We measured the concentration of PGE-MUM in spot urine by radioimmunoassay, and the concentration was compared between the two groups of patients. We also determined the optimal cut-off value of PGE-MUM to distinguish CEAS from CD by receiver operating characteristic (ROC) curve analysis.

RESULTS

Twenty Japanese patients with CEAS and 98 patients with CD were enrolled. PGE-MUM concentration in patients with CEAS was significantly higher than that in patients with CD (median 102.7 vs 27.9 $\mu\text{g/g} \times \text{Cre}$, $P < 0.0001$). One log unit increase in PGE-MUM contributed to 7.3 increase in the likelihood for the diagnosis of CEAS [95% confidence interval (CI) 3.2-16.7]. A logistic regression analysis revealed that the association was significant even after adjusting confounding factors (adjusted odds ratio 29.6, 95%CI 4.7-185.7). ROC curve analysis revealed the optimal PGE-MUM cut-off value for the distinction of CEAS from CD to be 48.9 $\mu\text{g/g} \times \text{Cre}$ with 95.0% sensitivity and 79.6% specificity.

CONCLUSION

PGE-MUM measurement is a convenient, non-invasive and useful test for the distinction of CEAS from CD.

Key words: Chronic enteropathy associated with *SLCO2A1* gene; Prostaglandin E major urinary metabolites; Chronic nonspecific multiple ulcers of the small intestine; Crohn's disease; Small intestine

©The Author(s) 2019. Published by Baishideng Publishing Group Inc. All rights reserved.

Core tip: Chronic enteropathy associated with *SLCO2A1* gene (CEAS) is a rare hereditary enteropathy associated with a gene encoding a prostaglandin transporter. It is sometimes difficult to distinguish CEAS from Crohn's disease (CD), because these diseases share some clinical features. We report the usefulness of prostaglandin E-major urinary metabolites (PGE-MUM) to differentiate CEAS from CD. PGE-MUM concentration was significantly higher in CEAS than in CD, and optimal cut-off value for the distinction of CEAS from CD to be 48.9 $\mu\text{g/g} \times \text{Cre}$ with sensitivity of 95.0% and specificity of 79.6%. In clinical practice, PGE-MUM measurement might be useful as a screening test for CEAS.

Citation: Matsuno Y, Umeno J, Esaki M, Hirakawa Y, Fuyuno Y, Okamoto Y, Hirano A, Yasukawa S, Hirai F, Matsui T, Hosomi S, Watanabe K, Hosoe N, Ogata H, Hisamatsu T, Yanai S, Kochi S, Kurahara K, Yao T, Torisu T, Kitazono T, Matsumoto T. Measurement of prostaglandin metabolites is useful in diagnosis of small bowel ulcerations. *World J Gastroenterol* 2019; 25(14): 1753-1763

URL: <https://www.wjgnet.com/1007-9327/full/v25/i14/1753.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v25.i14.1753>

Published online: April 14, 2019

P-Reviewer: Caboclo JF, Slomiany BL, Tsoulfas G**S-Editor:** Yan JP**L-Editor:** A**E-Editor:** Song H

INTRODUCTION

The use of capsule endoscopy and balloon-assisted endoscopy has enabled the precise observation of small bowel mucosal lesions in gastrointestinal disease pathology, such as Crohn's disease (CD), intestinal tuberculosis, and nonsteroidal anti-inflammatory drug (NSAID)-induced enteropathy^[1-5]. While characteristic endoscopic findings of small bowel inflammation have been reported, it is sometimes difficult to correctly diagnose patients by endoscopic findings alone in cases lacking characteristic features of the disease.

We previously reported an independent disease entity termed chronic nonspecific multiple ulcers of the small intestine (CNSU) characterized by multiple small intestinal ulcers of nonspecific histology^[6-9]. In addition, we recently found that CNSU is caused by recessive inheritance of the *SLCO2A1* gene and proposed a new nomenclature for this disease: Chronic enteropathy associated with *SLCO2A1* gene (CEAS)^[10]. Therefore, it is now possible to diagnose CEAS based on a combination of clinical features and genetic analysis. However, it is sometimes difficult to differentiate CEAS from CD because these diseases share clinical features such as persistent anemia and hypoproteinemia, susceptible age in puberty, and a predominance of ileal involvement. The differentiation between CEAS and CD is clinically essential to avoid ineffective therapies because patients with CEAS do not respond to treatment with corticosteroids or biologics^[7,9]. Moreover, since the patients with CEAS usually show low inflammatory markers in blood test^[11], it is a particularly important problem to distinguish CEAS from CD with low disease activity. In clinical practice, however, it is unrealistic to perform genetic screening for all patients suspected as CD or CEAS.

The *SLCO2A1* gene encodes a prostaglandin transporter that mediates the cellular uptake of prostaglandins. *SLCO2A1* is also the causative gene of primary hypertrophic osteoarthropathy (PHO)^[12-14]. An increase in urinary levels of prostaglandin E2 (PGE2) and prostaglandin E metabolites in *SLCO2A1*-deficient PHO individuals has been confirmed^[14]. PGE2 is known to have the protective effect on the mucosal injury, however it is difficult to directly measure PGE2 level in local tissues or in the blood because of its extremely rapid metabolism^[15]. Prostaglandin E-major urinary metabolite (PGE-MUM) is known to be stable and we have also previously demonstrated an increase in the urinary levels of prostaglandin E metabolites in CEAS patients^[10]. Therefore, this study investigated whether PGE-MUM are useful for differentiating CEAS from CD.

MATERIALS AND METHODS

Subjects

Among the participants of our previous studies^[10,16,17] and the patients diagnosed as CEAS or CD at Kyushu University hospital, we enrolled 20 Japanese patients with CEAS and 98 patients with CD for the present investigation. A diagnosis of CEAS was confirmed based on published clinical criteria^[18,19] with genetic analysis (Supplementary Table 1). The diagnosis of CD was based on the Japanese diagnostic criteria for CD^[20]. No patients were administered prostaglandin analogues or NSAIDs at the time of urine sample collection. All urine and blood samples were collected after obtaining written informed consent. This study was approved by the institutional review board at each collecting site in accordance with the Declaration of Helsinki Principles.

Clinical characteristics

Clinical information including age, gender, history of surgery, concomitant drugs, serum C reactive protein (CRP) levels, disease location, and disease duration was collected from each participating institution. Because the PGE2 metabolite concentration was reported to increase in cases of thiazide-induced hyponatremia^[21] and laxative sennoside administration^[22], we collected information about the use of these medications. Because the PGE-MUM concentration was reported to be influenced by age, gender, smoking habit, and pulmonary inflammatory conditions including chronic fibrosing interstitial pneumonia^[15,23], we also collected information about smoking habit and pulmonary disease. Age was determined at the time of urine sample collection. Smoking habit was defined as never, current (> 6 mo on a daily basis), and former (cessation of smoking > 6 mo). Pulmonary diseases included chronic respiratory disease, diffuse lung disease, and any other lung or airway disease. History of surgery was defined as a history of intestinal resection at the time of study enrolment. The site of involvement was determined by radiographic and/or

Table 1 Clinical characteristics of the study subjects

	CEAS (n = 20)	CD (n = 98)	P value ¹
Age, yr	56 (41-63)	44 (38-54)	0.098
Gender			
Male	5 (25)	66 (67)	0.0008
Female	15 (75)	32 (33)	
Smoking			
Never smoker	16 (80)	68 (69)	0.42
Current smoker	1 (5)	13 (13)	
Former smoker	3 (15)	17 (18)	
Pulmonary disease	1 (5)	5 (5)	1
History of surgery	13 (65)	59 (60)	0.8
CRP, g/dL	0.30 (0.11-0.53)	0.13 (0.03-0.58)	0.26
Disease location			
L1, ileal		34 (35)	
L2, colonic		14 (14)	
L3, ileocolonic		50 (51)	
Disease duration, yr	33 (19-42)	20 (13-29)	0.0055

¹Mann-Whitney *U*-test or Fisher's exact test. Data are the number (%) or median (interquartile range). Significant *P* values are indicated in bold. CEAS: Chronic enteropathy associated with *SLCO2A1* gene; CD: Crohn disease; CRP: C reactive protein.

endoscopic findings, and these patients were categorized according to the Montreal classification^[24]. Disease duration was defined as the period from the onset of symptoms until study enrolment.

Genetic analysis

We used genomic DNA samples extracted from peripheral blood in our previous studies^[10,16,17,25]. In advance, we checked and confirmed that none of the patients with CD had homozygous or compound heterozygous *SLCO2A1* mutations with regards to the major six sites in our previous report^[10], including c.421G>T, c.664G>A, c.940+1G>A, c.1372G>T, c.1461+1G>C, and c.1807C>T. The diagnosis of CEAS was confirmed with 90.2% sensitivity by genotyping these 6 mutations on the basis of our previous investigation.

Measurement of prostaglandin E major urinary metabolite concentrations

All PGE-MUM concentrations were measured at SRL Inc. (Tokyo, Japan) by using a bicyclic PGE-MUM radioimmunoassay kit (Institute of Isotopes Co., Ltd. Budapest, Hungary). PGE-MUM, 7 α -hydroxy-5, 11-diketotetranorprosta-1, 16-dioic acid derived from PGE1 and PGE2 can be transformed to bicyclic PGE-MUM by alkali treatment. Each urine sample was collected and submitted individually according to the manufacturer's protocol described by Fujiwara *et al*^[22]. The urine samples were frozen and stored at -20 °C until the assay. PGE-MUM concentrations were corrected by the urinary creatinine concentration.

Statistical analyses

Categorical variables were expressed as the number (%) and continuous variables were expressed as the median (interquartile range). Comparison of the variables between CEAS and CD were performed using the chi-squared test, Fisher's exact probability test, or the Mann-Whitney *U*-test, where appropriate. The associations between clinical factors and the PGE-MUM concentration were estimated using univariable linear regression analyses. Because of the skewed distribution of the PGE-MUM concentration, values were log-transformed and used for regression analysis.

For the analysis of the clinical significance of PGE-MUM concentration to differentiate CEAS from CD, the logistic regression model was used to estimate the odds ratios (ORs) with 95% confidence intervals (CIs). The diagnostic accuracy of PGE-MUM concentration was evaluated using the area under the curve (AUC) with the receiver operating characteristic (ROC). The optimal cut-off value of PGE-MUM concentration for the diagnosis of CEAS was determined based on the maximum Youden Index (sensitivity + specificity - 1). In addition, sensitivity and specificity were compared using several cut-off values to confirm the diagnostic accuracy of this

cut-off value. Statistical analyses were performed using JMP version 13.0 (SAS Institute, Cary, NC, United States). A *P* value of 0.05 or smaller was considered statistically significant.

RESULTS

Baseline patient characteristics

The enrolled subjects consisted of 20 patients with CEAS (CEAS group) and 98 patients with CD (CD group). The clinical characteristics of the patients are shown in [Table 1](#). In the CEAS group, female patients were more frequent compared with the CD group. The disease duration was significantly longer in the CEAS group than in the CD group (median 33 *vs* 20 years, *P* = 0.0055). There was no significant difference in smoking status or medical history of pulmonary disease. Because most patients with CD were in clinical remission at the time of urine sample collection, the CRP levels remained low in the CD group.

PGE-MUM concentrations

Overall, 118 urinary samples were obtained from CEAS and CD patients. The PGE-MUM concentration ranged from 7.7 to 402.0 (median 34.1) $\mu\text{g/g} \times \text{Cre}$. [Table 2](#) shows the result of univariable linear regression analysis. A positive history of pulmonary disease and history of surgery were significantly associated with a higher PGE-MUM concentration (*P* < 0.0001 and *P* = 0.029, respectively). Serum CRP value was marginally associated with PGE-MUM concentration (*P* = 0.078). Factors with a *P* value < 0.1 were included in the multivariate analysis considering the possibility of confounding factors.

The PGE-MUM concentration was significantly higher in the CEAS group than in the CD group [median interquartile range (IQR): 102.7 (62.0-155.0) $\mu\text{g/g} \times \text{Cre}$ *vs* 27.9 (19.6-40.0) $\mu\text{g/g} \times \text{Cre}$, *P* < 0.0001, [Figure 1](#)]. Crude OR of 1 log unit increase in the PGE-MUM concentration for the distinction between CEAS and CD was 7.3 (95%CI 3.2-16.7, [Table 3](#)). This prediction model showed good differentiating ability between CEAS and CD, with an AUC of 0.90 (95%CI 0.82-0.95). This association was significant even after adjusting for possible confounding factors including age, gender, medical history of pulmonary disease, history of surgery, disease duration, and serum CRP value (model 3). The multivariate-adjusted OR was 29.6 (95%CI 4.7-185.7), with an AUC of 0.95 (95%CI 0.86-0.98).

Correlation between PGE-MUM and CD phenotype

The PGE-MUM concentration in the colonic CD group (L2) was significantly lower than in the other groups (median: 16.5 $\mu\text{g/g} \times \text{Cre}$ *vs* 29.5 $\mu\text{g/g} \times \text{Cre}$, *P* = 0.036), and the PGE-MUM concentration of the ileal CD group (L1) was significantly higher than in the colonic CD group (median: 32.9 $\mu\text{g/g} \times \text{Cre}$ *vs* 16.5 $\mu\text{g/g} \times \text{Cre}$, *P* = 0.017). When we confined the subjects to those with ileal CD, the PGE-MUM concentration in the CEAS group was still significantly higher than that in the ileal CD group (median 102.7 $\mu\text{g/g} \times \text{Cre}$ *vs* 32.9 $\mu\text{g/g} \times \text{Cre}$, *P* < 0.0001).

PGE-MUM cut-off level

[Figure 2](#) shows the ROC curve of the PGE-MUM concentration to differentiate CEAS from CD. Based on the ROC analysis, the optimal cut-off value was identified as 48.9 $\mu\text{g/g} \times \text{Cre}$, which differentiated CEAS from CD with 95.0% sensitivity and 79.6% specificity ([Supplementary Table 2](#)). The AUC was 0.90 (95%CI, 0.82-0.95).

Genetic screening in CD patients with a high PGE-MUM concentration

Among the CD group, six patients had particularly high PGE-MUM concentrations (> 100 $\mu\text{g/g} \times \text{Cre}$). Although we confirmed that they did not harbor the six major mutations in the *SLCO2A1* gene, we screened all 14 coding exons and intron-exon boundaries of the *SLCO2A1* gene using Sanger sequencing to exclude a possible misdiagnosis of CEAS. No homozygous or compound heterozygous *SLCO2A1* mutations were found in the six CD patients.

DISCUSSION

CEAS was initially reported as a novel disease entity referred to as "C" in the Japanese literature in the 1960s^[26]. We recently reported it is an autosomal recessive inherited disease^[8] caused by loss-of-function mutations in the *SLCO2A1* gene^[10]. Small bowel lesions in CEAS are characterized by multiple shallow ulcers with a circular or

Table 2 Factors associated with prostaglandin E major urinary metabolite concentrations

	Univariable linear regression (<i>P</i> value)
Age	<i>P</i> = 0.47 (0.067)
Gender (1, female; 0, male)	<i>P</i> = 0.26 (0.10)
Smoking (1, Current or former smoker; 0, Never smoker)	<i>P</i> = 0.17 (-0.13)
Pulmonary disease	<i>P</i> < 0.0001 (0.36)
History of surgery	<i>P</i> = 0.029 (0.20)
CRP	<i>P</i> = 0.078 (0.16)
Disease duration	<i>P</i> = 0.0040 (0.26)

The values of prostaglandin E major urinary metabolite were log-transformed. Significant *P* values are indicated in bold. CRP: C reactive protein.

eccentric oblique configuration^[18]. In contrast, CD typically develops longitudinal ulcers and cobblestoning, and ulcerations are predominantly located on the mesenteric side. However, both CEAS and CD involve the ileum with occasional ileal stenoses, and the clinical features of CEAS and CD are similar; therefore, it is not easy to differentiate CEAS from CD based on clinical manifestations alone^[11]. Although immunohistochemical staining for SLCO2A1 protein in biopsy specimens from the GI tract was reported to be useful to distinguish CEAS from CD^[27,28], a more convenient and non-invasive screening test is required.

PGE-MUM, a major urinary metabolite derived from PGE2, is clinically used to assess the severity of inflammatory conditions such as interstitial pneumonia^[23], cystic fibrosis^[29], and ulcerative colitis^[30,31]. Similarly, urinary PGE2 metabolite concentrations in patients with CD are higher than those in patients with colorectal neoplasms and healthy controls^[32]. Indeed, PGE-MUM concentrations in patients with CD (median 27.9, IQR 19.6-40.0 µg/g × Cre) in the present study were higher than those in Japanese healthy controls (median 13.1, IQR 10.3-17.2 µg/g × Cre) as reported in previous studies^[15,23]. Nevertheless, the PGE-MUM concentration, a surrogate marker of physiological PGE2 levels^[30], in patients with CEAS was significantly higher than in patients with CD. This difference was still significant even after the adjustment for confounding factors. In addition, we demonstrated that the provisional cut-off value (48.9 µg/g × Cre) of PGE-MUM had a differentiating ability with 95.0% sensitivity and 79.6% specificity (AUC 0.90). Therefore, PGE-MUM is useful to distinguish CEAS from CD.

Although prostaglandins are mediators of active inflammation, many experimental studies have indicated that they play pivotal roles in maintaining mucosal homeostasis^[33] and mucosal repair^[34] of the intestine. PGE2 is mainly derived from arachidonic acid by cyclooxygenase and PGE synthase and PGE2 acts through four different receptor subtypes (EP1, EP2, EP3 and EP4), which partly determines the various physiological effects of PGE2 including the regulation of acid secretion, bicarbonate secretion, mucus production, and mucosal blood flow^[35]. In particular, dimethyl PGE2 (dmPGE2), a stable derivative of PGE2, was reported to exert a potent inhibitory effect against indomethacin-induced intestinal injuries in a rat model *via* activation of EP3 and EP4 receptors^[36]. It was also shown that the deleterious effect of indomethacin on the healing of intestinal ulcers was induced by an EP4 antagonist and suppressed by the co-administration of PGE2 as well as an EP4 agonist^[34]. These results indicate that the protective effect of PGE2 on mucosal injury of the intestine is caused by EP3 and EP4 activation.

PHO, which develops skin and bone disorders caused by impaired prostaglandin metabolism, is classified into two types: PHOAR2 caused by the *SLCO2A1* gene^[12], and PHOAR1 caused by the *HPGD* gene^[37]. Prostaglandin-signaling is terminated by the cellular uptake of prostaglandins *via* the prostaglandin transporter, followed by intracellular oxidation by 15-ketoprostaglandin dehydrogenase (15-PGDH) encoded by the *HPGD* gene^[38]. Thus, cellular uptake of prostaglandins is the first critical step in inactivating prostaglandins. The *SLCO2A1* gene mediates the degradation of prostaglandins *via* their intracellular uptake^[39,40]. The *HPGD* gene is also related to the degradation of prostaglandins by encoding intracellular oxidation enzymes of 15-PGDH. In both conditions, significantly elevated plasma PGE2 levels have been observed^[13,37]. Because PGE2 promotes the proliferation of osteoblasts in bone tissue *via* the expression of vascular endothelial growth factor^[41,42], various clinical manifestations of PHO such as digital clubbing, periostosis, acroosteolysis, painful joint enlargement, and thickened skin, are suggested to be the consequence of an




Figure 1 Comparison of Prostaglandin E-major urinary metabolites concentrations between chronic enteropathy associated with *SLCO2A1* gene ($n = 20$) and Crohn's disease ($n = 98$) groups. Prostaglandin E-major urinary metabolite concentrations in the chronic enteropathy associated with *SLCO2A1* gene group were significantly higher than those in the Crohn's disease group [median (interquartile range): 102.7 (62.0-155.0) $\mu\text{g/g} \times \text{Cre}$ vs 27.9 (19.6-40.0) $\mu\text{g/g} \times \text{Cre}$, $P < 0.0001$]. Mann-Whitney *U*-test. CEAS: Chronic enteropathy associated with *SLCO2A1* gene; CD: Crohn's disease; PEG-MUM: Prostaglandin E-major urinary metabolite.

overabundance of PGE2. Although PGE2 is protective against gastrointestinal mucosal injuries^[35,42], multiple ileal ulcers occur in patients with CEAS, in whom systemic levels of PGE2 are elevated. Considering our previous result that the prostaglandin transporter was expressed on the cellular membrane of vascular endothelial cells^[10], it can be presumed that impaired prostaglandin use in the intestinal mucosa caused by mutation of the *SLCO2A1* gene is a major factor in the pathogenesis of intestinal ulcers in CEAS.

Cryptogenic multifocal ulcerous stenosing enteritis (CMUSE) is another disease entity that causes gastrointestinal ulcerations resembling CEAS^[43,44]. Recently, recessive mutations in the *PLA2G4A* gene, encoding cytoplasmic phospholipase A2- α (cPLA2 α), which hydrolyses cellular membrane phospholipids into arachidonic acids, have been identified as a cause of CMUSE^[44,45]. Because the loss-of-function of cPLA2 α decreases systemic levels of PGE2 (approximated by PGE-MUM) and thromboxane A2^[45], multiple small bowel ulcers, as well as the dysfunction of platelet aggregation are observed in patients with CMUSE^[46]. Although precise analyses of the morphologic features of gastrointestinal lesions in CMUSE have not been reported yet, it seems likely that CEAS and CMUSE share morphologic features regarding the gastrointestinal lesions. In this sense, the measurement of PGE-MUM concentrations can be assumed to be useful to differentiate CEAS from CMUSE, although genetic testing is mandatory to confirm the diagnosis.

This study had several limitations. First, the sample size of the patients with CEAS was relatively small because CEAS is a rare disease. Therefore, we could not validate the cut-off value for the distinction between the two diseases. Second, because most of our CD patients were in clinical remission, the cut-off value may not be appropriate for the distinction of CEAS from active CD. Indeed, some patients with quiescent CD had markedly high PGE-MUM concentrations. Third, we could not completely rule out the possibility that the patients with CD had recessive mutations in the *SLCO2A1* gene. Although we checked the major six mutations of the *SLCO2A1* gene in patients with CD, 9.8% of the mutations might be overlooked based on our previous result^[10].

In conclusion, the measurement of PGE-MUM might be a convenient, non-invasive and useful test to differentiate CEAS from CD, although genetic analysis is mandatory for the confirmation of CEAS. Further studies are necessary to clarify the pathogenesis of CEAS to explore therapeutic targets of this disease.

Table 3 Convert odds ratio for prostaglandin E major urinary metabolite

	Model 1	Model 2	Model 3
Odds ratio of 1 log unit change	7.3	6	29.6
(95%CI)	(3.2-16.7)	(2.6-13.9)	(4.7-185.7)
AUC (95%CI)	0.90 (0.82-0.94)	0.93 (0.87-0.96)	0.95 (0.86-0.98)

CI: Confidence interval; AUC: Area under the curve; Model 1: Prostaglandin E-major urinary metabolite; Model 2: Prostaglandin E-major urinary metabolite + Age + Gender; Model 3: Prostaglandin E-major urinary metabolite + Age + Gender + Medical history of pulmonary disease + History of surgery + Disease duration+ C reactive protein.




Figure 2 Receiver operating characteristic curve of Prostaglandin E-major urinary metabolites concentrations for differentiation between chronic enteropathy associated with *SLCO2A1* gene and Crohn's disease. The area under the receiver operating characteristic curve was 0.90 (95% confidence interval, 0.82-0.95). The optimal cut-off value was 48.9 $\mu\text{g/g} \times \text{Cre}$.

ARTICLE HIGHLIGHTS

Research background

Chronic enteropathy associated with *SLCO2A1* gene (CEAS) is a rare small intestinal disease, but distinct clinical condition from Crohn's disease (CD). To date, we can make a correct diagnosis of CEAS by combination of clinical features and genetic analysis. However, CEAS is sometimes misdiagnosed as CD, because these diseases have some clinical features in common.

Research motivation

A more convenient and non-invasive screening test will help to diagnose CEAS correctly.

Research objectives

To investigate the usefulness of prostaglandin E-major urinary metabolites (PGE-MUM) to distinguish CEAS from CD.

Research methods

Participants were diagnosed CEAS and CD by clinical diagnostic criteria and genetic analysis. All participants' PGE-MUM concentrations were measured by radioimmunoassay. We analyzed differentiating ability of PGE-MUM measurement between CEAS and CD.

Research results

Twenty patients with CEAS and 98 patients with CD were enrolled. It was found that the PGE-MUM concentrations of patients with CEAS were significantly higher than those of patients with CD and this correlation was still statistically significant after adjusting the possible confounding factors. Additionally, the present study showed the provisional cut-off value (48.9 $\mu\text{g/g} \times \text{Cre}$) of PGE-MUM had a differentiating ability with 95.0% sensitivity and 79.6% specificity (area under the curve 0.90).

Research conclusions

The measurement of PGE-MUM can serve as a useful biomarker to differentiate CEAS from CD, although genetic analysis is mandatory for the diagnosis of CEAS.

Research perspectives

Our study showed the usefulness of PGE-MUM to differentiate CEAS from CD. In the present research, the sample size of the patients with CEAS were relatively small, the cut-off value for

the distinction of CEAS from CD should be validated with large sample size of patients with CEAS in the future research.

ACKNOWLEDGEMENTS

We appreciate the patients for participating in this study. We also appreciate Ms. Risa Tsuneyoshi for technical assistance and Drs. Tomohiko Moriyama, Shin Fujioka, Naoya Kubokura, Yoichiro Nuki, Ema Washio, and Shinichi Kawano for their assistance with clinical characterization.

REFERENCES

- Iddan G**, Meron G, Glukhovskiy A, Swain P. Wireless capsule endoscopy. *Nature* 2000; **405**: 417 [PMID: 10839527 DOI: 10.1038/35013140]
- Yamamoto H**, Sekine Y, Sato Y, Higashizawa T, Miyata T, Iino S, Ido K, Sugano K. Total enteroscopy with a nonsurgical steerable double-balloon method. *Gastrointest Endosc* 2001; **53**: 216-220 [PMID: 11174299 DOI: 10.1067/mge.2001.112181]
- Matsumoto T**, Moriyama T, Esaki M, Nakamura S, Iida M. Performance of antegrade double-balloon enteroscopy: Comparison with push enteroscopy. *Gastrointest Endosc* 2005; **62**: 392-398 [PMID: 16111958 DOI: 10.1016/j.gie.2005.04.052]
- Matsumoto T**, Kudo T, Esaki M, Yano T, Yamamoto H, Sakamoto C, Goto H, Nakase H, Tanaka S, Matsui T, Sugano K, Iida M. Prevalence of non-steroidal anti-inflammatory drug-induced enteropathy determined by double-balloon endoscopy: A Japanese multicenter study. *Scand J Gastroenterol* 2008; **43**: 490-496 [PMID: 18365915 DOI: 10.1080/00365520701794121]
- Esaki M**, Matsumoto T, Watanabe K, Arakawa T, Naito Y, Matsuura M, Nakase H, Hibi T, Matsumoto T, Nouda S, Higuchi K, Ohmiya N, Goto H, Kurokawa S, Motoya S, Watanabe M. Use of capsule endoscopy in patients with Crohn's disease in Japan: A multicenter survey. *J Gastroenterol Hepatol* 2014; **29**: 96-101 [PMID: 24354993 DOI: 10.1111/jgh.12411]
- Matsumoto T**, Iida M, Matsui T, Yao T, Watanabe H, Yao T, Okabe H. Non-specific multiple ulcers of the small intestine unrelated to non-steroidal anti-inflammatory drugs. *J Clin Pathol* 2004; **57**: 1145-1150 [PMID: 15509673 DOI: 10.1136/jcp.2003.015735]
- Matsumoto T**, Iida M, Matsui T, Yao T. Chronic nonspecific multiple ulcers of the small intestine: A proposal of the entity from Japanese gastroenterologists to Western enteroscopists. *Gastrointest Endosc* 2007; **66**: S99-107 [PMID: 17709045 DOI: 10.1016/j.gie.2007.01.004]
- Matsumoto T**, Kubokura N, Matsui T, Iida M, Yao T. Chronic nonspecific multiple ulcer of the small intestine segregates in offspring from consanguinity. *J Crohns Colitis* 2011; **5**: 559-565 [PMID: 22115375 DOI: 10.1016/j.crohns.2011.05.008]
- Esaki M**, Umeno J, Kitazono T, Matsumoto T. Clinicopathologic features of chronic nonspecific multiple ulcers of the small intestine. *Clin J Gastroenterol* 2015; **8**: 57-62 [PMID: 25788296 DOI: 10.1007/s12328-015-0559-x]
- Umeno J**, Hisamatsu T, Esaki M, Hirano A, Kubokura N, Asano K, Kochi S, Yanai S, Fuyuno Y, Shimamura K, Hosoe N, Ogata H, Watanabe T, Aoyagi K, Ooi H, Watanabe K, Yasukawa S, Hirai F, Matsui T, Iida M, Yao T, Hibi T, Kosaki K, Kanai T, Kitazono T, Matsumoto T. A Hereditary Enteropathy Caused by Mutations in the SLCO2A1 Gene, Encoding a Prostaglandin Transporter. *PLoS Genet* 2015; **11**: e1005581 [PMID: 26539716 DOI: 10.1371/journal.pgen.1005581]
- Umeno J**, Esaki M, Hirano A, Fuyuno Y, Ohmiya N, Yasukawa S, Hirai F, Kochi S, Kurahara K, Yanai S, Uchida K, Hosomi S, Watanabe K, Hosoe N, Ogata H, Hisamatsu T, Nagayama M, Yamamoto H, Abukawa D, Kakuta F, Onodera K, Matsui T, Hibi T, Yao T, Kitazono T, Matsumoto T; CEAS study group. Clinical features of chronic enteropathy associated with SLCO2A1 gene: A new entity clinically distinct from Crohn's disease. *J Gastroenterol* 2018; **53**: 907-915 [PMID: 29313109 DOI: 10.1007/s00535-017-1426-y]
- Seifert W**, Kühnisch J, Tüystüz B, Specker C, Brouwers A, Horn D. Mutations in the prostaglandin transporter encoding gene SLCO2A1 cause primary hypertrophic osteoarthropathy and isolated digital clubbing. *Hum Mutat* 2012; **33**: 660-664 [PMID: 22331663 DOI: 10.1002/humu.22042]
- Zhang Z**, Xia W, He J, Zhang Z, Ke Y, Yue H, Wang C, Zhang H, Gu J, Hu W, Fu W, Hu Y, Li M, Liu Y. Exome sequencing identifies SLCO2A1 mutations as a cause of primary hypertrophic osteoarthropathy. *Am J Hum Genet* 2012; **90**: 125-132 [PMID: 22197487 DOI: 10.1016/j.ajhg.2011.11.019]
- Zhang Z**, He JW, Fu WZ, Zhang CQ, Zhang ZL. Mutations in the SLCO2A1 gene and primary hypertrophic osteoarthropathy: A clinical and biochemical characterization. *J Clin Endocrinol Metab* 2013; **98**: E923-E933 [PMID: 23509104 DOI: 10.1210/jc.2012-3568]
- Okayasu I**, Ohnishi H, Sarandi I, Shojima J, Komatsu J, Oritsu M, Sasabe M, Nanami KO, Matsuura M, Azumi J, Ito S, Fujiwara M. Significant increase of prostaglandin E-major urinary metabolite in male smokers: A screening study of age and gender differences using a simple radioimmunoassay. *J Clin Lab Anal* 2014; **28**: 32-41 [PMID: 24375858 DOI: 10.1002/jcla.21640]
- Hirano A**, Yamazaki K, Umeno J, Ashikawa K, Aoki M, Matsumoto T, Nakamura S, Ninomiya T, Matsui T, Hirai F, Kawaguchi T, Takazoe M, Tanaka H, Motoya S, Kiyohara Y, Kitazono T, Nakamura Y, Kamatani N, Kubo M. Association study of 71 European Crohn's disease susceptibility loci in a Japanese population. *Inflamm Bowel Dis* 2013; **19**: 526-533 [PMID: 23388546 DOI: 10.1097/MIB.0b013e31828075e7]
- Yamazaki K**, Umeno J, Takahashi A, Hirano A, Johnson TA, Kumasaka N, Morizono T, Hosono N, Kawaguchi T, Takazoe M, Yamada T, Suzuki Y, Tanaka H, Motoya S, Hosokawa M, Arimura Y, Shinomura Y, Matsui T, Matsumoto T, Iida M, Tsunoda T, Nakamura Y, Kamatani N, Kubo M. A genome-wide association study identifies 2 susceptibility Loci for Crohn's disease in a Japanese population. *Gastroenterology* 2013; **144**: 781-788 [PMID: 23266558 DOI: 10.1053/j.gastro.2012.12.021]
- Matsumoto T**, Nakamura S, Esaki M, Yada S, Koga H, Yao T, Iida M. Endoscopic features of chronic nonspecific multiple ulcers of the small intestine: Comparison with nonsteroidal anti-inflammatory drug-

- induced enteropathy. *Dig Dis Sci* 2006; **51**: 1357-1363 [PMID: 16868823 DOI: 10.1007/s10620-006-9080-x]
- 19 **Hosoe N**, Ohmiya N, Hirai F, Umeno J, Esaki M, Yamagami H, Onodera K, Bamba S, Imaeda H, Yanai S, Hisamatsu T, Ogata H, Matsumoto T; CEAS Atlas Group. Chronic Enteropathy Associated With SLCO2A1 Gene [CEAS]-Characterisation of an Enteric Disorder to be Considered in the Differential Diagnosis of Crohn's Disease. *J Crohns Colitis* 2017; **11**: 1277-1281 [PMID: 28510689 DOI: 10.1093/ecco-jcc/jjx068]
- 20 **Hisabe T**, Hirai F, Matsui T, Watanabe M. Evaluation of diagnostic criteria for Crohn's disease in Japan. *J Gastroenterol* 2014; **49**: 93-99 [PMID: 23546557 DOI: 10.1007/s00535-013-0798-x]
- 21 **Ware JS**, Wain LV, Channavajhala SK, Jackson VE, Edwards E, Lu R, Siew K, Jia W, Shrine N, Kinnear S, Jalland M, Henry AP, Clayton J, O'Shaughnessy KM, Tobin MD, Schuster VL, Cook S, Hall IP, Glover M. Phenotypic and pharmacogenetic evaluation of patients with thiazide-induced hyponatremia. *J Clin Invest* 2017; **127**: 3367-3374 [PMID: 28783044 DOI: 10.1172/JCI89812]
- 22 **Fujiwara M**, Okayasu I, Oritsu M, Komatsu J, Yoshitsugu M, Katoh Y, Bandoh T, Toyoshima H, Kase Y, Sugihara K, Kanno J, Hayashi Y. Significant increase in prostaglandin E-main urinary metabolite by laxative administration: Comparison with ulcerative colitis. *Digestion* 2000; **61**: 201-206 [PMID: 10773726 DOI: 10.1159/000007758]
- 23 **Horikiri T**, Hara H, Saito N, Araya J, Takasaka N, Utsumi H, Yanagisawa H, Hashimoto M, Yoshii Y, Wakui H, Minagawa S, Ishikawa T, Shimizu K, Numata T, Arihiro S, Kaneko Y, Nakayama K, Matsuura T, Matsuura M, Fujiwara M, Okayasu I, Ito S, Kuwano K. Increased levels of prostaglandin E-major urinary metabolite (PGE-MUM) in chronic fibrosing interstitial pneumonia. *Respir Med* 2017; **122**: 43-50 [PMID: 27993290 DOI: 10.1016/j.rmed.2016.11.017]
- 24 **Satsangi J**, Silverberg MS, Vermeire S, Colombel JF. The Montreal classification of inflammatory bowel disease: Controversies, consensus, and implications. *Gut* 2006; **55**: 749-753 [PMID: 16698746 DOI: 10.1136/gut.2005.082909]
- 25 **Fuyuno Y**, Yamazaki K, Takahashi A, Esaki M, Kawaguchi T, Takazoe M, Matsumoto T, Matsui T, Tanaka H, Motoya S, Suzuki Y, Kiyohara Y, Kitazono T, Kubo M. Genetic characteristics of inflammatory bowel disease in a Japanese population. *J Gastroenterol* 2016; **51**: 672-681 [PMID: 26511940 DOI: 10.1007/s00535-015-1135-3]
- 26 **Okabe H**, Sakimura M. "Hi Tokui-sei Tahatsu-sei Shouchou Kaiyou-shou". *Stomach Intest* 1968; **3**: 1539-1549
- 27 **Yamaguchi S**, Yanai S, Nakamura S, Kawasaki K, Eizuka M, Uesugi N, Sugai T, Umeno J, Esaki M, Matsumoto T. Immunohistochemical differentiation between chronic enteropathy associated with SLCO2A1 gene and other inflammatory bowel diseases. *Intest Res* 2018; **16**: 393-399
- 28 **Yanai S**, Yamaguchi S, Nakamura S, Kawasaki K, Toya Y, Yamada N, Eizuka M, Uesugi N, Umeno J, Esaki M, Okimoto E, Ishihara S, Sugai T, Matsumoto T. Distinction of Chronic Enteropathy associated with SLCO2A1 Gene from Crohn's Disease. *Gut Liver* 2018; **13**: 62-66
- 29 **Jabr S**, Gartner S, Milne GL, Roca-Ferrer J, Casas J, Moreno A, Gelpi E, Picado C. Quantification of major urinary metabolites of PGE2 and PGD2 in cystic fibrosis: Correlation with disease severity. *Prostaglandins Leukot Essent Fatty Acids* 2013; **89**: 121-126 [PMID: 23791427 DOI: 10.1016/j.plefa.2013.06.001]
- 30 **Arai Y**, Arihiro S, Matsuura T, Kato T, Matsuoka M, Saruta M, Mitsunaga M, Matsuura M, Fujiwara M, Okayasu I, Ito S, Tajiri H. Prostaglandin E-major urinary metabolite as a reliable surrogate marker for mucosal inflammation in ulcerative colitis. *Inflamm Bowel Dis* 2014; **20**: 1208-1216 [PMID: 24846719 DOI: 10.1097/MIB.0000000000000062]
- 31 **Arai Y**, Matsuura T, Matsuura M, Fujiwara M, Okayasu I, Ito S, Arihiro S. Prostaglandin E-Major Urinary Metabolite as a Biomarker for Inflammation in Ulcerative Colitis: Prostaglandins Revisited. *Digestion* 2016; **93**: 32-39 [PMID: 26788915 DOI: 10.1159/000441665]
- 32 **Johnson JC**, Schmidt CR, Shrubsole MJ, Billheimer DD, Joshi PR, Morrow JD, Heslin MJ, Washington MK, Ness RM, Zheng W, Schwartz DA, Coffey RJ, Beauchamp RD, Merchant NB. Urine PGE-M: A metabolite of prostaglandin E2 as a potential biomarker of advanced colorectal neoplasia. *Clin Gastroenterol Hepatol* 2006; **4**: 1358-1365 [PMID: 16996805 DOI: 10.1016/j.cgh.2006.07.015]
- 33 **Otani T**, Yamaguchi K, Scherl E, Du B, Tai HH, Greifer M, Petrovic L, Daikoku T, Dey SK, Subbaramaiah K, Dannenberg AJ. Levels of NAD(+)-dependent 15-hydroxyprostaglandin dehydrogenase are reduced in inflammatory bowel disease: Evidence for involvement of TNF-alpha. *Am J Physiol Gastrointest Liver Physiol* 2006; **290**: G361-G368 [PMID: 16195422 DOI: 10.1152/ajpgi.00348.2005]
- 34 **Hatazawa R**, Ohno R, Tanigami M, Tanaka A, Takeuchi K. Roles of endogenous prostaglandins and cyclooxygenase isozymes in healing of indomethacin-induced small intestinal lesions in rats. *J Pharmacol Exp Ther* 2006; **318**: 691-699 [PMID: 16699067 DOI: 10.1124/jpet.106.103994]
- 35 **Takeuchi K**, Kato S, Amagase K. Prostaglandin EP receptors involved in modulating gastrointestinal mucosal integrity. *J Pharmacol Sci* 2010; **114**: 248-261 [PMID: 21041985 DOI: 10.1254/jphs.10R06CR]
- 36 **Kunikata T**, Tanaka A, Miyazawa T, Kato S, Takeuchi K. 16,16-Dimethyl prostaglandin E2 inhibits indomethacin-induced small intestinal lesions through EP3 and EP4 receptors. *Dig Dis Sci* 2002; **47**: 894-904 [PMID: 11991626 DOI: 10.1023/A:1014725024519]
- 37 **Uppal S**, Diggle CP, Carr IM, Fishwick CW, Ahmed M, Ibrahim GH, Helliwell PS, Latos-Bieleńska A, Phillips SE, Markham AF, Bennett CP, Bonthron DT. Mutations in 15-hydroxyprostaglandin dehydrogenase cause primary hypertrophic osteoarthropathy. *Nat Genet* 2008; **40**: 789-793 [PMID: 18500342 DOI: 10.1038/ng.153]
- 38 **Nakanishi T**, Tamai I. Roles of Organic Anion Transporting Polypeptide 2A1 (OATP2A1/SLCO2A1) in Regulating the Pathophysiological Actions of Prostaglandins. *AAPS J* 2017; **20**: 13 [PMID: 29204966 DOI: 10.1208/s12248-017-0163-8]
- 39 **Kanai N**, Lu R, Satriano JA, Bao Y, Wolkoff AW, Schuster VL. Identification and characterization of a prostaglandin transporter. *Science* 1995; **268**: 866-869 [PMID: 7754369 DOI: 10.1126/science.7754369]
- 40 **Nomura T**, Lu R, Pucci ML, Schuster VL. The two-step model of prostaglandin signal termination: *In vitro* reconstitution with the prostaglandin transporter and prostaglandin 15 dehydrogenase. *Mol Pharmacol* 2004; **65**: 973-978 [PMID: 15044627 DOI: 10.1124/mol.65.4.973]
- 41 **Harada S**, Nagy JA, Sullivan KA, Thomas KA, Endo N, Rodan GA, Rodan SB. Induction of vascular endothelial growth factor expression by prostaglandin E2 and E1 in osteoblasts. *J Clin Invest* 1994; **93**: 2490-2496 [PMID: 8200985 DOI: 10.1172/JCI117258]
- 42 **Rao R**, Redha R, Macias-Perez I, Su Y, Hao C, Zent R, Breyer MD, Pozzi A. Prostaglandin E2-EP4 receptor promotes endothelial cell migration via ERK activation and angiogenesis in vivo. *J Biol Chem*

- 2007; **282**: 16959-16968 [PMID: [17401137](#) DOI: [10.1074/jbc.M701214200](#)]
- 43 **Shoesmith JH**, Tate GT, Wright CJ. Multiple Strictures Of The Jejunum. *Gut* 1964; **5**: 132-135 [PMID: [14159400](#) DOI: [10.1136/gut.5.2.132](#)]
- 44 **Perlemuter G**, Guillevin L, Legman P, Weiss L, Couturier D, Chaussade S. Cryptogenetic multifocal ulcerous stenosing enteritis: An atypical type of vasculitis or a disease mimicking vasculitis. *Gut* 2001; **48**: 333-338 [PMID: [11171822](#) DOI: [10.1136/gut.48.3.333](#)]
- 45 **Adler DH**, Cogan JD, Phillips JA, Schnetz-Boutaud N, Milne GL, Iverson T, Stein JA, Brenner DA, Morrow JD, Boutaud O, Oates JA. Inherited human cPLA(2alpha) deficiency is associated with impaired eicosanoid biosynthesis, small intestinal ulceration, and platelet dysfunction. *J Clin Invest* 2008; **118**: 2121-2131 [PMID: [18451993](#) DOI: [10.1172/JCI30473](#)]
- 46 **Brooke MA**, Longhurst HJ, Plagnol V, Kirkby NS, Mitchell JA, Rüschemdorf F, Warner TD, Kelsell DP, MacDonald TT. Cryptogenic multifocal ulcerating stenosing enteritis associated with homozygous deletion mutations in cytosolic phospholipase A2- α . *Gut* 2014; **63**: 96-104 [PMID: [23268370](#) DOI: [10.1136/gutjnl-2012-303581](#)]



Published By Baishideng Publishing Group Inc
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-2238242
Fax: +1-925-2238243
E-mail: bpgoffice@wjgnet.com
Help Desk: <http://www.f6publishing.com/helpdesk>
<http://www.wjgnet.com>

