Germ cell development in male *Perinereis nuntia* and gamete spawning mechanisms in males and females

マリア, ジャヌアリ, ピーター

<http://hdl.handle.net/2324/4110548>

出版情報 : 九州大学, 2020, 博士 (農学), 課程博士
バージョン :
権利関係 :

**Germ cell development in male *Perinereis wilsoni* and gamete
spawning mechanisms in males and females**

*Dissertation submitted in accordance with the requirement of Kyushu University for the
Degree of Doctor of Philosophy*

Maria January Peter

Fisheries Research Laboratory

Graduate School of Bioresource and Bioenvironmental Sciences

Faculty of Agriculture

Kyushu University

2020

TABLE OF CONTENTS

LIST OF ABBREVIATIONS

LIST OF FIGURES

LIST OF PUBLICATIONS

	Page No.
CHAPTER 1	
General introduction on polychaetes	1
1.1 Taxonomy and distribution	2
1.2 Morphology, habitat and feeding	2
1.3 Reproduction	3
1.3.1 Gametogenesis	4
1.3.2 Endocrine Control	5
1.3.3 Environmental control	5
1.4 Embryonic development and segmentation	6
1.5 Importance of polychaetes	7
1.6 Experimental animal (<i>Perinereis wilsoni</i>) classification	8
1.7 Figures	9
CHAPTER 2	
Germ cells development in male <i>Perinereis wilsoni</i>	10
2.1 Abstract	11
2.2 Introduction	12
2.3 Materials and Methods	14
2.3.1 Animals	14

2.3.2 cDNA cloning of <i>Perinereis wilsoni</i> - <i>piwi</i> gene	14
2.3.3 <i>In situ</i> hybridization	14
2.3.4 Histology	15
2.4 Results	16
2.4.1 Germ cell localization in male <i>P. wilsoni</i>	16
2.4.2 Sexual development of gametes in <i>P. wilsoni</i>	17
2.4.3 Figures and Table	19
2.5 Discussions	33
2.5.1 Germ cells localization and gametes production in <i>P. wilsoni</i>	33
2.6 Conclusion	36
CHAPTER 3	
Mechanisms of gamete releasing in males and females <i>P. wilsoni</i>	37
3.1 Abstract	38
3.2 Introduction	39
3.3 Materials and Methods	41
3.3.1 Observation of spawning sites	41
3.3.2 Histology	41
3.4 Results	42
3.4.1 Spawning sites in <i>P. wilsoni</i>	42
3.4.2 Gamete spawning sites in male <i>P. wilsoni</i>	42

3.4.3 Gamete spawning sites in female <i>P. wilsoni</i>	42
3.4.4 Figures	44
3.5 Discussions	51
3.5.1 Gamete spawning sites in <i>P. wilsoni</i>	51
3.5.2 The eosinophilic structure associated with gamete during spawning	52
3.6 Conclusion	53
CHAPTER 4	
General discussion	55
4.1 Germ cells localization in male and gamete production <i>P. wilsoni</i>	55
4.2 Gametes releasing sites in males and females <i>P. wilsoni</i>	58
ACKNOWLEDGEMENTS	62
REFERENCES	64

LIST OF ABBRIVIATIONS

NBT	Nitro Blue Tetrazodium
BCIP	5-Bromo-4-Chloro-3-Indoly Phosphate
PBS	Phosphate Buffered Saline
<i>Pl. dumerilii</i>	<i>Platynereis dumerilii</i>
<i>P. wilsoni</i>	<i>Perinereis wilsoni</i>
<i>Pw-vasa</i>	<i>Perinereis wilsoni -vasa</i>
<i>Pw-piwi</i>	<i>Perinereis wilsoni - piwi</i>
SAZ	Segment Addition Zone

LIST OF FIGURES

CHAPTER 1

Figure. 1. Experimental animal, *Perinereis wilsoni*

Figure. 2. Polychaete (*Perinereis wilsoni*) aquaculture in Oita prefecture, Japan

CHAPTER 2

Figure. 1. Whole mount *in situ* hybridization with the *Pw-piwi* probe in an adult male worm in late August.

Figure. 2. Horizontal sections of the hybridized worm showing the localization of the *Pw-piwi* expression in an adult male in late August.

Figure. 3. Specificity of the *Piwi* probe in *in situ* hybridization.

Figure. 4. Germ cell clusters in a juvenile worm.

Figure. 5. Germ cells before and after the start of vitellogenesis.

Figure. 6. Germ cells fill body cavities of each segment in March (mid-body region).

Figure. 7. Monthly development of male germ cells.

Figure. 8. Whole mount *in situ* hybridization with *Pw-piwi* in early March.

Figure. 9. Whole mount *in situ* hybridization of *Pw-piwi* in end of March.

Figure. 10. Whole mount *in situ* hybridization of *Pw-piwi* in April; swimming epitoke.

Figure. 11. *Pn-piwi* expression in monthly development of male germ cells.

Figure. 12. Schematic diagram on the gametogenic cycle of *P. wilsoni*.

Figure. 13. Schematic diagram on the timing of the germ cells migration from the parapodia to the deeper body cavity during segments addition and the start of the vitellogenesis.

CHAPTER 3

Figure. 1. Observation of spawning sites in *P. wilsoni*. (b) Male (c) female

Figure. 2. Horizontal sections of swimming male worm just prior to spawning in April.

Figure. 3. Sperm spawning area in a tail region of a male.

Figure. 4. Histological observation of sperm releasing sites in *P. wilsoni*.

Figure. 5. Horizontal sections of swimming male worm just prior to spawning in April.

Figure. 6. Egg spawning area in female.

Figure. 7. The eosinophilic structure associated with gamete release indicated by arrows.
(A)

Spawning male. (B) Spawning female. H&E stain.

LIST OF PUBLICATIONS

Peer-reviewed scientific paper:

Maria January Peter, Mercedes Maceren-Pates, Gaudioso Pates Jr, Michiyasu Yoshikuni Yoshihisa Kurita (2020). Germ cell development in male *Perinereis wilsoni* and gamete spawning mechanisms in males and females. Zoological Science

Scientific conference abstract presentation

Maria January Peter and Maceren-Pates Mercedes: A study of germ cells development in *Perinereis wilsoni* 90th International conference. Biodiversity, Ecology and Evolution (2019), Amakusa Marine Biology Laboratory, Kyushu University, Japan.

CHAPTER 1

GENERAL INTRODUCTION

1.1 Taxonomy and distribution of Polychaetes

Annelida is the phylum of the segmented worms. It is one of the most diverse phyla in the animal kingdom. The phylum comprises more than 20000 species which widely distributed in ocean water, fresh water and in moist terrestrial environment (Fauchald and Rouse, 1997; Kriska, 2013). Over centuries, annelids had been classified into three classes; Polychaeta (marine worms), Oligochaeta (Earthworms) and Hirudinea (Leeches). Among these, polychaeta is the largest group in the phylum, mostly comprised by marine worms and few (about 2%) are fresh water species (Glasby and Timm, 2008; Shoobs *et al.*, 2016). There are about 11,456 species of polychaetes which are distributed in a wide range of habitats; wetlands, sandflats, intertidal zones and deep seas (Kriska, 2013; Pamungkas *et al.*, 2019). Based on the development of the anterior appendages and life habits, polychaetes had been classified into two major groups, Errantia and Sedentaria. Errantia includes the active scavengers, the free living and crawler worms of the family Nereididae and Phyllodocida. Sedentaria are permanent burrowers or tube dwellers, for example, the Capitella. (Hartman, 1968, 1969; Kriska, 2013).

1.2 Polychaetes morphology, habitat and feeding

A body of polychaete is divided into three parts of head, trunk and tail region. The head part includes a prostomium which bears palps, eyes, tentacles and antenna and a peristomium where the mouth is located. The body length ranges from 1mm to 3m with the average size of 5cm to 10cm (Kriska, 2013). Polychaetes occupy a wide range of habitats; burrowing in sediments or sands, living under rocks or corals and some are tube-dwellers which includes some members of the family Oweniidae (Dauvin and Thiébaud, 1994). The diversity in life style of polychaetes is greatly influenced by morphological variations and feeding habits as well. There are active hunters such as Phyllodocidae, Syllidae and Nereidae, some are

deposit feeders such as Capitellids and Arenicolids while others are filter feeders like Lumbrids, Serpulids and Sabellids (Fauchald, 1977).

1.3 Polychaetes reproduction

Majority of the polychaetes are gonochoric and only few are hermaphrodites. The reproduction mode varies among species depending on their life cycle. In dioecious reproduction, gametes mature in separate sexes. Spawning can be broadcasting with external fertilization as in nereidids (Fauchald, 1977) or viviparity development shown by Syllidae (Franke, 1999). Asexual reproduction involves splitting of the adult worm into two or more individuals or by budding off the new individual from the parent. In most case, budding occurs at the posterior region (reproductive region) and the parent organism will regenerate and survive to reproduce another time. Examples are *Pristina leidy*, syllids and oweniids (Oliver, 1984; Franke, 1999; Özpölat and Bely, 2015).

Nereidids are errant polychaetes with unique reproductive strategy called epitoky. Epitoky is a form of reproduction in which sexually immature worm (atoke) transform into sexually mature worm (epitoke), and epitokes emerge from benthic to short pelagic life in spawning season (Fong, 1993; Bartels *et al.*, 1990; Franke, 1999; Hébert-Chatelain *et al.*, 2008). There are two forms of epitoky, Epigamy and Schizogamy. Epigamy is a sexual reproduction when the whole body transforms into the epitoke. Schizogamy is another form of reproduction in which only part of the body will transform into sexual form and detach from asexual part by fission (Pamungkas and Glasby, 2015). In breeding season, ripe animals release gametes after a swimming dance (nuptial) and die shortly within hours (Fong, 1993; Bartels *et al.*, 1990; Hébert-Chatelain *et al.*, 2008). However, some species had evolved into iteroparous and can survive more than one spawning season in a life time such as *Eunice fucata* (Fischer, 1999). Sexes of mature individuals can easy be differentiated due to differences in gamete

colours; for examples in *P. wilsoni* male has whitish or cream colour and female look greenish, in *P. dumerilii*, male has cream/whitish colour and female has yellow (Rebscher *et al.*, 2007) and in *Nereis fucata* lilac or blue (Brown- Gilpin, 1959).

Transformation into pelagic life is energy consuming process which involves tremendous changes in body system so as to adapt the swimming life. Those are addition sets of setae (bristles), enlarged parapodia, histolysis of the body wall, reorganisation of the body musculature, growth of vascular system and respiratory surface, which are the morphological and physiological changes reported in the previous studies (Fischer, 1999; Hébert-Chatelain *et al.*, 2008).

1.3.1 Gametogenesis

In many polychaetes, the developing germ cells are usually observed floating in the coelomic cavity. The location of the gonads varies among species, and in some species it is not yet identified (Rice and Eckelbarger, 1989; Gambi, 2000). In *P. dumerilii*, the gonad settles at the neck region where germ cells proliferate and move to fill other body segments (Rebscher, 2007). In *Nicole zostericola*, the gonad is located beneath the peritoneum in the thoracic region of the body (Eckelbarger, 1975). In *P. wilsoni*, primordial germ cells settles at the distal end of each parapodium to proliferate before migration to the deeper body cavity (Maceren-Pates *et al.*, 2015).

Vitellogenesis in polychaetes develops intraovarian or extraovarian. In intraovarian development, female germ cells develop and mature inside the ovary and are released after maturation. In extraovarian vitellogenesis, female germ cells are released singly from the ovary to develop in the coelomic cavity, or released as the cluster to develop and then separated as an individual oocyte after development (Eckelbarger, 1983). The start of vitellogenesis can be used as one of the criteria for sex determination due to enlargement of germ cells as the result of vitellogenin accumulation (Fischer, 1974; Gambi, 2000).

1.3.2 Endocrine control

The timing of gametogenic cycle of numerous polychaetes is controlled by and endocrine system in coordination with the environmental cues. The process of the germ cells development itself is said to be solely regulated by the endocrine and neural systems responding to external stimulus (Golding, 1967; Bentley, 1992; Lawrence, 1996) reported that a juvenile hormone secreted by a supraphrægeal ganglion induced accumulation of oogonia in the coelomic cavity but inhibit the sexual maturation. In *Nereis divoscolor*, cerebral endocrine activities were observed to be high in most time of the gametogenic cycle and declined as the life cycle terminates (Dilding, 1983). A similar endocrine trend was observed during oocytes development in *Nephtys hombergii* (Olive and Bentley, 1980). Fischer and Hausenschild (1974) suggested that a brain hormone might be responsible in controlling the extent of proliferation of oocytes and the inhibin regulated the homogenous development of the oocytes in late stages of maturation in *P. dumerilii*. Lawrence (1996) found that gonadotropin promoted egg synthesis in *Harmothoe imbricate*.

1.3.3 Environmental factors

Gametogenic cycle of numerous polychaetes is highly synchronized with the environmental cues such as the start of their breeding season corresponds with the rise in sea temperature. Studies on temperate species show that the start of gamete development has been synchronised with the low temperature and short photoperiod in the winter season and the gamete reaches the maturity size in the next spring (Olive, 1978; Garwood, 1980; Bartels-Hardege and Zeeck, 1990; William, 1997; Clark 1998). Rise in sea temperature and increase in day length have been reported to correspond with nutrients productivity and increase in metabolic activities which in turn favour the growth and development of the newly hatched larvae and growing juveniles (Newell *et al.*, 1982; Bartels-Hardege and Zeeck, 1990; William, 1997; Lawrence and Saome, 2009). Spawning events have been also found

to relate with certain phases of the lunar cycle. For instance, *P. dumerilii* and *Nereis succinea* spawning events happen 18 days after new moon (Rasmussen, 1973).

1.4 Embryonic development and segmentation

In Nereidids, gametes fertilize in the external environment after broadcast spawning. Within 1 hour post fertilization (hpf), the fertilized eggs covered by the jelly-coat for protection settle to the substratum prior to start the cleavage. The embryos undergo spiral cleavage forming a ball of mass cells. The embryos hatch within 7-8 days post fertilization (Hardege and Bartels-Hardege, 1996).

The larvae undergo three stages; trochophore, metatrochophore and nectochaete. Trochophore is the free swimming and non-feeding larvae formed 2-3 days post fertilization. Metatrochophore at 4-6 dpf starts the parapodium protrusion accompanied with the formation of chaetae, but does not actively move. In this stage, the larvae accomplished to form the adult eyes. Nectochaete (7 dpf) has a three fully-segmented body with well-developed parapodia, ciliary bands and chaetae. Nectochaete larva do not feed and have a mixed pelagic and benthic life style (Fischer *et al.*, 2010).

Segments are added from the pygidium, accordingly, anterior segments are older than posterior segments. In errant polychaetes, the segment addition is active in a late larvae and juvenile stage, and slows down as the animal reaches the sexual maturity stage. The rate of segment addition varies among species. However, the growth and enlargement continues as the germ cells develop (Clack and Clack, 1962; Reish, 1993). The process of segment addition was reported to be controlled by growth-promoting hormones secreted by a supraesophageal ganglion (Clack, 1963). Each segment bears one pair of parapodia on each side of the body which can be used as means for locomotion, sensation, respiration and excretion (Giangrande and Gambi, 1998). In some species, parapodium can also act as a

temporal gonad as recent described in *P. wilsoni* (Maceren *et al.*, 2015). Segments are separated each other by the complete intersegmental septum (Maceren *et al.*, 2015) or the incomplete septum (Fischer, 2004).

1.5 Importance of Polychaetes

Polychaetes play the important roles in the ecosystem as food source for shrimps, fishes and shore birds. Some species are commonly used as live baits by anglers in fishing (Lawrence *et al.*, 1998). Currently, polychaetes play great contributions in the aquaculture industry of crustaceans and fishes because polychaetes are rich in nutrients, especially the polysaturated fatty acids. Several reports have identified that the feeding polychaetes enhances the gonad maturation in shrimps (Olive, 1999; Techaprempreecha *et al.*, 2011; Kumar *et al.*, 2017; Bischoff, 2014;). Aside nutrients provision, polychaetes have been reported to be useful in the treatment of the wastewater in the mariculture system and bioremediation (Giangrande *et al.*, 2005; Palmer, 2010; Stabili *et al.*, 2010; Ito *et al.*, 2011). A case study conducted on *Nereis diversicolor* reported that polychaetes enhanced the growth of microorganisms (bacteria) which in return increased decomposition of organic sediments in fish farms (Jadao *et al.*, 2008). Moreover, Lucciano *et al.*, (2007, 2010) described that the ability of filter feeder polychaetes (*Branchiommma luctuosum* and *Sabella spallanzanii*) to accumulate bacteria is useful as bioremediator as well as the bioindicator of the microbe pollution in water. In regard with this importance, several studies have suggested that polychaetes can be useful in the integrated aquaculture, nutrient recycling and also in the removal of wastes (Jadao *et al.*, 2008). Nevertheless, Hutchings (1998) reported that the burrowing and feeding activities of polychaetes enhance sedimentation and nutrient recycling in the ecosystem. Additionally, benthic dweller and scavenger polychaetes such as

Capitella sp. provide suitable specimens for the biomonitoring and marine toxicological studies (Bryan and Gibbs, 1987; Nugteren *et al.*, 2009; Aller R, 2001).

1.6 Experimental animal (*Perinereis wilsoni*) classification and distribution

Kingdom	:	Animalia
Phylum	:	Annelida
Class	:	Polychaeta
Order	:	Phyllodocida
Family	:	Nereididae
Genus	:	<i>Perinereis</i>
Species	:	<i>P. wilsoni</i>

Perinereis wilsoni (*P. wilsoni*) is the nereidid polychaete which is commonly found in the intertidal reef flats or rock shores. It is widely distributed Taiwan, Japan, South China Sea and South Korea (Glasby and Hwey- Lian, 2006).

Fig. 1. Experimental animal, *Perinereis wilsoni*

Fig. 2. Polychaete (*Perinereis wilsoni*) aquaculture in Oita prefecture, Japan

CHAPTER 2

Germ cell development in male *Perinereis wilsoni*

2.1 Abstract

Perinereis wilsoni is a fully segmented worm with complete intersegmental septa. A previous study of females revealed that germ cells of this animal originated in the tail end segment, called pygidium. In females, germ cells were duplicated in the pygidium, transferred to a newly generating segment, and then settled in the parapodia. Within each segment, the settled germ cells proliferated in the parapodia and then migrated into a body cavity area to begin meiotic development. However, there is no information about germ cell development in males. In this study, I analyzed the male germ cell development by *in situ* hybridization with use of germ cell markers, *Pw-piwi* and *Pw-vasa*. The marker signals were first detected in the distal areas of the parapodia on both sides of each segment and formed a large germ cell cluster in each parapodium. The large germ cell cluster separated into smaller clusters and the small clusters migrated to the deeper body cavity area during growth of animals. Until the female germ cells began vitellogenesis, it was difficult to identify the sex of germ cells by morphological observations. Thus, male and female *P. wilsoni* have the same mechanism of germ cell provision to all segments.

2.2 Introduction

Annelida is a diverse and widely distributed animal phylum. Polychaeta, the largest class of Annelida, comprises two subclasses: Errantia (active scavengers or free-moving worms) and Sedentaria (permanent burrowers). Most polychaete species are marine, although a few are freshwater worms (Hartman, 1969; Fauchald and Rouse, 1997; Bakken and Wilson, 2005; Kriska, 2013). Polychaetes play an important role in the marine ecosystem as a food resource for crustaceans, fishes, and shore birds (Pamungkas and Glasby, 2015). Polychaetes are also the essential food in shrimp aquaculture for induction of shrimp gonadal maturation (Olive, 1999; Binh *et al.*, 2008; Techaprempreecha *et al.*, 2011). Most polychaete species live as scavengers, decomposing and recycling nutrients in the marine ecosystem (Fauchald and Rouse, 1997; Nugteren *et al.*, 2009; Ito *et al.*, 2011).

Due to divergence in ecological features such as their habitats, morphologies, and behaviors, polychaetes have evolved various reproductive strategies including hermaphroditism, viviparity, and external fertilization (Pocklington and Hutcheson, 1983; Bartels-Hardege and Zeeck, 1990; Fong, 1993; Simon *et al.*, 2014). Nereididae, one of several families of polychaetes, has long been studied for its unique reproductive strategy, called epitoky. As the breeding season approaches, a sexually immature worm (atoke) transforms into a sexually mature worm (epitoke) with a characteristic pelagic morph. In time with the spawning, the entire body of the epitoke is filled with mature gametes. Epitokes release eggs or sperm with a typical swimming behavior and die shortly after the spawning (Bartels-Hardege and Zeeck, 1990; Fong, 1993; Fischer, 1999; Hébert *et al.*, 2008).

Despite the intensive studies of the reproductive biology of the polychaete, knowledge about gametogenesis and release is still limited, especially in fully segmented species. The previous study described the oogenesis of *Perinereis wilsoni* (Errantia: Nereididae), which has body segments that are completely compartmentalized by intersegmental septa

(Maceren-Pates *et al.*, 2015). During growth, each new segment is added at the anterior border of the pygidium, the tail end segment. Thus, anterior segments are older than posterior segments (Fischer, 1999; Fischer and Dorrestijn, 2004; Bakken and Wilson, 2005). It was found that the pygidium was the only source of germ cells in *P. wilsoni*, providing germ cells to each new segment at the time of segment generation (Maceren-Pates *et al.*, 2015). This may be a common mechanism of germ cell distribution to each body segment in fully segmented species of polychaetes. Within each segment, the distributed germ cells migrate and settle in distal areas of parapodia at both sides of the segment, then proliferate to form one large germ cell cluster in each parapodium. The cluster separates into smaller clusters, which migrate to a body cavity area to begin meiotic development (Maceren-Pates *et al.*, 2015). However, compared to female germ cell development, male germ cell development has never been well described. Furthermore, the mechanisms of gamete release after epitoky are still unclear. To address these matters, I examined 1) germ cell development in males in comparison to oogenesis using histological analyses and gene expression observations and 2) the characteristics of gamete release in males and females.

2.3 Materials and methods

2.3.1 Animals

P. wilsoni (Polychaeta – Errantia – Phyllodocida – Nereididae) was purchased from a local hatchery in Oita Prefecture and reared in the Fisheries Research Laboratory (FRL) of Kyushu University. Animals were maintained in filtered running seawater with daily feeding. The natural breeding season in Oita Prefecture is in early April, and animals die shortly after spawning in the next Apr. with one-year life cycle (Fig. 12). The embryos and young juveniles were reared at the FRL. Animals grew by the segment addition at the anterior border of the tail-end body segment, pygidium. This extension growth began just after the spawning in April and declined at the end of July. The segment addition was estimated to occur once per day and reached up to 101 ± 7.5 (standard deviation; $n = 30$) segments in an adult. About 15 animals were sacrificed at least once per month for the analysis of germ cell development. Prior to fixation, animals were anesthetized with 0.3% ethylene glycol monohexyl ether in seawater and then fixed in 4% paraformaldehyde in phosphate-buffered saline. The photos used in this manuscript show the representative figures of similar animals.

2.3.2 *In situ* hybridization

Vasa, *nanos* and *piwi* genes have been long used as the germline marker in various annelids species (Cox *et al.*, 1998; Kuramochi-Miyagawa *et al.*, 2004; Rebscher *et al.*, 2007; Giani *et al.*, 2011; Beyret and Lin, 2011; Rebscher, 2014; Özpölat and Bely, 2015; Ponz-Segrelles *et al.*, 2018). In our previous experiment, I used the *vasa* gene to trace germ cells in female *P. wilsoni* (Maceren-Pates *et al.*, 2015). In this experiment, I cloned the *piwi* gene and used to trace germ cells for an improved reliability of germ cell detection. For *Pw-piwi* cloning in this study, primers were designed from the mRNA sequence of a closely related species, *Platynereis dumerilii* (Rebscher *et al.*, 2007) (forward: 5'-accacgactccaagcagaag-3'; reverse:

5'-atcttggtgggcgtattggca-3'). Total RNA was extracted from unfertilized eggs of mature *P. wilsoni* using TRIzol Reagent (Invitrogen) and purified using a RNeasy Kit (Qiagen). cDNA was synthesized using a Prime Script RT-PCR Kit (Takara). Using a pGEM-T Easy Vector System I kit, the 613-bp PCR product was ligated into a plasmid and transformed into *Escherichia coli* DH5. Probes were labelled with digoxigenin and used for whole-mount *in situ* hybridization, as described previously (Maceren-Pates *et al.*, 2015). A cDNA sequence of *Pw-piwi* is registered in DDBJ (Accession No. LC564863).

2.3.3 Histological analysis

After whole-mount *in situ* hybridization, animals were sectioned for histological observations to detect germ cells inside the body segment, as described previously (Maceren-Pates *et al.*, 2015). The same batch of animals were also prepared for staining with hematoxylin and eosin (H&E). Sections were cut at 7 µm.

2.4 Results

2.4.1 Germ cell localization in male *P. wilsoni*

To examine the localization of germ cells in the body segments of males for comparison with the previous observations of female *P. wilsoni*, I used *in situ* hybridization with *Pw-piwi* and *Pw-vasa* anti-sense probes on adult males at first. Although both probes showed similar expression patterns in whole-mount *in situ* hybridization, the *Pw-piwi* probe tended to give a stronger expression compared with the *vasa*-probe. Therefore, I determined to use the *Pw-piwi* probe in this study. The *Pw-piwi* probe was strongly expressed at the distal areas of parapodia on both sides of the segments (Fig. 1). The signal strength gradually decreased from the posterior to the mid-body, then disappeared in the anterior region of the body (Fig. 1). Horizontal sections of worms stained by *in situ* hybridization showed cell clusters at the parapodia in the posterior region, but no signal was detected in the trunk area (Fig. 2A). In the medial body segments, I observed large cell clusters with the *Pw-piwi* signal at the distal areas of the parapodia, and several smaller clusters were separated from the large cluster and several *Pw-piwi* signals in trunk and basal areas of parapodia (Fig. 2B). In the anterior segments, the large cell clusters were not detected in the parapodia, but several signals were observed in the trunk areas of the parapodia and in the deeper body cavity area (Fig. 2C). The *Pw-piwi* signals were decreased at the trunk and deeper body cavity areas compared to the clusters at the distal areas of the parapodia. Figure 3 shows no signal detected with use of a sense probe. *Piwi* signals were still observed at the distal area of the parapodia of the

posterior region even in early March (Fig. 8). At the end of March, only a few segments in the posterior were stained weakly with the *piwi* probe (Fig. 9). No *piwi* signal was detected in the parapodia in the transformed animal (epitoke) in early April (Fig. 10). Moreover, *piwi* was expressed in early stages of germ cell maturation (spermatogonia and spermatocytes) and was undetectable in spermatid and sperm stages (Fig. 11).

2.4.2 Sexual development of gametes in *P. wilsoni*

To study the development and translocation of germ cells, histological observations of *P. wilsoni* were conducted from May to the following April. The germ cell clusters formed in the parapodia divided into smaller clusters and migrated toward deeper areas of the body (Fig. 2B). The migration was observed in the early specimens of whole mount *in situ* hybridization in May, which indicated the migration already began in young juveniles by 40 dpf (Fig. 4), as those were formed earlier than posterior segments. During this migration period, it was not possible to distinguish the sex of each animal because there was no morphological difference in germ cells (Fig. 5A, A').

In September, the female germ cell became discriminable from the male germ cell by its enlargement with an accumulation of vitellogenin (Fig. 5B, B'). After September, 30 more animals were analyzed and were equally divided into males and females; thus, the sex ratio of *P. wilsoni* was assumed to be 1:1. The vitellogenesis began simultaneously in all body segments, but the size of vitellogenic germ cells varied, even within the same segment during this season. However, nearly all eggs were observed to attain equal size just before the spawning season (Fig. 6A). By contrast, male germ cells were observed to have a longer proliferation period (Fig. 5C), which lasted until March, and they entered meiosis at the same time, in late March, in all body segments (Fig. 6B, B'). Figure 13 is the schematic diagram which explains the timing of the germ cells migration and the start of the vitellogenesis.

The developing male germ cells were categorized according to morphological changes in the clusters at each stage (Fig. 7). In the early stage of proliferation, nuclei and nucleoli were clearly visible, and the cytoplasm was filled with granular particles (Fig. 7A). These cells were observed in September and categorized as spermatogonia I. In this stage, the average cell size varied among clusters, even within the same segment. In November, each cluster was filled with proliferating germ cells (Fig. 7B), and by January, the clusters enlarged in size due to the highly proliferating germ cells; the cells decrease in size and the nuclei and nucleoli could not be distinguished due to chromosome condensation. This cluster was categorized as spermatogonia II (Fig. 7C). The enlarged clusters loosened throughout February, divided into smaller clusters (Fig. 7D), and then transitioned into spermatocytes in March (Fig. 7E). In April, a few spermatids were observed as tetrads, although the majority were single sperm cells (Fig. 7F).

Fig. 1. Whole-mount *in situ* hybridization with the *Pw-piwi* probe in an adult male worm in late August. Strong signals were detected at distal ends of the parapodia in a posterior region and decreased toward an anterior region (A–G). White arrows indicate the *Pw-piwi* signals at parapodia (H). Body length = 10 cm. The total number of body segments in this individual is 98. Scale bars for A–G and H are 200 and 100 μ m, respectively.

Fig. 2. Horizontal sections of the hybridized worm showing the localization of the *Pw-piwi* expression in an adult male in late August. (A) Posterior region; strong *Pw-piwi* signals were observed at large cell clusters in distal areas of parapodia. No clusters were detected in trunk areas. (B) Mid region; the *Pw-piwi* signals in distal areas were decreased, and several smaller clusters were detected in trunk areas. (C) Anterior region; the *Pw-piwi* signal was undetectable in distal areas. Smaller clusters were observed in trunk and basal regions of parapodia. Arrows indicate smaller clusters of germ cells. Scale bar, 200 μm .

Fig. 3. Specificity of the *Piwi* probe in *in situ* hybridization. (A) Germ cell clusters were shown by *in situ* hybridization with use of the anti-sense probe of *Pw-piwi*. (B) No signal shown with use of the sense probe. Arrows indicate germ cell clusters.

Fig. 4. Germ cell clusters in a juvenile worm. (A) Germ cell clusters shown by *in situ* hybridization at parapodia of a juvenile worm at 40 dpf. (B) Enlarged parapodia of the posterior region indicated by a black square (b) in Fig. 4A. Germ cell clusters are shown at distal areas of each parapodia. (C) Enlarged parapodia of the anterior region indicated by a black square (c) in Fig. 4A. Germ cell clusters are not observed at parapodia.

Fig. 5. Germ cells before and after the start of vitellogenesis. (A) It is difficult to distinguish sex with morphological characteristics in germ cells of 4-month-old juveniles in early August. (B) Growing female germ cells with vitellogenesis in September. (C) Proliferating male germ cells in clusters in September. A', B', and C' are enlarged images of the frames in A, B and C, respectively. GT: Gut tract. Stained with H&E.

Fig. 6. Germ cells fill body cavities of each segment in March (mid-body region). (A) Female body segments are filled with fully-grown oocytes. (B) Male segments are filled with many germ cell clusters. B' is the enlarged image of the frame in B. Stained with H & E.

Fig. 7. Monthly developments of male germ cells. (A) Early spermatogonia in Sept. (B) Proliferating spermatogonia in Nov. (C) Proliferating spermatogonia in Jan. (D) A large cluster in early Mar. (E) Spermatocytes in middle March. (F) Spermatids and sperms in Apr. (breeding season). N: Nucleus, n: nucleolus, Cy: Cytoplasm. Black arrows in F show a tetrad spermatid and a single sperm. Stained with H & E. Scale bar; 25 μ m.

Fig. 8. Whole mount *in situ* hybridization with *Pw-piwi* in early March. Strong signals were still detected at the distal ends of the parapodia in the posterior region (A), even though germ cell inside the body cavity were in spermatocytes stage. Scale bar; 200 μ m.

Fig. 9. Whole mount *in situ* hybridization of *Pw-piwi* in end of March when the animals were transforming into epitoke. Weak signals were still detected at the distal ends of the parapodia in few segments of the posterior region (A). Scale bar; 200 μ m.

Fig. 10. Whole mount *in situ* hybridization of *Pw-piwi* in April; swimming epitoke. No *piwi* signals detected at the distal ends of the parapodia region (A). Scale bar; 200 μ m.

Fig. 11. *Pw-piwi* expression in monthly development of male germ cells. (A, B) Early spermatogonia in Sept. (C, D) Proliferating spermatogonia in Nov. (E, F) Proliferating spermatogonia in Jan. (G, H) Spermatocytes in middle March. Both genes were undetectable in spermatid and sperm stage. Scale bar A-H, 25µm.

Fig. 12. Schematic diagram on the gametogenic cycle of *P. wilsoni*. (A) Hypothetical germ cells migration from the parapodia to the deeper body cavity in *P. wilsoni*.

Fig. 13. Schematic diagram on the timing of the germ cells migration from the parapodia to the deeper body cavity during segments addition and the start of the vitellogenesis. In April, germ cells from the pygidium are supplied to every newly formed segment and settles at the distal end of the parapodia. May, germ cells in the anterior segments started migration to the deeper body cavity. In August, germ cells in the anterior segments have translocated into the deeper body cavity. However, sexes could not be distinguished due to immature state of the germ cells. In Sept., germ cells have reached the deeper body cavity in almost all body segments and start the vitellogenesis simultaneously. During this during period sexes could be differentiated due to enlargement of the oocytes. Black dots represents the hypothetical migrating germ cell clusters. (AS); Anterior segments.

Table 1. Number of segments in adult worm *P. wilsoni*

Samples	No. of segments in adult worm	Samples	No. of segments in adult worm
1	94	16	98
2	112	17	107
3	104	18	103
4	94	19	97
4	107	20	108
6	99	21	111
7	104	22	114
8	107	23	98
9	107	24	103
10	107	25	100
11	92	26	101
12	84	27	84
13	101	28	98
14	103	29	92
15	91	30	104
Sum =	3024		
Count =	30		
Average =	101		
STDEV =	7.53		

2.5 Discussion

2.5.1 Germ cells localization and gametes production in *P. wilsoni*

In this study, we compared germ cell development between males and females of *P. wilsoni*. *P. wilsoni* grows by adding new body segments at the anterior boarder of the tail-end segment, the pygidium, and all segments are completely separated by intersegmental septa. In the previous report, an interesting mechanism by which germ cells are allocated to all body segments was identified, that is, a newly generating segment receives copies of germ cells from the pygidium, the only source of germ cells in *P. wilsoni* (Maceren-Pates *et al.*, 2015). In each segment, germ cells settled first at the parapodia on each side of the segment and proliferated to form a large cell cluster. Then, the large cluster separated into several smaller clusters, which migrated to the inner body cavity area (Maceren-Pates *et al.*, 2015).

During this period, germ cells proliferated and developed in each segment. In the previous paper, it was found that the large germ cell clusters were already formed in the anterior segments of young juveniles at 10 dpf (Fig. 9 of Maceren-Pates *et al.*, 2015). However, at 40 dpf, the *Pw-piwi* signal disappeared from the parapodia of the anterior segments (Fig. 4). The germ cell clusters translocated into the deeper body cavity area might be difficult to be observed because the *Pw-piwi* signal decreased during migration (Fig. 2). These observations indicate that germ cells proliferate to form a large cluster in the parapodia and then migrate to the inner body cavity area within 30 days after the cluster formation. However, at this stage, it was impossible to distinguish the sex of animals because there were no morphological differences between males and females (Fig. 5A, A').

In September, all germ cells began vitellogenesis in females, which was recognizable by the increase in size (Fig. 5B, B'). By contrast, male germ cells continued proliferation (Fig. 5C, C'). With these differences, animals could be identified as male or female. At this time,

the sex ratio of *P. wilsoni* was 1:1, and it was the same in the breeding season, similar to other species (Reish, 1954, 1957; Barels-Hardege and Zeeck, 1990; Fischer and Dorresteyn, 2004). Before the start of vitellogenesis, although I did not identify the sex of the animals used for the histological analysis, I assumed that the animals included males and females at a ratio of 1:1. Therefore, I may conclude here that there is no clear morphological difference between male and female germ cells in the early developmental phase.

In late September, germ cells had reached the inner body cavity area in all body segments and simultaneously began vitellogenesis. Naturally, the germ cells in the anterior segments reached the body cavity area earlier than those in the posterior segments. However, the anterior germ cells delayed vitellogenesis while waiting for the posterior germ cells to reach the body cavity area. This may suggest that there is some unidentified mechanism to control the initiations of migration and vitellogenesis in female germ cells. As the oocytes reached a maximum size before the spawning season, vitellogenesis seemed to continue until just before spawning. The oocytes varied in size in the early phase of vitellogenesis, and vitellogenesis was not well synchronized among germ cells, even in the same segment (Fig. 5B, B'). This variation in oocyte sizes in the early stages has been observed in several polychaete species (Olive and Garwood, 1981; Fischer, 1974; Golding and Yuwono, 1994). However, as the breeding season approached, all oocytes attained equal size (Fig. 6A).

By contrast, males had a longer germ cell proliferation phase to maximize the number of sperm produced, and they entered meiosis in late March, as shown in Figure 6B & B'. The developing male germ cells in *P. wilsoni* were clearly distinct in morphology between stages (Fig. 7). In many polychaetes, morphological characteristics have been used as the criteria for germ cell development. In *Nereis virens*, spermatogenesis was categorized into five stages based on the morphology of germ cells; spermatogonia I, spermatogonia II, spermatocytes, tetrads, and spermatozoa (Schenk and Hoeger, 2010). In this study, the

observed *P. wilsoni* spermatogenesis stages closely resembled those of *N. virens* (Schenk and Hoeger, 2010). In *P. wilsoni*, male germ cells in September might be categorized as spermatogonia I by clear visibility of big nuclei and nucleoli (Fig. 7A). However, as proliferation continued through January, it resulted in the formation of thick and tightly packed clusters of germ cells, which were categorized as late spermatogonia II (Fig. 7B, C). In February, this tightly packed cluster (spermatogonia II) loosened and began to form several smaller clusters; these were clearly separated by March and were categorized as spermatocytes (Fig. 7D, E). In April, the breeding season, some germ cells were observed in groups of four spermatids, whereas the majority were individual sperm cells (Fig. 7F). However, although the *Pw-piwi* signal was clearly detected in the stages of spermatogonium and spermatocyte, it was significantly decreased in the stages of spermatid and sperm (Fig. 11). The decrease of the *Pw-piwi* signal in the late stages of spermatogenesis especially in sperms, had been also reported in other animals (Lin and Spradling, 1997; Houwing and Berezikov, 2008; Xiao *et al.*, 2013; Hempfling *et al.*, 2017)

In *P. wilsoni* segments addition ceases towards the end of July and by September, most of the germ cells has finished migrating to the deeper body cavity and ready to enter vitellogenesis. However, not all germ cells in the posterior segments translocated to the deeper body cavity, as revealed by the detection of *piwi* signals at the distal area of the parapodia throughout March (Fig. 8). Probably there was not enough time for all the posterior germ cells to migrate since the younger segments were formed lately before the start for the vitellogenesis. *Piwi* signals weakens as the worms transforms into epitoke (Fig. 9). *Piwi* signals were not detectable in epitokes (Fig. 10), the reason for disappearance of signals is still unknown, and however it might be related to the transformation process.

2.6 Conclusion

This study explored the process of the gamete development in details in a fully segmented worm, *Perinereis wilsoni*. The findings provide new insights on understanding the timing of the germ cells translocation from the parapodia to deeper body cavity and the start of gamete maturation. The male has shown similar features of germ cell distribution, proliferation and translocation in each segment to those of the female. Germ cells translocate to the deeper body cavity during growth (segment addition phase). The translocation correlates with age of the segments, however, the further development in the body cavity area starts simultaneously in all body segments and all mature gametes are spawn at one time in the breeding season. This may suggest that there should be under the some regulating mechanism to synchronize gamete development among all segments of the whole body.

CHAPTER 3

Mechanisms of gamete spawning in males and females

Perinereis wilsoni

3.1 Abstract

In the previous studies, I described the germ cell development and gamete maturation in a fully segmented worm. In *P. wilsoni*, germ cells proliferate, differentiate and mature separately in each segment. Like other nereidids, *P. wilsoni* also undergoes transformation into epikoke for the breeding season. The present research aimed to understand the characteristics of gamete releasing sites in males and females by histological techniques. I found that the complete septa were broken, gametes were mixed between segments prior to spawning and released from the posterior region of the body. However, the releasing sites were differently allocated between males and females. Sperms were released from nephridiopores at 2nd through 15th segments from the pygidium, while eggs were released through raptures of skin of 2 – 3 segments between 10th and 30th segments from the tail. The rapturing point varied among females.

3.2 Introduction

Nereidids are errant polychaetes whose spawning behaviours had been long studied due to their unique reproductive strategy called epitoky. Epitoky is a form of reproduction in which sexually immature worm (atoke) transform into sexually matured worm (epitoke) and emerge from benthic to short pelagic life in spawning season (Bartels-Hardege and Zeeck, 1990; Fong, 1993; Franke, 1999; Hébert-Chatelain *et al.*, 2008; Fischer and Dorresteijn, 2004).

Nereidids are broadcasting spawners as many of other marine invertebrates. To achieve successful fertilization, the effective communication among partners is of great importance. Majority of species use the chemical pheromone to insure the synchronous spawning. In polychaetes, the pheromone is found in the coelomic cavity of the ripe epitoke. The pheromone is released and spreads in water, and detected by the opposite partners which will release more pheromones (Jorg *et al.*, 1998). The pheromone is species specific to have the interspecies effects. For instance, Zeeck *et al.*, (1988) identified a ketone compound (5-methyl-3-heptanone) in three species, *P. dumerilii*, *Nereis succinea* and *Pereinereis var. brevicirrus*. The compound induces the swarming behaviour (nuptial dance) and also enables to recognize other individuals of the same species. In *Nereis japonica*, a different ketone compound (3,5-octadiene-2-one) was found which had the same function to that found in *P. dumerilii* (Hardege *et al.*, 1995). Along with the pheromones communication, many species release gametes in strings, streams or plumes with mucus to reduce gametes dilution by rapid and active water mixing (Denny and Shibata, 1989; Olive and Pillai, 1983b; Thomas, 1994; Eckelbarger *et al.*, 2001; Lotterhos and Levitan, 2010). In breeding season, some of the polychaetes such as syllids modify few nephridia of the posterior into gonoduct (metanephromixia) for gametes release (Franke, 1999; Goodrich, 2009). On the other hand,

majority of the nereidids have been reported to release gametes through ruptures of the body wall (Fischer and Dorresteijn, 2004).

Despite the fact that, spawning mechanisms on several nereidids have been explore, the knowledge on gamete releasing sites is still limited especially to a fully segmented worm. The present research was undertaken to characterize the gamete spawning sites in *P. wilsoni*.

3.3 Materials and Methods

3.3.1 Spawning sites observation

Swimming epitokes (2 – 4 males per female) were placed in a transparent container filled with filtered seawater and observed until spawn occurred naturally. This observation was repeated several times to locate the spawning site in the body. Some epitokes were quickly fixed at the beginning of spawning to analyse the spawning site. After collecting fertilized eggs, we reared embryos for further use in the experiments.

3.3.2 Histology

Two sets of samples were collected for histological analysis of internal morphology of the spawning; swimming worms (ready to spawn) and spawning worms. In both cases, males and females were sampled separately. Animals were anesthetized with 0.3% ethylene glycol monohexyl ether in sea water prior to fixation and then fixed in 4% paraformaldehyde in PBS fixed in 4% paraformaldehyde for 24 hrs at 4°C. Fixed animals were cut into about 1cm blocks from the anterior to the posterior, then dehydrated in a series of ethanol (70-100) % and embedded in paraffin. Serial sections were cut in 7 µm and stained with haematoxylin and eosin (H&E) for the microscopic observations.

3.4 Results

3.4.1 Spawning sites in *P. wilsoni*

P. wilsoni body segments are completely separated each other by intersegmental septa. Each segment is filled with proliferated germ cells by the breeding season, and all those germ cells are released during the characteristic swimming. However, it has never been clearly described how *P. wilsoni* releases gametes. To examine the spawning mechanism, we observed the animal behaviour during spawning to identify the gamete releasing site, and also examined the fine structure of the releasing site.

The intersegmental septa were broken during the swimming at spawning and gametes were mixed between segments in male and female (Fig. 2, 6). Moreover, the release of gametes was observed to associate with unidentified structures of varies size which stained dark pink with H &E (Fig. 7).

3.4.2 Gamete spawning sites in male *P. wilsoni*

In male, sperms were continuously released as streams from the several segments in the tail region (Fig. 3A). The total numbers of segments with sperm release were varied among individuals. Those were 10 – 14 segments from 2nd up to 15th segments from the tail end (Fig. 3B). The gamete releasing site were located near the intersegmental septum at the anterior side of each segment (Figs 4A, A'), and a tube like structure from the body cavity was observed connecting to this opening site which could possibly be a “nephridium” (Fig. 4B). The sites were opened/splitted during sperm release (Fig. 4C).

3.4.2 Gamete spawning sites in female *P. wilsoni*

In female, eggs were released from only 2 – 3 segments in the tail region (Fig. 4A). Exact segments were different between individuals, but were in the range of 10th to 30th from the pygidium. As the epithelial structure was severely ruptured by egg spawning, it was difficult to identify the exact point and its detailed structure for egg release (Figs 6B, c, d). Ten

animals were fixed and sectioned to analyse the internal structures for each sex. Figures 3 to 6 show typical examples.

Fig. 1. Observation of spawning sites in *P. wilsoni*. (b) Male (c) female spawning.

Fig. 2. Horizontal sections of a swimming male worm just prior to spawning in April. Arrows indicate broken septa. Stained with H&E.

Fig. 3. Sperm spawning area in the tail region of a male. (A) The square indicates the area of sperm spawning. (B) A horizontal section of the tail indicated in (A). Arrows indicate sperm released from both sides of the limited segments. Stained with H&E.

Fig. 4. Histological observation of sperm release sites in *P. wilsoni*. (A) A horizontal section of sperm-releasing segments. (A') An enlarged sperm release site indicated by the frame in (A). (B) A sperm release site with a tube-like structure indicated by a black arrow. (C) Sperm release from the opening of a structure similar to that shown in (A'). The square indicates the opening point. Stained with H&E.

Fig. 5. Horizontal sections of a swimming female worm just prior to spawning in April. Arrows indicate broken septa. Stained with H&E.

Fig. 6. Histological observation of egg release sites in *P. wilsoni*. (A) The white arrow indicates the segments where eggs were released. (B) The egg release sites are indicated by black frames a, b. Eggs are released from breaks at both sides of the segment. a' and d' are enlarged images of (a) and (b). Black arrows indicate the break points of the egg-releasing segment. Animals were fixed during spawning. Stained with H&E.

Fig. 7. Eosinophilic structure associated with gamete release indicated by arrows. (A) Spawning male. (B) Spawning female. H&E stain.

3.5 Discussion

3.5.1 Gametes spawning sites in males and females *P. wilsoni*

P. wilsoni has the complete septum to separate each segment of the body. Therefore, I had thought *P. wilsoni* would spawn mature gametes from various area of the body. But, in this study, both male and female were found to release gametes from the limited regions in the posterior body (Fig. 3, 6). Males released sperms from 10 – 13 segments and females released eggs from only a few segments. Most germ cells were spawn from those segments during a typical spawning swimming. The spawning swimming was characterized by a strong and flowing serpentine motion of the body. I presume that this strong motion may help to break the intersegmental septa and move germ cells toward the posterior region. This septum breakage looked to precede the germ cell release in both sexes (Fig. 2). If the serpentine motion effectively pushed germ cells into the posterior region, it would increase the internal pressure to break the skin to release germ cells. In female, the germ cell release occurred from only a few neighbouring segments, but those appeared in the back one third region of the body (10 – 30th segments from the tail). This may indicate the egg release site is formed randomly in female. In contrast, in males, sperms were released from about 10 segments in the tail region (Fig. 4). It seemed no shift in location of the release sites among individuals, which suggested there was some common structure related to the sperm release unlike female. There are several precedent papers reporting the site of germ cell release in polychaetes. Most of the nereidids are reported to spawn by rupturing their body wall (Fischer and Dorresteijn, 2004) or by dehiscence (Goodrich, 1895). However, in Syllidae, metanephridia of the fertile segments are modified into metanephromixia during sexual maturity to act as the gonoducts (Franke, 1999; Goodrich, 1895). In this study, I observed that sperms were released by splitting/opening at the “nephridipore” area in *P. wilsoni* (Fig. 4). It was quite similar to Syllidae. However, I could not identify the detailed structures as

metanephromixia. Although these opening sites was basically set in the most segments, the sperm release occurred at the limited “nephridipore” in the tail 10 segments. Therefore, in male, there might be some unknown mechanism involved those limited nephridipore to open for the sperm release.

3.5.2 The eosinophilic structure associated with gamete during spawning

In the present study, I observed irregular structures of different shapes which associated with gametes release. The structures stained dark pink with H&E stain (Fig. 7A, B). In many broadcasting marine invertebrates including polychaetes, the release of gametes had been reported to associate with some adaptive factors which could be biological, chemical or physical factors which offers physical support to minimize gametes dilution in active water movement which will decrease fertilization rates (Denny and Shibata, 1989, Crimaldi and Browning, 2004). In *P. n. brevicirrus* eggs and sperm bond within 2 threads of mucus (Hardege and Bartels-Hargedede, 1996)). In *Phragmatopoma californica*, sperms are released in strings, streams or plumes (Thomas, 1994). Similarly, in the present study the release of gametes were observed to associate with physical elongated structures. In male, the structures infused inside the sperm cluster while in female the structures surrounded eggs and some in between eggs (Fig. 7). The release of gametes in companion with those structures might suggest that, probably they have a supportive role of keeping the gametes intact along the water column to ensure successful fertilization.

3.6 Conclusion

The present study, managed to identify the gamete releasing sites in fully segmented worm. *P. wilsoni* gametes were released from the posterior region regardless of the fact that, during gametogenesis, germ cells develop and mature separately in each segment. Males released sperms from nephridiopores at 2nd through 15th segments from the pygidium, while female released eggs through ruptures of skin of 2 – 3 segments between 10th and 30th segments from the tail. The rupturing point varied in between females.

CHAPTER 4

GENERAL DISCUSSION

4. General discussion

The present study describes the germ cell development in male *Perinereis wilsoni* in comparison with characteristics of the female germ cell studied in the previous study by Mercedes *et al.*, (2015). This study also analysed the mechanisms of gamete spawning in males and females.

4.1 Germ cell localization in male *P. wilsoni* and gamete production

In the previous study in females, it was revealed that the body segments of *P. wilsoni* were separated by the complete intersegmental septa. Germ cells of this species are originated from the pygidium and distributed to each body segment during the segment addition phase and settled at the distal ends of each parapodium on both sides of the segments. During gamete development in females, germ cells at the parapodia proliferated and formed clusters which then separated into a single female germ cell as they moved towards the coelomic cavity where they started vitellogenesis (Maceren Pates *et al.*, 2015). The present study aimed to focus on understanding the germ cell development in male *P. wilsoni* in comparison with the female findings so as to expand the knowledge on reproductive biology of *P. wilsoni* as a representative of fully segmented polychaetes with the complete intersegmental septa.

To find out the localization of germ cells in male, this study analysed a mRNA expression of a germ cell marker gene in adult males of 5 months old, since the sex of this species can be distinguished by the enlargement of germ cells by vitellogenesis. Sampling was carried out through the observation of the coelomic cavity under a microscope and females were distinguished from males by the presence of growing vitellogenic oocytes. Two germline markers were used for whole mount *in situ* hybridization. The *Pw-vasa* was cloned in the previous study held by Maceren Pates *et al.* (2015) and the *Pw-piwi* was cloned in this study (Chapter 2). In the result, both genes showed the similar patterns of expressions as the strong signals at the distal area of the parapodia and the strength of the signal decreased from the

posterior towards the anterior region. These expressions were quite similar to the previous study of females (Maceren Pates *et al.*, 2015). These results indicate that both male and female germ cells in *P. wilsoni* have the same starting points at the distal area of the parapodia. In the posterior region, strong signals of *piwi* were detected only at distal area of the parapodia. In the mid body region, the strength of the signals decreased at the parapodia and several small clusters were detected in the trunk and deeper coelomic cavity. Strong signals at the parapodia and small cluster inside the body cavity suggests that the smaller clusters might have divided from the large cluster at the parapodia and migrated into the deeper body cavity and this migration corresponds with the age of the segment (Chapter 2). To clarify the beginning of germ cell migration and vitellogenesis, the monthly histological observations were conducted through May to September. In May, the *piwi* signals began to decrease first in the anterior body region about 40 days after fertilization. This decreasing of the *piwi* signal occurred as a result of germ cell migration into the inner body cavity area from the parapodia, which was shown by the vertical section of hybridized worms. The migrating germ cells were thought to be in a primordial germ cell stage and could not be distinguished the sex. The migration process continues through the months of June to September. In September, most of the germ cells reached the deeper body cavity area in all segments and starts vitellogenesis, simultaneously. Even though the germ cells in the anterior segments migrated early, they looked to arrest the start of vitellogenesis in Sept. This arrest might be the waiting for the posterior germ cells until their reach in the body cavity area. This arrest of the anterior germ cells may suggest that there is some unknown mechanism to arrest germ cells to enter the vitellogenesis phase. As the beginning of vitellogenesis is related to the cell stage changing from the oogonium to the oocyte, there may be a hormonal control for the arrest of beginning of the vitellogenesis until Sept. (Chapter 2)

In *P. wilsoni*, growth of animals by the segment addition began at Apr. and continued to the end of July. The addition of the segment was one segment per day. While the segment addition continued three months from Apr. to Jul., the adult worm finally reached to have more than 90 segments. The posterior segments were formed by the end of Jul. As the germ cell migration finish within 30 days in the anterior region, the germ cells in the posterior region should also finish the migration into the inner body cavity area before the beginning of vitellogenesis in Sept. However, surprisingly, the *Pw-piwi* signals were still detected at the parapodia of posterior region even in the mid-March. There was an unexpected delay of the germ cell migration forward inner body cavity area in the posterior region. However, the *piwi* signal in the posterior region finally disappeared in early Apr. when the animals transformed into epitokes for spawning. The reason of this delay of germ cell migration in the posterior region is unknown. After the migration, the germ cells were transferred from the primordial germ cell stage to the gonium stage in the inner body cavity area, then changed to the oocyte stage with starting vitellogenesis in females. All these changes of germ cells have been thought under the hormonal control of the central nervous system in various animals. Although there is no any specific evidences, there may be some causal relationship among the delay of germ cell migration in the posterior region and body segmentation in this species. (Chapter 2)

The process of gamete maturation have been documented in many polychaetes (Gambi, 2000; Fischer, 1974). In *P. dumerilii*, the early stage of vitellogenesis was reported to be heterogeneous whereby pre-vitellogenic and vitellogenic oocytes were observed at a particular time. The variation in size gradually disappears as the breeding season approaches, whereby all oocytes attained an equal full-grown size (Fischer, 1974). In the present study, *P. wilsoni* has shown similar vitellogenic characteristics. Although the start of vitellogenesis was simultaneously in all body segments, there were a variation in size of oocytes even in a

same cluster especially in early stages. The enlarged oocytes eventually separated each other and developed freely in the coelomic cavity. The equality in size was gradually observed towards the end of Feb. and all oocytes had grown to full maturity size by the end of March. Unlike in the female, male germ cells had extended a long proliferation period from Sept. to Jan. As in many other polychaetes, male germ cells in *P. wilsoni* proliferated and differentiated within clusters which changed in number and morphology during the maturation progressed. These changes of cluster morphology between each stage have been used as one of criteria in classification of the maturation stages. Based on this criteria, Schenk (2010) categorized five stages of spermatogenesis in *Nereis virens*; spermatogonia I, spermatogonia II, spermatocytes, tetrads and spermatozoa. In present study, the developing male germ cells were observed to have resemblance characteristics to those described in *N. virens*. Spermatogonia I was observed in Sept. when the female germ cells already entered vitellogenesis. Spermatogonia II was observed from Nov. to Jan. when the cell size of germ cells decreased while the number increased many times to form thick and tightly packed clusters of germ cells. Spermatocytes could be seen in Mar. as small clusters resulted from the breakage of the large cell clusters of Spermatogonia II. In Apr., many spermatids were observed as a single cell through the tetrad form stage. (Chapter 2)

4.2 Gamete releasing sites in males and females *P. wilsoni*

Spawning behaviours and gamete releasing mechanisms have been well described in several polychaetes species (Bartels-Hardege and Zeeck, 1990; Fong, 1993; Franke, 1999; Fischer and Dorresteyn, 2004). Gamete maturation is one of the important step towards the creation of new offsprings. Successful gamete release increases the chance of fertilization especially in the broadcasting spawning. The way of gametes release in polychaetes is not described well as in higher animals, although several studies have shown that gametes are

mostly released from the posterior region (Franke, 1999; Goodrich, 2009). For instance, members of the family syllidae change some of the posterior nephridia into metanephromixia which act as the gonoducts during spawning (Franke, 1999). In contrast, most of nereidids are known to release gametes through rupturing/tearing of the body wall in the posterior region (Goodrich, 2009).

Since *P. wilsoni* is a fully segmented worm and germ cells develop and mature separately in each segment, the way of gamete release in this species is quite interested. This uniqueness of the complete segmentation easily comes up an imagination that gametes are released through the ruptures in segments all over the body. To confirm this speculation, at first I carefully observed the spawning behaviour of several spawning males and females to identify the area of gamete releasing. Prior to spawning in many nereidids, the epitokes swarm in the circular motion for several minutes (Bartels-Hardege and Zeeck, 1990; Fong, 1993; Franke, 1999; Hébert *et al.*, 2008; Fischer and Dorresteijn, 2004). In both males and females of *P. wilsoni*, germ cells were released from the posterior region as reported previously, although the releasing sites were in the different location between males and females. At first, males started to release sperms as the milky streams from the posterior segments. After a short while, females spawned eggs as streams from the two ruptured points. Males continued to release sperms with the rapid circulating motion until the end of egg spawning by females. The rupture point of egg release varied among females. The egg spawning continued for 25 seconds, and sperm spawning continued for 40seconds. The longer sperm spawning period by males looked to cover the whole egg release event by females. After the spawning, animals died within a few hours.

Further histological experiments were conducted to specify the exact spawning segments and observe the internal morphology during spawning, including the status of the complete intersegmental septum and the structure of the gamete releasing sites. Histological

observations of the swimming epitokes showed that the septa were broken and gametes were mixed among segments. The breakage of the septa may be caused by the intense swimming behaviours during spawning and the body actions may help germ cells to move forward posterior region of the body. Observation of the gamete releasing area found that, sperms were coming out from the splitting/opening of nephridiopores at 2nd through 15th segments from the pygidium, while eggs released through ruptures of skin of 2 – 3 segments between 10th and 30th segments from the tail.

In conclusion, this study provided the detailed descriptions of the male germ cell development of a fully segmented worm, *Perinereis wilsoni* a nereidid polychaete. These findings were necessary to fulfil the understanding of the *P. wilsoni* gonads locations regarding that the female data had been reported from the previous study. Likewise, information on the timing of the germ cells translocation and the start of vitellogenesis together with the description of the gamete releasing sites have provided the additional knowledge in understanding the reproductive biology of fully segmented worms.

P. wilsoni is a good experimental animal to understand the reproduction of the fully segmented polychaetes. Further analyses are required to better understand the reproduction of this species as follows;

1) Further classification of the specific stages of the migrating germ cells. Currently the specific gametogenic stage of migrating germ cells is still unknown. As for the current data, the migrating germ cells do not look to differentiate sexually. The germ cells become distinguishable in Sept. by showing the different development of vitellogenesis or proliferation in females and males, respectively. The sexual bipotentiality in early stages of germ cells have been reported in many animals. In *P. wilsoni*, the migrating germ cell may also have sexual bipotentiality, and the sex of the germ cell may be determined in Sept. by some signals such as a hormone and other cellular factors. In any event, the identification of

the specific differentiation stage of the migrating germ cells is one of the great issues to be urgently resolved. Some genetic markers such as Oct4 and Stella (Linher *et al.*, 2009) may help to recognize the germ cells on this point.

2) Live imaging study to reveal the germ cell migration. The horizontal section all through the body of one animal from head to tail will show a believable image of germ cell migration. There is another way to show a live image of germ cell migration. A cell permeable fluorescent substrate of alkaline phosphatase is currently used to chase a living primordial germ cell (Hahnel *et al.*, 1990; Linher *et al.*, 2009). The reagent can be injected into juveniles of 1-4 months old to chase the germ cells as they are in a period of active germ cell migration and also suitable to fluorescent observation because of the low density of body pigments. The direct tracing of living germ cells will help to analyse the details of germ cell migration.

3) Histological and physiological mechanisms involved in the formation of gamete releasing sites. In *P. wilsoni*, the sperm releasing site is very possibly the nephridium shown by this study and former studies. However, it is still unknown why the limited segments in a posterior region will involve in sperm release. As the sperms are pushed out by the rising internal body pressure, the nephridia involving sperm release must be ready to break just before spawning. Further comparative analyses between normal and sperm-releasing nephridia will show the mechanism of the formation of sperm releasing sites.

ACKNOWLEDGEMENT

I would like to take this opportunity to express my deepest gratitude to my thesis supervisor, Dr. Michiyasu Yoshikuni, Professor, Kyushu University Fisheries Research Laboratory (FRL) for his timeless supervision and excellent guidance throughout my research period and thesis writing. Indeed, his continuous support and constructive suggestions have been of great contribution to the completion of the present thesis.

I am thankful to Dr. Norio Onikura, Professor, FRL Kyushu University and Dr. Noritaka Mochioka, Associate Professor, Animal and Marine Bioresource Science Laboratory, Kyushu University for their evaluation, comments and suggestions to improve the quality of the present thesis.

I am grateful to Dr. Yoshihisa Kurita, Assistant Professor, FRL Kyushu University for his kind feedbacks, suggestions and encouragements during laboratory research and thesis writing.

I thank Dr. Masanori Sato and Dr. Hiroaki Tosuji, Professors at Kagoshima University for their excellent support to analyse DNA for species identification.

I thank Dr. Mercedes Maceren-Pates, former member of the FRL for teaching the molecular and histology techniques and for her friendly support. I extend my thanks to Mr. Gaudioso Pates Jr, PhD student FRL, for sharing his research experience and kind assistance in rearing polychaetes.

I wish to thank all staffs of the FRL, Mr. Masaaki Tsujigawa, and all FRL students for their friendly support and cooperation during laboratory research and my stay in Japan.

I would like to give my sincere thanks to the Ministry of Education, Culture, Sports and Technology (MEXT) for the financial provision which enabled me to pursue my study smoothly.

I thank Mr. Yasunobu Masuo, the owner of the polychaete hatchery in Oita Prefecture Japan for the supply of animals during the full course of my research study.

My special thanks to my family for their love, encouragement and support.

Last but not least, I am thankful to almighty God for giving me strength, courage and the ability to undertake my research. Thank you God.

REFERENCES

- Aller RC (2001). Transport and reactions in the bioirrigated zone, in the Benthic Boundary Layer. Transport Processes and Biogeochemistry. B Boudreau and B Jorgensen eds. Oxford University Press, oxford: 269–301
- Bakken T and Wilson RS (2005). Phylogeny of nereidids (Polychaeta, Nereididae) with paragnaths. Zool Scripta 34: 507–547
- Bartels-Hardege HD and Zeeck E (1990). Reproductive behaviour of *Nereis diversicolor* (Annelida: Polychaeta). Mar Biol 106: 409–412
- Beyret E and Lin H (2011). Pinpointing the expression of piRNAs and function of the PIWI protein subfamily during spermatogenesis in the mouse. Dev Biol 355: 215–226
- Bentley MG, Pacey AA (1992). Physiological and environmental control of reproduction in polychaetes. Oceanogr Mar Biol Ann Rev 30: 443–481
- Bentley MG, Olive PJW, Garwood PR and Wright NH (1984). The spawning and spawning mechanism of *Nephtys caeca* (Fabricius, 1780) and *Nephtys hombergi* Savigny, 1818 (Annelida: Polychaeta), Sarsia, 69: 63–68
- Binh NT, Shikawa M, Yokoyama S, Michael FR, Sakiyama K and Koshio S (2008). Effects of Polychaete Meal Supplementation to the Maturation Feed on Kuruma Shrimp (*Penaeus japonicus*) Female Broodstocks. Aquaculture Sci 56: 523–530

Brown- Gilpin L, (1959). The reproduction and larva development in *Nereis fucata* (Savigny). J Mar Bio Associ UK 38: 65–80

Bryan GW and Gibbs PE (1987). Polychaetes as indicators of heavy-metal availability marine deposits, p. 37–49. In J.M Capunzo and D. R Kester. Oceanic processes in Marine Pollution I. Biological Processes and Wasters in the Ocean Krieger Publishing, Florida, USA

Hasse C, Rebscher N, Reiher, W, Sobjinski K, Moerschel E, Beck L, Tessmar-Raible K, Arendt D and Hassel, M (2010). Three Consecutive Generations of Nephridia Occur during Development of *Platynereis dumerilii* (Annelida, Polychaeta). Developmental Dynamics 239: 1967–1976.

Cox DN, Chao A, Baker J, Chang L, Qiao D and Lin H (1998). A novel class of evolutionarily conserved genes defined by piwi are essential for stem cell self-renewal. Genes Dev 12: 3715–3727

Dauvin JC nad Thiébaut E (1994). Is *Owenia fusiformis* Delle Chiaje a cosmopolitan species? Memories du Museum National d'Histoire Naturelle 162: 383–404

Dannenberg LC, Seaver EC (2018). Regeneration of the germline in the annelid *Capitella teleta*. Dev Biol 440: 74–87

Denny MW and Shibata MF (1989). Consequences of Surf-Zone Turbulence for Settlement and External Fertilization. The American Naturalist, 134: 859–889

- Eckelbarger KJ, Young CM, Llodra RE, Brooke S and Tyler P (2001). Gametogenesis, spawning behavior, and early development in the “iceworm” *Hesiocaeca methanicola* (Polychaeta): Hesionidae) from methane hydrates in the Gulf of Mexico. *Mar Biol* 138: 761–775
- Eckelbarger KJ (1975). A light and electron microscope investigation of gametogenesis *Nicolea zostericola* (Polychaeta: Terebellidae). *Mar Biol* 30: 353–370
- Eckelbarger KJ (1983). Evolutionary radiation in polychaete ovaries and vitellogenic mechanisms: their possible role in life history patterns. *Can J Zool* 61: 487–504
- Fauchald K and Rouse G (1997). Polychaete systematics: Past and present. *Zool Scripta* 26: 71–138
- Fischer A (1974). Stages and stage distribution in early oogenesis in the Annelid, *Platynereis dumerilii*. *Cell Tissue Res* 156: 35–45
- Fischer A (1999). Reproductive and developmental phenomena in annelids: a source of exemplary research problems. *Hydrobiologia* 402: 1–20
- Fischer A and Dorresteijn A (2004). The polychaete *Platynereis dumerilii* (Annelida): a laboratory animal with spiralian cleavage, lifelong segment proliferation and a mixed benthic /pelagic life cycles. *BioEssays* 26: 314–25

- Fong PP (1993). Lunar control of epitokal swarming in the polychaete *Platynereis bicanaliculata* (Baird) from Central California. Bull Mar Sci 52: 911–924
- Franke HD (1999). Reproduction of the Syllidae (Annelida : Polychaeta). Hydrobiologia 402: 39–55
- Garwood PR (1980). The role of temperature and day length in the control of the reproductive cycle of the *Harmothoe imbricate* (L) (Polychaeta: Polynoidae). J Exp Mar Biol Ecol 47: 35–53
- Giani VC, Yamaguchi E, Boyle MJ and Seaver EC (2011). Somatic and germline expression of piwi during development and regeneration in the marine polychaete annelid *Capitella teleta*. Evo Devo 2: 10
- Giangrande A, Cavallo A, Licciano M, Mola E, Pierri C and Trianni L (2005). The utilization of the filter feeder polychaete *Sabella spallanzanii* (Sabelledae) as bioremediator in aquaculture. Aquaculture International. 13: 129–136
- Glasby JC and Timm T (2008). Global diversity of polychaetes (Polychaeta; Annelida) in freshwater. Hydrobiologia 595: 107–115
- Glasby J Christopher and Hwey- Lian Hsieh (2006). New species and new records of the *Perinereis nuntia* species group. (Nereididae: Polychaeta) from Taiwan, and other Indo-west pacific shores. Zoological Studies 45: 553–577

- Golding DW (1967). Endocrinology, regeneration and maturation in *Nereis*. Biol Bull Mar Biol Lab Woods Hole 133:567–577
- Golding DW and Yuwono E (1994). Latent capacities for gametogenic cycling in the semelparous invertebrate *Nereis*. Proc Natl Acad Sci USA 91: 11777–11781
- Goodrich E (1945). The study of nephridia and genital ducts since 1895. Q J Microsc Sci 86: 113–301
- Hartman O (1969) Atlas of Sedentariate Polychaetous Annelids from California, Allan Hancock Foundation, University of Southern California, Los Angeles
- Hardege JD and Bartels-Hargedde HD, (1996). Spawning behaviour and development of *Perinereis nuntia* var. *brevicirrus*. Invert Bio 114:39–15
- Hardege DJ, Müller TC, Beckmann M, BartelShardege DH and Bentley GM (1998). Timing of reproduction in marine polychaetes: The role of sex pheromones, Écoscience, 5: 395–404
- Hahnel CA, Rappolee DA, Millan LJ, ManesT, Ziomek AC, Theodosiou NG, Werb Zena, Pendersen AR and Gilbert A (1990). Two alkaline phosphatase genes are expressed during early development in the mouse embryo. Development 564: 555–564.
- Hébert Chatelain É, Breton S, Lemieux H and Blier PU (2008). Epitoky in *Nereis* (*Neanthes*) *virens* (Polychaeta: Nereididae): A story about sex and death. Comp Biochem and Physiol B: Biochem Mol Biol 149: 202–208

- Hutchings PA (1998). Biodiversity and functioning of polychaetes in benthic sediments. *Biodiversity & Conservation* 7: 1133–1145
- Ito K, Nozaki M, Kunihiro T, Miura C, Miura T (2011) Study of sediment cleanup using polychaetes. In “Interdisciplinary Studies on Environmental Chemistry-Marine Environmental Modeling & Analysis”, Ed by K Omori, X Guo, N Yoshie, N Fujii, IC Handoh, A Isobe, S Tanabe, TERRAPUB, Tokyo, pp133–139
- Jansen HM, Hansen PK, Brennan N, Dahlgren TG, Fang J, Nederlof MAJ, Strohmeier T, Strand Ø (2019). Enhancing opportunistic polychaete communities under fish farms : an alternative concept for integrated aquaculture. 11: 331–336
- Kozin VV and Kostyuchenko P (2015). Vasa, PL10 and Piwi gene expression during caudal regeneration of the polychaete *Allita virens*. *Dev Genes Evol* 225: 129–138
- Kriska G (2013) Segmented worms – Annelida. In “Fresh Water Invertebrates in Central Europe. A field Guide”, Springer-Verlag, Wien, pp 104–115
- Kumar SS, Reena S, Perumal M, Samikkannu M and Kandaramy K (2017). Biochemical studies on the live feed polychaete, *Nereis* sp, in relation to maturity stages, 46: 591–596
- Kuramochi-Miyagawa S, Kimura T, Ijiri TW, Isobe T, Asada N, Fujita Y, Ikawa M, Iwai N, Okabe M, Deng W, Lin H, Matsuda Y and Nakano T (2004). Mili, a mammalian member of piwi family gene, is essential for spermatogenesis. *Development (Cambridge)* 131: 839–849

- Kunihiro T, Miyazaki T, Uramoto Y , Kinahita K, Inoue A, Tamaki S, Hama D, Tsutsuki H and Ohwada K (2008). The succession of microbial community in the organic rich fish-farm sediment during bioremediation by introducing artificially mass-cultured colonies of a small polychaete, *Capitella sp. I*, Mar Pollut Bull 57: 68–77
- Lawrence A, (1996). Environmental and Endocrine Control of Reproduction in Two Species of Polychaete: Potential Bio-Indicators for Global Climate Change. J Mar Biol Associ UK 76(1): 247–250.
- Lawrence AJ and Soame JM (2009). The endocrine control of reproduction in Nereidae: A new multi-hormonal model with implications for their functional role in a changing environment. Philos Trans R Soc London B: Biol Sci 364: 3363–3376
- Licciano M, Stabili L, Giangrande A and Cavallo AR (2007). Bacterial accumulation by *Branchiommma luctuosum* (Annelida: Polychaeta): A tool for biomonitoring marine systems and restoring polluted waters. 63: 291–302
- Lin H and Spradling AC (1997). A novel group of pumilio mutations effects of asymmetric division of the germ cells in the Drosophila Ovary. Development 124(12): 2463–2476
- Linhaer K, Dyce P and Li J (2009). Primordial Germ Cell-Like Cells Differentiated In Vitro from Skin-Derived Stem Cells. PLoS One 4(12)
- Lotterhos EK and Levitan RD (2010). Gamete Release and Spawning Behavior in Broadcast Spawning Marine Invertebrates. The Evolution of Primary Sexual Characters in Animals pp 99–120

- Maceren-Pates M, Kurita Y, Pates G and Yoshikuni M (2015). A model for germ cell development in a fully segmented worm. *Zool Letters* 1: 34
- Newell RIE, Hilbish TJ, Koehn RK, Newell CJ, (1982). Temporal variation in the reproductive cycle of *Mytilus edulis* L (Bivalvia, Mytilidae) from localities on the coast of the United States. *Biol Bull* 162: 229–310
- Nugteren PV, Herman PMJ, Moodley L, Middelburg JJ, Vos M and Heip CHR (2009). Spatial distribution of detrital resources determines the outcome of competition between bacteria and a facultative detritivorous worm. *Limnol Oceanogr* 54: 1413–1419
- Olive PJW (1999). Polychaete aquaculture and polychaete science: a mutual synergism. *Hydrobiologia* 402: 175–183
- Olive PJW and Garwood PRG (1981). Gametogenic cycle and population structure of *Nereis* (Hediste) *diversicolor* and *Nereis* (Nereis) *pelagica* from north-east England. *Journal of the Mar Biol Assoc UK* 61: 193–213
- Olive PJW and Bentley MG (1980). Hormonal control of oogenesis, ovulation and spawning in the annual reproductive cycle of the polychaete, *Nephtys hombergii* Sav (Nephtyidae), *Intl J Invert Reprod.* 2: 205–221
- Olive PJW and Pillai G (1983b). Reproductive biology of the polychaete *Kefersteinia cirrata* *Keferstein* (Hesionidae). II. The gametogeneic circle and evidence for photoperiodic control of oogenesis. *Intl J Invert Reprod*, 6: 295–306

- Oliver SJ (1984). Selection for asexual reproduction in an Antarctic polychaete worm. *Mar. Ecol. Prog. Ser* 19:33–38
- Özpolat BD and Bely AE (2015). Gonad establishment during asexual reproduction in the annelid *Pristina leidy*. *Dev Biol* 405: 123–136
- Pamungkas J and Glasby CJ (2015). Taxonomy of reproductive Nereididae (Annelida) in multispecies swarms at Ambon Island, Indonesia. *ZooKeys* 520: 1–25
- Pfannenstiel Hans-Dieter and Grünig Charlotte (1982). Primordial germ cells and early stages of oogenesis in *Ophryotrocha puerilis* (Polychaetes, Dorvillidae). *Zoomorphology*, 100: 203–215
- Pfannenstiel Hans-Dieter and Grünig Charlotte (1990). Spermatogenesis and sperm ultrastructure in the polychaete genus *Ophryotrocha* (Dorvilleidae) 171: 159–171
- Pierre F (1932). Annelida polychaeta of the Indian museum, Calcutta. *Memoirs of the Indian museum, Calcutta*. 12:1–262
- Ponz-Segrelles G, Bleidorn C and Aguado MT (2018). Expression of vasa, piwi and nanos during gametogenesis in *Typosyllis antoni* (Annelida, Syllidae). *Evol Dev* 20: 132–145
- Pocklington P and Hutcheson MS (1983). New record of viviparity for the dominant benthic Invertebrate *Exogone hebes* (Polychaeta: Syllidae) from the Grand Banks of Newfoundland. *Mar Ecol Prog Ser* 11: 239–244

- Rebscher N, Zelada-González F, Banisch TU, Raible F and Arendt D (2007). Vasa unveils a common origin of germ cells and of somatic stem cells from the posterior growth zone in the polychaete *Platynereis dumerilii*. *Dev Biol* 306: 599–611
- Rebscher N (2014). Establishing the germline in spiralian embryos. *Int J Dev Biol* 58: 403–411
- Reish (1954). The life history and ecology of the polychaetous annelid *Nereis grubei* (Kinberg). *Allan Hancock Found Occas* 14
- Reish DJ (1957) The life history of the polychaetous annelid *Neanthes caudata* (delle Chiaje), including a summary of development in the family Nereidae. *Pac Sci* 11: 216–228
- Willian SR, (1997). The control of oogenesis in *Nereis* (*Neanthes*) *virens* (Sars). Interactions of environmental and endogenous mechanisms. The University of Newcastle upon Tyne
- Rice SA and Eckerlbarger JK, (1989). An ultrastructural investigation of spermatogenesis in the holopelagic polychaetes *Vanadis formosa* and *Krohnia lepidota* (Polychaeta: Alciopidae): *Biol Bull* 176: 123–134
- Schenk S and Hoeger U (2010). Lipid accumulation and metabolism in polychaete spermatogenesis : Role of the large discoidal lipoprotein. *Mol Reprod Dev* 77: 710–719

- Simon C, Martín GS and Robinson G (2014). Two new species of *Syllis* (Polychaeta: Syllidae) from South Africa, one of them viviparous, with remarks on larval development and vivipary. *J Mar Biol Assoc UK* 94: 729–746
- Stabili L, Schirosi R, Lucciano M, Mola E, Giangrande A (2010). Bioremediation of bacteria in aquaculture waste using the polychaetes *Sabella spallanzanii*. *New Biotechnology*. 27:774–81
- Shoobs FN, Schmidt ER and McMullin RE (2016). A new record of the freshwater polychaete *Namanereis hummelincki* (Polychaeta: Nereididae) from epigeal waters of Montserrat, (January)
- Techaprempreecha S, Khongchareonporn N, Chaicharoenpong C, Aranyakananda P, Chunhabundit S and Petsom A (2011). Nutritional composition of farmed and wild sandworms, *Perinereis nuntia*. *Anim Feed Sci Technol* 169: 265–269
- Thomas FIM (1994). Transport and mixing of gametes in three free-spawning polychaete annelids, *Phragmatopoma californica* (Fewkes), *Sabellaria cementarium* (Moore), and *Schizobranchia insignis* (Bush). *J Exper Mar Biol Ecol* 179: 11–27
- Weigert A, Helm C, Hausen H, Zakrzewski AC, Bleidorn C (2013). Expression pattern of Piwi-like genes in adult *Myzostoma cirriferum* (Annelida). *Dev Genes Evol* 223(5): 329–334
- Wilson RS (1993). Systematic of the *Perinereis nuntia* complex (Polychaeta: Nereididae)

from south-eastern Australia. Records of the Australian museum. 45: 241–252

Zeeck E, Hardege J, Bartels-Hardege H and Wesselmann G (1988). Sex pheromone in a marine polychaete: determination of the chemical structure. J Exp Zool 246: 285–292