

Development and evaluation of the Sedentary Behavior and Light-Intensity Physical Activity Questionnaire

田中, るみ

<https://hdl.handle.net/2324/4474993>

出版情報 : Kyushu University, 2020, 博士 (看護学), 課程博士
バージョン :
権利関係 :

Doctoral Dissertation

Title: Development and evaluation of the Sedentary Behavior and Light-Intensity Physical Activity Questionnaire

Department of Health Sciences Graduate School of
Medical Sciences Kyushu University

Rumi Tanaka

Supervisor: Kimie Fujita

2021

Abstract

[Purpose]

To describe our newly developed Sedentary Behavior and Light-Intensity Physical Activity Questionnaire and examine its reliability and validity.

[Participants and Methods]

We identified and selected self-reported items through a literature review and interviews with 11 inactive individuals. Thirty-one individuals with lower limb prostheses and an expert panel assessed the content validity of the integrated items and identified 17 items. Patients who had undergone lower limb surgeries were regarded as inactive individuals, and 112 patients completed the questionnaire twice for test-retest reliability and wore an accelerometer for criterion validity. The ethics committee of Kyushu University approved this study (2019-126 and 2019-273).

[Results]

Item analysis was revised to the Sedentary Behavior and Light-Intensity Physical Activity Questionnaire-10 (six light-intensity physical activity and four sedentary behavior items) because of the floor effect. The test-retest correlation coefficient showed high reliability. Moderate to weak correlation coefficient was observed between the questionnaire and accelerometer (light-intensity physical activity: 0.43 and sedentary behavior: 0.20), and the Bland–Altman plots indicated no bias.

[Conclusion]

The Sedentary Behavior and Light-Intensity Physical Activity Questionnaire-10 had acceptable validity and reliability among inactive individuals and it could be used for studying light-intensity physical activity.

Key words: physical activity; sedentary; validity

INTRODUCTION

Physical activity is classified into three categories depending on the intensity of the activity: sedentary behavior, light-intensity physical activity, and moderate- to vigorous-intensity physical activity (MVPA)¹⁾. There is strong evidence of an association between MVPA and health benefits²⁾. However, some people have factors, such as health problems, feeling tired, and lacking motivation, that prevent them from participating in physical activity³⁾ and people with mobility limitations were found to be more sedentary than those without⁴⁾. Long-term sedentary behavior causes muscle weakness, metabolic dysfunction, cardiovascular disease^{5, 6)}, and increased mortality⁷⁾.

The focus of physical activity health benefits has expanded to include light-intensity physical activity, which is easy to conduct even for people who face barriers to physical activity. In recent years, several studies have revealed that light-intensity physical activity has a positive effect not only physically, by preventing non-communicable diseases and improving lipid and glucose metabolism⁸⁾, but also psychosocially, by decreasing depression⁹⁾. Furthermore, light-intensity physical activity decreases the risk of physiological indicators such as the number of co-morbidities¹⁰⁾. Three hundred minutes per week of light-intensity physical activity is recommended¹¹⁾.

Although most light-intensity physical activity studies used accelerometers^{8, 12-14)}, these devices needed to be worn continuously, which sometimes resulted in a high attrition rate¹⁵⁾. A questionnaire is a suitable measurement method for a large sample survey as it is inexpensive and versatile¹⁶⁾. Only two scales have focused on light-intensity physical activity measurements to date. The 7-day Sedentary and Light Intensity Physical Activity Log was the first scale developed for evaluating light-intensity physical activity and sedentary behavior¹⁷⁾; however, its adaptability to middle-aged and older people has not yet been established because the mean age of survey participants was 26.5 years. Second, the Community Health Activities Model Program for Seniors,

a self-report scale developed as a physical activity measurement scale for older individuals, has recently been modified to measure light-intensity physical activity^{18, 19)} but was found to have low validity. A systematic review reported that scales to measure physical activity included a limited number of light-intensity physical activity²⁰⁾, and validity and/or reliability of these scales was only tested in the general population not in physically inactive people²¹⁻²³⁾.

Patients with lower limb arthroplasty have lower levels of physical activity than individuals in the general population, even long after surgery^{24, 25)}, and could be regarded as inactive people. We aimed to develop a questionnaire to characterize sedentary or light-intensity physical activity and to examine its reliability and validity in middle-aged and older adults who tend to have an inactive lifestyle.

PARTICIPANTS AND METHODS

To develop the Sedentary Behavior and Light-Intensity Physical Activity Questionnaire (SLPAQ), we extracted 265 light-intensity physical activity and sedentary behavior items from the 2011 Compendium of Physical Activities (Japanese version)²⁶⁾. It contains common activities converted into metabolic equivalent units (METs)^{26, 27)}. Then, a panel of experts, comprising physical activity researchers, clinical nurses, and orthopedic surgeons, chose 163 activities that were thought to be performed by inactive adults out of the 265 light-intensity and sedentary activities.

To identify additional activities, we conducted a review of literature on physical activity scales. We identified 25 light-intensity and sedentary activities from 22 studies. To identify any unreported physical activity items in community-dwelling middle-aged and older persons, we recruited a convenience sample of 11 community-dwelling persons 40–90 years old (age: 65.8 ± 13.5 years; 6 males and 5 females) who did not exercise regularly. We interviewed the participants regarding their activities over the previous 4 days. We identified 18 additional activities from this survey. The

expert panel grouped the 206 activities by similarity into 58 items, comprising five domains: housework, leisure, work, transport, and self-care.

To confirm the questionnaire's content validity—its ability to capture activities that accurately represent sedentary behavior and light-intensity physical activity—we recruited patients with artificial lower limb joints from Kyushu University Hospital and expert panel members from the previous survey. We asked 31 community-dwelling outpatients aged 40 years or older with artificial lower limbs joints (age: 68.3 ± 12.9 years; 14 males and 17 females) to report how often they performed each of the 58 activities using a 4-point Likert scale, ranging from 1 (very infrequent) to 4 (very frequent). Items for which >50% of responses reported 4 (very frequent) were included in the draft version of the scale. Then, 7 expert panel members participated in an item content validity index survey, using a 4-point Likert scale ranging from 1 (not relevant) to 4 (very relevant), and a content validity ratio survey, using a 3-point Likert scale ranging from 1 (not necessary) to 3 (essential)²⁸). We used an item content validity index cut-off point of 0.8 and a content validity ratio cut-off point of 0.99, where a value greater than the cut-off indicated acceptable validity²⁹). Of 58 items, 17 activity items were retained—the SLPAQ-17.

The SLPAQ-17 asked the frequency that each physical activity had been performed in the previous week and the duration of each physical activity per day; the total time spent on light-intensity physical activity and sedentary behavior per week could then be calculated. Item responses for activity duration had five categorical options (the algorithm conversion values are included in parentheses): <1 h (0.5 h), 1–2.5 h (1.75 h), 3–4.5 h (3.75 h), 5–6.5 h (5.75 h), >7 h (7.75 h). For activities typically performed for longer durations (e.g., lying down or sitting down), items used the same format with eight categories: <1 h (0.5 h), 1–2.5 h (1.75 h), 3–4.5 h (3.75 h), 5–6.5 h (5.75 h), 7–8.5 h (7.75 h), 9–10.5 h (9.75 h), 11–12.5 h (11.75 h), >13 h (13.75 h).

We compared the performance reliability of two new and conventional physical activity monitoring accelerometers, the Active style PRO HJA-750 (Omron Healthcare Co., Ltd., Kyoto, Japan³⁰) and Lifecorder (Suzuken Co., Ltd., Nagoya, Japan). Active style PRO captures physical activity and classifies it into walking physical activity and non-walking physical activity³⁰. We recruited 22 individuals (age: 57.6 ± 17.4 years; 13 males and 9 females; BMI 22.0 ± 2.8 kg/m²) from the general population. After the individuals were informed about the study, those who agreed to participate provided signed consent forms. Each was asked to simultaneously wear an Active style PRO accelerometer and a Lifecorder accelerometer. The intraclass correlation coefficient (ICC) for Active style PRO walking activity detection and Lifecorder physical activity detection were 0.71 for light-intensity physical activity and 0.79 for MVPA.

To test the validity and reliability of the SLPAQ-17, we selected the Active style PRO accelerometer because it was more sensitive to light-intensity physical activity than the Lifecorder accelerometer, which tended to underestimate non-walking activities^{31, 32}) such as standing, sitting while doing laundry, or dishwashing³³). Active style PRO captured these light-intensity physical activities and correctly classified them as non-walking.

To test the validity and reliability of the SLPAQ-17, patients with lower limb arthroplasty were recruited from Kyushu University Hospital. Participants who had undergone hip or/and knee arthroplasty ≥ 6 months prior and who were aged between 40 and 90 years were eligible. The planned sample size was 100 participants, in accordance with Consensus-based Standards for the Selection of Health Measurement Instruments³⁴). Eligible outpatients who consented to participate were given two copies of the SLPAQ-17 questionnaire and an Active style PRO accelerometer. The participants were asked to complete the questionnaire (Time 1) and wear the accelerometer for 4 days, except while sleeping. The test–retest reliability of the questionnaire survey was conducted 1 week after the Time 1 survey (Time 2). The questionnaires and accelerometer were returned by

mail. Data from those who wore the accelerometer ≥ 10 h/day for at least 4 days were included in the analysis⁸⁾. Physical activity intensities were defined as follows: 0.9 MET < sedentary ≤ 1.5 METs, 1.5 < light intensity ≤ 3 METs, and MVPA > 3 METs.

Item analyses assessed ceiling and floor effects of the scale items. When >15% of participants had the highest possible value (>7 or >13 h \times 7 days) or the lowest possible value (0 h \times 7 days), the ceiling or floor effect, respectively, was considered to be present.

Test-retest reliability was assessed using Spearman rank correlation coefficients (ρ) and ICCs between Time 1 and Time 2 SLPAQ variable. Spearman rank correlation and Bland-Altman analyses of SLPAQ and accelerometer-based results (h/day) were used to evaluate item content validity. For testing proportional biases, Pearson correlation of the differences between and averages of the activity durations estimated by the accelerometer and SLPAQ were calculated.

Correlation coefficients (Spearman ρ , Pearson and ICC) were calculated using SPSS version 24 (IBM Corp, Armonk, NY), and Bland-Altman analyses were performed in Excel (Microsoft Inc). The level of significance for all statistics was set at $p < 0.05$. This study was approved by the ethics committee of Kyushu University (first SLPAQ version: 2019-126; validation and reliability study: 2019-273).

RESULTS

Overall, 138 patients were asked to participate in the survey. Of these, 13 declined to participate. The response rate for the SLPAQ-17 was 96.8%. Out of 121 respondents, 9 failed to wear the accelerometer. The remaining 112 patients participated in the validation study. The mean age of the participants was 67.5 ± 10.6 years, and 79 were female. The mean BMI was 24.5 ± 3.9 kg/m², and most participants had undergone total hip arthroplasty ($n = 100$, 89.3%). All participants had

received non-cemented prostheses. The mean number of steps was 4644 ± 3187 steps/day. The mean durations of MVPA, light-intensity physical activity, and sedentary behavior were 0.9 ± 0.5 h/day, 4.9 ± 1.6 h/day, and 6.6 ± 1.8 h/day, respectively. The proportions of MVPA, light-intensity physical activity, and sedentary behavior were 7.3%, 39.6%, and 53.1%, respectively.

No ceiling effect was observed, while a floor effect was demonstrated for 11 items. Of these 11 items, 2 work and 2 transport activity items were retained in the questionnaire because a previous study had classified physical activity in work and transportation in general people as crucial³⁵.

These four retained items and the six items without floor effects were included in the final version of the questionnaire, SLPAQ-10 (Table 1), comprising six light-intensity physical activity and four sedentary behavior items.

[Table 1 near here]

Table 2 shows test–retest reliability and correlations between Time 1 and Time 2 of the SLPAQ-10 data. Time 1 and Time 2 showed high correlations for light-intensity physical activity items ($\rho = 0.74$, $p < 0.01$) and moderate correlation for sedentary behavior items ($\rho = 0.66$, $p < 0.01$). There was also high to moderate agreement for light-intensity physical activity (ICC = 0.70) and sedentary behavior (ICC = 0.69).

Table 2 reports Spearman ρ for the SLPAQ-10 and accelerometer variables. A moderate and significant correlation between the SLPAQ-10 and light-intensity physical activity accelerometer variables ($\rho = 0.43$, $p < 0.01$) was observed. Correlation values for SLPAQ-10 and sedentary behavior accelerometer variables were low ($\rho = 0.20$, $p = 0.03$). Bland–Altman plots demonstrated that zero was within the 95% confidence interval (CI) of the mean difference between SLPAQ-based light-intensity physical activity (mean difference; 95% CI = 0.51; -0.10 to 1.12 h/day) and sedentary behavior (mean difference; 95% CI = 0.10; -0.76 to 0.79 h/day) and the accelerometer variables, indicating no bias (Fig. 1). Pearson correlation coefficients for differences and averages

of the accelerometer and the SLPAQ-10 variable were 0.69 ($p < 0.01$) and 0.64 ($p < 0.01$) for light-intensity physical activity and sedentary behavior, respectively.

[Table 2 and Fig. 1 near here]

DISCUSSION

We developed a self-report questionnaire—SLPAQ-10—to measure sedentary behavior and light-intensity physical activity among middle-age and older adults who had undergone lower limb arthroplasty and who mainly perform low-intensity activities. The current validation study showed the SLPAQ-10 had good reliability and validity compared with an accelerometer.

The SLPAQ-10 has some advantages over previous scales used to measure physical activity. First, the SLPAQ-10 had a high validity for light-intensity physical activity, while a systematic review found that most scales measuring physical activity that include light-intensity physical activity have a low validity for light-intensity physical activity²⁰). Second, the SLPAQ-10 is short and easy to complete; this study had a high response rate. Conversely, many physical activity scales require that participants report activity times in detail^{17, 21, 36}), which places a time burden on the participants.

Although there are light-intensity physical activity studies using accelerometers^{8, 12-14}), participant dropout is common because of the inconvenience of wearing an accelerometer¹⁵). For example, Matsunaga et al. researched physical activity using accelerometers and health-related quality of life using questionnaires in patients 5 years after total hip arthroplasty for; the dropout rate for physical activity measurement was 64.3% and that for the questionnaire was 43.0%³⁷). The SLPAQ-10 can be used in long-term physical activity follow-up studies.

In this study, the participants were inactive individuals because patients with lower limb arthroplasty mostly engage in light-intensity physical activity and sedentary behavior. Although

previous studies of light-intensity physical activity scales involved individuals from the general population^{17,19, 21-23}), our questionnaire could be applied to populations that find performing MVPA difficult. Meanwhile, a survey comparing the physical activity of community-dwelling adults aged ≥ 50 years reported that participants who spend more time performing light-intensity physical activity also tend to spend more time performing MVPA³⁸). The use of an additional MVPA questionnaire, such as the International Physical Activity Questionnaire²⁰), to capture MVPA is recommended as a complement to the SLPAQ-10 for physical activity studies.

The importance of light-intensity physical activity in health promotion has been gaining acceptance in recent years. According to a nationwide survey in the United States, 300 min/week of light-intensity physical activity are recommended for older adults¹¹). In an accelerometer-based retrospective survey in the United States, all-cause mortality risk was reduced by 14% for every increase of 60 min/day of light-intensity physical activity in those who self-reported that MVPA was difficult or impossible to perform without assistance¹⁰); therefore, light-intensity physical activity instead can be effective for people who have difficulty performing high-intensity physical activity, and the SLPAQ-10 is useful for capturing light-intensity physical activity. In an intervention study to decrease sedentary behavior, people with rheumatoid arthritis were instructed to increase the amount of light-intensity physical activity and reported a significant decrease in sedentary behavior³⁹). Though this study measured sedentary behavior with an accelerometer, our questionnaire could be used in such intervention studies.

While the SLPAQ-10 had higher Spearman correlations than those demonstrated by previous scales²⁰), and Bland–Altman plots demonstrated that 0 was within the 95% CI, Bland–Altman plots also showed proportional biases. This indicates the SLPAQ-10 variables were higher than the accelerometer variables. Additionally, in a previous study, participants tended to over-report physical activity because of factors such as social desirability⁴⁰).

This study has a couple of limitations. Sedentary behavior measured by the SLPAQ-10 had a low correlation with that measured by the accelerometer. However, sedentary behavior scales often show no significant correlation or low correlation with accelerometer data because of factors such as immobilization time (0 METs) unmeasured as sedentary behavior (1~1.5 METs) by the accelerometer^{20, 41}).

The generalizability of the SLPAQ-10 to all middle-aged and older adults with an inactive lifestyle is limited because this study was based on patients who had undergone hip and/or knee total joint replacement surgery. It will be necessary to investigate the adaptability of the SLPAQ-10 to inactive people other than patients with hip or/and knee total joint replacements.

Acknowledgments

We are much indebted to the patients who participated in this study as well as to Satoshi Hamai, PhD; Yukio Akasaki, PhD; Hideki Mizu-uchi, PhD; Shinya Kawahara, PhD; and Hidetoshi Tsushima, MD in the Department of Orthopedic Surgery and the Nursing Department at Kyushu University Hospital. This study was funded by the Grants-in-Aid for Scientific Research (Grant Number Research (C) 19K11141). We thank Coren Walters-Stewart, PhD, from Edanz Group (<https://en-author-services.edanzgroup.com/ac>) for editing a draft of this manuscript.

Conflict of interest

There are no conflicts of interest to declare.

REFERENCES

- 1) National University of Singapore [Internet]. What is MVPA?; [2020.08.01]. Available from: http://www.nus.edu.sg/uhc/docs/default-source/default-document-library/what-is-mvpa.pdf?sfvrsn=59cfa1de_2
- 2) World Health Organization. Global recommendations on physical activity for health. Geneva: WHO Press; 2010.
- 3) Justine M: Barriers to participation in physical activity and exercise among middle-aged and elderly individuals. 2013, 54: 581–586.
- 4) Loprinzi PD, Sheffield J, Tyo BM, et al.: Accelerometer-determined physical activity, mobility disability, and health. *Disabil Health J*, Elsevier Inc, 2014, 7: 419–425.
- 5) Tremblay MS, Colley RC, Saunders TJ, et al.: Physiological and health implications of a sedentary lifestyle. *Appl Physiol Nutr Metab*, 2010, 35: 725–740.
- 6) Booth FW, Roberts CK, Thyfault JP, et al.: Role of inactivity in chronic diseases: Evolutionary insight and pathophysiological mechanisms. *Physiol Rev*, 2017, 97: 1351–1402.
- 7) Martinez-Gomez D, Guallar-Castillon P, Rodríguez-Artalejo F: Sitting Time and Mortality in Older Adults With Disability: A National Cohort Study. *J Am Med Dir Assoc*, Elsevier Inc., 2016, 17: 960.e15-960.e20.
- 8) Füzéki E, Engeroff T, Banzer W: Health benefits of light-intensity physical activity: a systematic review of accelerometer data of the national health and nutrition examination survey (NHANES). *Sport Med*, 2017, 47: 1769–1793.
- 9) Buman MP, Hekler EB, Haskell WL, et al.: Objective light-intensity physical activity associations with rated health in older adults. *Am J Epidemiol*, 2010, 172: 1155–1165.
- 10) Frith E, Loprinzi PD: Accelerometer-assessed light-intensity physical activity and mortality among those with mobility limitations. *Disabil Health J*, Elsevier Ltd, 2018, 11: 298–300.

- 11) Loprinzi PD, Lee H, Cardinal BJ: Evidence to support including lifestyle light-intensity recommendations in physical activity guidelines for older adults. *Am J Heal Promot*, 2015, 29: 277–284.
- 12) Spartano NL, Davis-Plourde KL, Himali JJ, et al.: Association of Accelerometer-Measured Light-Intensity Physical Activity With Brain Volume: The Framingham Heart Study. *JAMA Netw open*, 2019, 2: e192745.
- 13) Stubbs B, Chen L, Chang C, et al.: Accelerometer-assessed light physical activity is protective of future cognitive ability : A longitudinal study among community dwelling older adults. *Exp Gerontol*, Elsevier Inc., 2017, 91: 104–109.
- 14) Seol J, Abe T, Fujii Y, et al.: Effects of sedentary behavior and physical activity on sleep quality in older people: A cross-sectional study. *Nurs Heal Sci*, 2020, 22: 64–71.
- 15) Pedi Ž, Bauman A: Accelerometer-based measures in physical activity surveillance : current practices and issues. *Br J Sports Med*, 2015, 49: 219–223.
- 16) Pols MA, Peeters PHM, Kemper HCG, et al.: Methodological aspects of physical activity assessment in epidemiological studies. *Eur J Epidemiol*, 1998, 14: 63–70.
- 17) Barwais FA, Cuddihy TF, Washington T, et al.: Development and Validation of a New Self-Report Instrument for Measuring Sedentary Behaviors and Light-Intensity Physical Activity in Adults. *J Phys Act Heal*, 2014, 11: 1097–1104.
- 18) Stewart AL, Mills KM, King AC, et al.: CHAMPS Physical Activity Questionnaire for. 1989.
- 19) Hekler EB, Haskell WL, Cain KL, et al.: Reliability and Validity of CHAMPS Self-Reported Sedentary-to-Vigorous Intensity Physical Activity in Older Adults. *J Phys Act Heal*, 2016, 9: 225–236.

- 20) Helmerhorst HJF, Brage S, Warren J, et al.: A systematic review of reliability and objective criterion-related validity of physical activity questionnaires. *Int J Behav Nutr Phys Act*, 2012, 9: 103.
- 21) Nicolaou M, Gademan MGJ, Snijder MB, et al.: Validation of the SQUASH physical activity questionnaire in a multi-ethnic population: The HELIUS study. *PLoS One*, 2016, 11: 1–15.
- 22) Strath SJ, Bassett DR, Swartz AM: Comparison of the College Alumnus Questionnaire Physical Activity Index with objective monitoring. *Ann Epidemiol*, 2004, 14: 409–415.
- 23) Chinapaw MJM, Sloopmaker SM, Schuit AJ, Van Zuidam M, Van Mechelen W. 2109 Reliability and validity of the activity questionnaire for adults and adolescents (AQuAA). *BMC Med Res Methodol*. 2009;9.
- 24) King LK, Kendzerska T, Waugh EJ, et al.: Impact of Osteoarthritis on Difficulty Walking: A Population-Based Study. *Arthritis Care Res*, 2018, 70: 71–79.
- 25) Arnold JB, Walters JL, Ferrar KE: Does physical activity increase after total hip or knee arthroplasty for osteoarthritis? A systematic review. *J Orthop Sports Phys Ther*, 2016, 46: 431–442.
- 26) Ministry of Health, Labour and Welfare [Internet]. Revised version of Codes and MET Values; 2012 [2020.08.01]. Available from: <https://www.nibiohn.go.jp/files/2011mets.pdf>. Japanese.
- 27) Ainsworth BE, Haskell WL, Herrmann SD, et al.: 2011 compendium of physical activities: A second update of codes and MET values. *Med Sci Sports Exerc*, 2011, 43: 1575–1581.
- 28) Zamanzadeh V, Ghahramanian A, Rassouli M, et al.: Design and Implementation Content Validity Study: Development of an instrument for measuring Patient-Centered Communication. *J Caring Sci*, 2015, 4: 165–178.

- 29) Davis LL: Instrument review: Getting the most from a panel of experts. *Appl Nurs Res*, 1992, 5: 194–197.
- 30) Ohkawara K, Oshima Y, Hikiyara Y, et al.: Real-time estimation of daily physical activity intensity by a triaxial accelerometer and a gravity-removal classification algorithm. *Br J Nutr*, 2011, 105: 1681–1691.
- 31) Hikiyara Y, Tanaka S, Ohkawara K, et al.: Validation and comparison of 3 accelerometers for measuring physical activity intensity during nonlocomotive activities and locomotive movements. *J Phys Act Heal*, 2012, 9: 935–943.
- 32) Matthews CE: Calibration of accelerometer output for adults. *Med Sci Sports Exerc*, 2005, 37.
- 33) Oshima Y, Kawaguchi K, Tanaka S, et al.: Classifying household and locomotive activities using a triaxial accelerometer. *Gait Posture*, 2010, 31: 370–374.
- 34) COSMIN: COSMIN checklist with 4-point scale. *Cosmin*, 2011, 6.
- 35) Pettee Gabriel KK, Morrow JR, Woolsey ALT: Framework for physical activity as a complex and multidimensional behavior. *J Phys Act Health*, 2012, 9 Suppl 1: 11–18.
- 36) Hargreaves MK, Cohen SS, Matthews CE, et al.: Evaluation of a Questionnaire to Assess Sedentary and Active Behaviors in the Southern Community Cohort Study. *J Phys Act Heal*, 2016, 9: 765–775.
- 37) Matsunaga-Myoji Y, Fujita K, Makimoto K, et al.: Three-Year Follow-Up Study of Physical Activity, Physical Function, and Health-Related Quality of Life After Total Hip Arthroplasty. *J Arthroplasty*, Elsevier Ltd, 2020, 35: 198–203.
- 38) Blodgett J, Theou O, Kirkland S, et al.: The association between sedentary behaviour, moderate-vigorous physical activity and frailty in NHANES cohorts. *Maturitas*, Elsevier Ireland Ltd, 2015, 80: 187–191.

- 39) Thomsen T, Aadahl M, Beyer N, et al.: Sustained long - term efficacy of motivational counselling and text message reminders on daily sitting time in patients with rheumatoid arthritis? Long - term follow - up of a randomized, parallel - group trial. *Arthritis Care Res (Hoboken)*, 2019, 0–2.
- 40) Adams SA, Matthews CE, Ebbeling CB, et al.: The Effect of Social Desirability and Social Approval on Self- Reports of Physical Activity. *Am J Epidemiol*, 2005, 161: 389–398.
- 41) Van Cauwenberg J, Van Holle V, De Bourdeaudhuij I, et al.: Older adults' reporting of specific sedentary behaviors: Validity and reliability. *BMC Public Health*, 2014, 14: 1–10.

Legends to Figures

Figure 1. Bland–Altman plots of accelerometer and Light-Intensity Physical Activity (upper panel) and Sedentary Behavior (lower panel) Questionnaire (SLPAQ) variables

Table 1. Physical activity measures derived from Time 1 survey using the SLPAQ (h/day)

Table 2. Reliability of the SLPAQ-10 administered at Time 1 and Time 2 and validity coefficients for the SLPAQ-10 and accelerometer




Figure .1 Bland-Altman plots of the accelerometer variables and the Sedentary and Light Intensity Physical Activity Questionnaire (SLPAQ)

Table 1. Physical activity measures derived from Time 1 survey using the SLPAQ (h/day)

		n = 112		
Items		Mean ± SD	Ceiling effect (%)	Floor effect (%)
Housework	Preparing and clearing meals [†]	2.2 ± 1.8	0.0	7.1
	Cleaning up [†]	0.7 ± 0.8	0.0	13.4
Leisure	Lying down or sitting down [‡]	4.1 ± 3.2	5.4	0.0
Work	Sitting at work [‡]	1.0 ± 2.1	0.0	67.9
	Standing or walking slowly at work [†]	0.8 ± 1.9	0.0	78.6
Transport	Transporting to destination [†]	0.7 ± 1.1	0.0	33.0
	Sitting in a vehicle [‡]	0.3 ± 0.6	0.0	52.7
Self-care	Eating meals [‡]	1.2 ± 0.8	0.0	4.5
	Morning preparation [†]	0.7 ± 0.5	0.0	0.0
	Preparation and tidying up before going to sleep [†]	0.4 ± 0.2	0.0	9.8

※SLPAQ: Sedentary Behavior and Light Intensity Physical Activity Questionnaire, SD: standard deviation, Ceiling and floor effect: the percentage of the people scoring highest (more than 7 or 13 hours × 7 days) or lowest (0 hours × 7 days) respectively, †Light intensity physical activity, ‡Sedentary behaviour

Table 2. Reliability of the SLPAQ-10 administered at Time 1 and Time 2 and validity coefficients for the SLPAQ-10 and accelerometer

n = 112						
	Reliability: the correlation between Time 1 and Time 2 of the SLPAQ			Validity : the correlation between Accelerometer physical activity and the SLPAQ		
	Spearman's ρ	p value	ICC	Spearman's ρ	p value	ICC
Light-intensity physical activity	0.74	< 0.01	0.70	0.43	< 0.01	0.32
Sedentary behavior	0.66	< 0.01	0.69	0.20	0.03	0.10

※ SLPAQ: Sedentary Behavior and Light Intensity Physical Activity Questionnaire

※Spearman's ρ: Spearman's rank correlation coefficient, ICC: Intraclass Correlation Coefficient