

(E)-N-(3,4-dimethylisoxazol-5-yl)-4-((4-hydroxy-2-oxo-2H-chromen-3-yl)diazenyl)Benzenesulfonamide; A New Synthetic Anti-Herpetic Agent for HSV-1

Kumar, Ashish

Department of Applied Science & Humanities, G.L. Bajaj Institute of Technology & Management

Brij Kishore Tiwari

Department of Applied Science & Humanities, G.L. Bajaj Institute of Technology & Management

Mishra, Mamta

Department of Applied Science & Humanities, G.L. Bajaj Institute of Technology & Management

Kumar, Navneet

Department of Applied Science & Humanities, Inderprastha Engineering College

<https://doi.org/10.5109/4480717>

---

出版情報 : Evergreen. 8 (2), pp.351-357, 2021-06. 九州大学グリーンテクノロジー研究教育センターバージョン :

権利関係 : Creative Commons Attribution-NonCommercial 4.0 International

# (E)-N-(3,4-dimethylisoxazol-5-yl)-4-((4-hydroxy-2-oxo-2H-chromen-3-yl) diazenyl) Benzenesulfonamide; A New Synthetic Anti-Herpetic Agent for HSV-1

Ashish Kumar<sup>1,\*</sup>, Brij Kishore Tiwari<sup>1</sup>, Mamta Mishra<sup>1</sup>, Navneet Kumar<sup>2</sup>

<sup>1</sup>Department of Applied Science & Humanities, G.L. Bajaj Institute of Technology & Management, Greater Noida-201308, India

<sup>2</sup>Department of Applied Science & Humanities, Inderprastha Engineering College, Ghaziabad, India

\*Author to whom correspondence should be addressed:

E-mail:ashish2009chemistry@gmail.com

(Received November 9, 2020; Revised March 18, 2021; accepted April 9, 2021).

**Abstract:** (E)-N-(3,4-dimethylisoxazol-5-yl)-4-((4-hydroxy-2-oxo-2H-chromen-3-yl) diazenyl) benzene sulfonamide (HDF) was prepared by condensation of 4-hydroxy coumarin and 4-amino-N-(3,4-dimethylisoxazol-5-yl)benzene sulfonamide (sulphafurazole). The structure of synthesized compound was established using IR, <sup>1</sup>H and <sup>13</sup>C-NMR spectral analysis. On evaluation of *in-vitro* anti-herpetic activity of this compound against *HSV-1*, compound was found able to inhibit viral replication at late stages of infection by inhibiting the replication of HSV-1. This anti herpetic nature of HDF was evidenced by the reduction of specific gene products  $\beta$ -(ICP-6) and  $\gamma$ -(ICP-5 and gB) groups in western blotting.

**Keywords:** coumarin-sulphonamide, 4-hydroxy coumarin, HSV-1, immunoblot analysis, post infection

## 1. Introduction

Herpes Simplex Virus type-1 (HSV-1), an associated member of herpes viridae community, is an enveloped DNA virus that causes many types of contagious viral infections like cold sores.<sup>1)</sup> It is possible to prevent viral infection by administration of the vaccine, which is an antibody required for the organism's immune system to detect the virus or treat the disease. Some types of infections, such as HSV type-1 for which no vaccines are available, requiring the development of antiviral agents.<sup>2)</sup>

Various chemical substances are admissible to treat the infections begetted by Herpes Simplex Virus type-1 and for medication of such viral infections. Acyclovir (ACV) and its nucleoside derivatives such as Panivciclovir, Valacyclovir and Famciclovir are used. The mechanism of these antiviral agents is stationed on the inhibition of DNA polymerase enzyme of virus, which avert the replication of DNA of virus.<sup>3)</sup> Usually these antiviral agents require activation by phosphorylation which is relying on viral and kinases of host cell and capable to eliminate replication of virus in host cell.<sup>4)</sup> The long time uses of these compounds as antiviral agents can show the possibility to develop the immunity by virus for these chemical compounds.<sup>5)</sup> These limitations therefore highlight the need to develop new anti-HSV agents based on alternative mechanism of action.

Some synthetic and natural coumarin (2H-1-benzopyran-2-one) derivatives have outstanding chemical reactivity and bioactivities and they are able to show a paramount role to amend various viral infections.<sup>6)</sup>

4-Hydroxy coumarin and its various derivatives has proven a central ingredient in fabrication of an efficacious anti-coagulant drug named as Warfarin. The addition of aryl group to carbon of 4-hydroxy coumarin moiety is vastly pronounced in demonstrating capacious biological actions namely antiviral<sup>7)</sup> and antibacterial<sup>8)</sup> activities. In previous studies the substitution of aryl or azo group in 4-Hydroxy coumarin moiety has been reported which indicates admirable antimicrobial activities.<sup>9)</sup> Some thiazole, pyridine, triazole and phenyl thiazole derivatives of 4-hydroxy coumarin was synthesised by J. Sahoo *et al* and evaluate their *in-vitro* antibacterial activities against pathogens.<sup>10)</sup>

A Sabt *et al.* synthesized coumarin-6-sulfonamide derivatives and investigated their anti-proliferative nature and conclude that all synthesized compounds was able to diminish the growth of cancer cell by blocking topoisomerase-2 enzyme while doxorubin used as reference drug.<sup>11)</sup> So it can be stated that coumarin-sulphonamide derivatives can be acts as a compounds of biological interest.<sup>12)</sup>

Present paper focus on the synthesis of

coumarin-sulphonamide derivative by reaction with 4-hydroxy coumarin and sulphafurazole. The structure of synthesized compound established by using Infrared,  $^1\text{H}$  and  $^{13}\text{C}$ -N.M.R. spectroscopic methods and evaluation of its *in-vitro* antiviral activity against HSV-1 along with determination of its  $\text{CC}_{50}$  and  $\text{EC}_{50}$  value

## 2. Material and Methods

Sodium nitrite, Sulphafurazole, 4-hydroxy coumarin, and sodium hydroxide, hydrochloric acid, methanol, ethanol and Isopropanol, Dulbecco modified Eagle medium, Fetal Bovine Serum, Phosphate Buffer Saline, Glutamine, Sodium pyruvate, Penicillin and Streptomycin used in synthesis were purchased from Sigma Aldrich. All solvents were purified by distillation method. Lymph Buffer solution, Phenyl methacophenol Fluoride, Triton Solution, Protein antibodies ICP-5,6,8, anti gb and  $\beta$ -actin were purchased from Merck-BDH.


Melting point was determined by open capillary method in sulphuric acid bath. Elemental analysis for new synthesized compound was performed on a Perkin-Elmer series-II, 2400 elemental analyzer. Infra-Red spectra was recorded using Nucon IR spectrophotometer using potassium bromide pellets.

The  $^1\text{D}$  &  $^2\text{D}$   $^1\text{H}$  and  $^{13}\text{C}$ -N.M.R spectra were recorded by using Bruker Avance 500 spectroscope. Mixture of  $\text{CDCl}_3$  and  $\text{DMSO}-d_6$  was used as solvent and TMS as reference compound. The chemical shifts of synthesized compound were delineated on  $\delta$  scale/ppm.

## 3. Synthesis of HDF

4-(chlorodiazanyl)-N-(3,4-dimethylisoxazole-5-yl)benzenesulphonamide (II) was obtained by adding 3 mMol (0.207 gram) cold aqueous solution of sodium nitrite added in acidic solution of 3 mMol (0.801 gram) 4-amino-N-(3,4-dimethyl-isoxazol-5-yl) benzene sulphonamide (Sulphafurazole) (I) in ice bath at  $5^\circ\text{C}$  temperature.

Further mix this solution with 20 mL alkaline solution of 4-hydroxy coumarin (III) which prepared by dissolving 3mMol (0.545 gram) 4-hydroxy coumarin in 20 mL of 15% lye solution. Whole mixture was placed in ice bath for 45 minute at  $5^\circ\text{C}$ . Pale yellow crystal of (E)-N-(3,4-dimethylisoxazol-5-yl)-4-((4-hydroxy-2-oxo-2H-chromen-3-yl)diazanyl)benzene sulfonamide (HDF) (IV) was filtered and washed with demineralized water.<sup>13)</sup> (Scheme-1)



### 3.1 Microanalysis of HDF

Pale Yellow crystal, Melting point  $235-241^\circ\text{C}$ , Yield (80%), Analysis for  $\text{C}_{20}\text{H}_{16}\text{N}_4\text{O}_7\text{S}$  (molar mass 456) Calculated % C = 52.63, H = 3.53, N = 12.28, S = 7.03, O = 24.54, Found % C = 52.83, H = 3.51, N = 12.26, S = 7.02, O = 24.56, Soluble in DMSO, Acetone,  $m/z$  = 456 (100%), 457 (4.9%), 458 (3.9%), 459 (1.1%),

IR Spectral data:  $3435-3438\text{cm}^{-1}$  (-NH group),  $3179-3180\text{cm}^{-1}$  (-OH group),  $1689\text{cm}^{-1}$  (-C=O of Lactone carbonyl),  $1272\text{cm}^{-1}$  (O-C=O of coumarin),  $1585\text{cm}^{-1}$  (-N=N-),  $1129$  &  $1302\text{cm}^{-1}$  (symmetric and asymmetric vibration of  $\text{SO}_2$  group),  $1647\text{cm}^{-1}$  (methoxazole ring).

The  $^1\text{H}$ -N.M.R spectroscopic data of HDF which numbered as Figure-1, show the presence of one broad singlet at downfield value 4.0 ppm for proton adjacent with nitrogen of  $-\text{SO}_2\text{NH}-$  moiety present in HDF. The doublet of doublet appeared in aromatic region at 7.91 (2H,  $J$  = 6.6, 1.8, 0.5 Hz) ascribed to H-17 and H-13 show meta coupling with each other, likewise a new doublet of doublet appeared at 7.51 (2H,  $J$  = 6.6, 1.8, 0.5 Hz) due to H-16 exhibiting meta coupling with H-14. Moreover two sharp singlet of three protons embarked at downfield value 2.27 ppm and 2.25 ppm assigned to two methyl group existent in compound and the existence of all these characteristic proton signals reveals the presence of N-(3,4-dimethyl isoxazol-5-yl)benzene sulphonamide moiety in HDF. Further sharp singlet of single proton appeared at 15 ppm represents -OH group while four doublet of doublet appeared at 7.59 ppm ( $J$  = 8.3, 1.3, 0.5 Hz), 7.52 ppm ( $J$  = 8.3, 7.3, 1.5 Hz), 7.59 ppm ( $J$  = 8.3, 1.3, 0.5 Hz) and 8.13 ppm ( $J$  = 7.2, 1.6, 0.5 Hz) assigned to H-26, H-27, H-28 and H-29 of hydroxy coumarin ring present in HDF. (Figure-2)




Fig.1: Numbering of HDF




Fig.2:  $^1\text{H}$ -NMR chemical shift ( $\delta$ /ppm) of HDF

The  $^{13}\text{C}$ -NMR spectroscopic analysis of HDF (Figure-3) show signal of 20 carbon atom appeared at 159.9, 100.5, 158.9 ppm ascribed to C-3, C-4 and C-5 atom of N-(3,4-dimethyl isoxazole-5-yl)benzene sulphonamide moiety and two sharp singlet appeared at 11.1 and 7.5 ppm that two methyl group substituted on C-3 and C-4 of such group. Aromatic carbon signal appeared at 139.7 ppm (C-12), 127.3 ppm (identical C-13 and C-17 atom), 129.1 ppm (identical C-14 and C-16 atom) and 131.9 ppm (C-15) assigned to N-(3,4-dimethyl isoxazol-5-yl)benzene sulphonamide moiety present in HDF. The carbon signal appeared at 164.9 ppm (C-20), 156.2 ppm (C-21), 150.2 ppm (C-23), 117.5 ppm (C-24), 82 ppm (C-25), 121.5 ppm (C-26), 128.4 ppm (C-27), 125.5 ppm (C-28) 126.5 ppm (C-29) strongly suggested the presence of 4-hydroxy coumarin ring in HDF which was further confirmed by  $^1\text{H}$ - $^{13}\text{C}$ -HSQC spectrum




Fig.3:  $^{13}\text{C}$ -NMR chemical shift ( $\delta$ /ppm) of HDF

$^1\text{H}$ - $^1\text{H}$  COSY spectra of HDF show all  $^1\text{H}$ - $^1\text{H}$  scalar coupling which is more significant being H-13 $\leftrightarrow$ H-17, H-14 $\leftrightarrow$ H-16, H-26 $\leftrightarrow$ H-28 and H-27 $\leftrightarrow$ H-29 and all protons of  $-\text{CH}_3$  group (Figure-4)




Fig.4:  $^1\text{H}$ - $^1\text{H}$  coupling in HDF

The  $^1\text{H}$ - $^{13}\text{C}$  HSQC spectra of HDF show that out of twenty carbon atom, nine carbon atom were associated with coumarin ring, six were associated with benzene sulphonamide moiety and other five were of 3,4-dimethylisoxazole moiety present in HDF. Such spectra show coupling of H-13 ( $\delta_{\text{H}}$  7.91) with C-13 ( $\delta_{\text{C}}$  127.3), H-14 ( $\delta_{\text{H}}$  7.51) with C-14 ( $\delta_{\text{C}}$  129.1), H-16 ( $\delta_{\text{H}}$  7.51) with C-16 ( $\delta_{\text{C}}$  129.1), H-17 ( $\delta_{\text{H}}$  7.91) with C-17 ( $\delta_{\text{C}}$  127.3), H-7<sub>a</sub>, H-7<sub>b</sub>, H-7<sub>c</sub> ( $\delta_{\text{H}}$  2.27) with C-7 ( $\delta_{\text{C}}$  7.5), H-6<sub>a</sub>, H-6<sub>b</sub>, H-6<sub>c</sub> ( $\delta_{\text{H}}$  2.27) with C-6 ( $\delta_{\text{C}}$  11.1) in N-(3,4-dimethyl isoxazol-5-yl)benzene sulphonamide moiety, while H-26 ( $\delta_{\text{H}}$  7.59) with C-26 ( $\delta_{\text{C}}$  121.5), H-27 ( $\delta_{\text{H}}$  7.52) with C-27 ( $\delta_{\text{C}}$  128.4), H-28 ( $\delta_{\text{H}}$  7.59) with C-29 ( $\delta_{\text{C}}$  125.5), H-29 ( $\delta_{\text{H}}$  8.13) with C-26 ( $\delta_{\text{C}}$  126.8) in coumarin ring present in HDF, which was further confirmed by broad range correlation of C-H shown in Figure-2. On analyzing the  $^1\text{H}$ - $^{13}\text{C}$  HMBC correlation (Figure-5) observe that H-26 ( $\delta_{\text{H}}$  7.59) show cross peak with C-21 ( $\delta_{\text{C}}$  156.2) and C-25 ( $\delta_{\text{C}}$  82) further H-29 ( $\delta_{\text{H}}$  8.13) show cross peak with C-20 ( $\delta_{\text{C}}$  164.9) and C-23 ( $\delta_{\text{C}}$  150.2) and H-27 ( $\delta_{\text{H}}$  7.52) cross peak with C-24 ( $\delta_{\text{C}}$  117.5) in coumarin moiety present in HDF, other long range correlation of H-7<sub>a</sub>, H-7<sub>b</sub>, H-7<sub>c</sub> ( $\delta_{\text{H}}$  2.27) with C-5 ( $\delta_{\text{C}}$  158.9) and H-6<sub>a</sub>, H-6<sub>b</sub>, H-6<sub>c</sub> ( $\delta_{\text{H}}$  2.27) with C-3 ( $\delta_{\text{C}}$  159.9) and the existence of N-(3,4-dimethyl isoxazol-5-yl)benzene-sulphonamide moiety was confirmed by long range coupling of H-8 ( $\delta_{\text{H}}$  4.0) with C-17 ( $\delta_{\text{C}}$  127.3) and H-13 ( $\delta_{\text{H}}$  7.91) with C-5 ( $\delta_{\text{C}}$  158.9) <sup>14,15</sup>




Fig.5:  $^1\text{H}$ - $^{13}\text{C}$  HMBC coupling in HDF

## 4. Evaluation of antiviral activity of HDF

### 4.1 Cell line, HSV-1 Culture and Extent of infection

Sample Vero Cells of Cercopithecus aethiops were multiplied in Dulbecco modified Eagle Medium containing 10% fetal bovine serum and medium was supplemented in 2 mmole glutamine and 1 mmole sodium pyruvate alongwith antibiotic penicillin (100 unit/mL) and streptomycin (100  $\mu\text{g mL}^{-1}$ ). The viability of Vero cells were appraised by employing 0.02 % Trypan blue exclusion assay. After infection all infected Vero cells were centrifuged to remove cellular debris and remaining viral titer was used to measure the plaques formed by using plaque number reduction method. <sup>16)</sup>

### 4.2 Determination of $\text{CC}_{50}$ value of HDF

Infected Vero cells of Cercopithecus aethiops were

incubated with HDF at varied concentration of 01-500  $\mu\text{g mL}^{-1}$  in Dulbecco's modified Eagle Medium for 24 hours and add 50  $\mu\text{L}$  volume of MTT solution with 100  $\mu\text{L}$  of acidic solution of isopropanol (N/10 HCl dissolve in 10% isopropanol).

At room temperature after ensuring complete disposal of formazon crystals, all petri plates were placed in automatic plate reader at 690 nm with reference wavelength and 570 nm test wavelength and absorbance was observed.<sup>17)</sup>

$\text{CC}_{50}$  value was narrated as drug concentration solicited to reduce the viability of cell up to 50%

#### 4.3 Determination of $\text{EC}_{50}$ value of HDF

Antivirucidal nature of HDF was determined by observing plaque count in supernatants of infected Vero cells by adopting plaque number reduction method.<sup>18)</sup> Infected Vero cells after 60 minute of absorption at temperature of  $37^{\circ}\text{C}$  temperature, all the petri plates were douched with 5% ethanol and growing medium was supersede with RPMI-1640 medium containing carboxymethyl cellulose 0.5% Fetal Bovine Serum and (30  $\mu\text{g mL}^{-1}$ ) HDF was added with Vero cells. After 24 hours incubation, carboxymethyl cellulose was discarded and monolayers were treated with 10% formaldehyde dissolve in Phosphate Buffer Saline and cells staining were performed by 1% methanolic solution of crystal violet indicator. The concentration of HDF required reducing virus yield up to 50% ( $\text{EC}_{50}$ ) was reckoned by regression analysis of the dose-response curves. Selectivity index value<sup>19)</sup> of HDF was calculated by using the formula Selectivity Index Value (SI Value) =  $\text{CC}_{50}$  Value/ $\text{EC}_{50}$  Value

#### 4.4 Evaluation of Synergistic activity of HDF with Acyclovir

In synergistic study of HDF and Acyclovir minimum inhibitory concentration (EC value) of Acyclovir (0.5  $\mu\text{g mL}^{-1}$ ) and 2.5  $\mu\text{g mL}^{-1}$  of HDF was used.

To observe the synergistic activity of HDF and ACV at post infection (p.i.), infected Vero cells were incubated with HDF and Acyclovir for 60 minute at  $37^{\circ}\text{C}$  temperature. After centrifugation supernatants of infected Vero cells were separate out and used to observe the plaques while non treated infected Vero cells (CTR) were used as reference.<sup>20)</sup>

#### 4.5 Affectivity of HDF during various stages of HSV-1 infection

To determine the effect of HDF at different stages of infection of HSV-1, half minimum effective concentration of HDF (25.98  $\mu\text{g mL}^{-1}$ ) was added with Vero cells (i) before infection induced by HSV-1 (ii) during the adsorption period for 1 hour (iii) Immediately after the adsorption of HSV-1. In all steps Vero cells were incubated in RPMI supplemented 2% concentrated

solution of PBS for 24 hour by maintaining temperature  $37^{\circ}\text{C}$  and supernatants of infected Vero cells were used to identify the affectivity of HDF as antiviral agent.<sup>21)</sup>

#### 4.6 Assessment of anti-viral nature of HDF at early stages of post infection

Vero cells matured in 24-well plates induced an infection by HSV-1 (0.1 m.o.i) and incubated for 2 hour at  $4^{\circ}\text{C}$ . Further infected Vero cells were washed by phosphate buffered saline to remove viral debaris. 25.98  $\mu\text{g mL}^{-1}$  concentration of HDF was supplemented with infected Vero cells at various times of post infection for 0–1 hour, 1–3 hour, 4–6 hour and 6–8 hour. After this specified time lap the supernatants of untreated infected Vero cells and HDF treated infected cells were recovered and to assess anti herpetic behavior of HDF by plaque number reduction assay.

#### 4.7 Assessment of antiviral nature of HDF at late stages of post-infection


To evaluate the conseiences HDF on viral replication at post infection, firstly Vero-cells were incubated with HSV-1 at  $37^{\circ}\text{C}$  temperature for 5-6 minute to allow virus to penetrate Vero cell membrane and allow replication of virus for 24 hours. Infected Vero cells were treated with 25.98  $\mu\text{g mL}^{-1}$  concentration of HDF for 0-24 hour, 15 minute-24 hour, 45 minute-24 hour, 2-24 hour, 3-24 hour and 4-24 hour while untreated Infected Vero cells (CTR) were used as control.<sup>22)</sup>

#### 4.8 Western Blotting Analysis

To explore the attribute of HDF to intrupt protein expression of HSV-1, western blotting technique was employed.<sup>24,25)</sup> Vero cells, previously washed with phosphate buffer saline centrifuged for 15 minutes and pellets were adjourned in cold lymph buffer solution containing 10mM phenyl methacophenol fluoride phosphatase as inhibitor mixture along with 2% triton solution, incubated for 30 minutes at  $0^{\circ}\text{C}$  temperature and supernatants were arrassed to negotiate protein expression. The protein contents were aloofed by gel electrophoresis technique in 10% S.D.S. polyacrylamide and electroblotted on to nitrocellulose membrane. After blocking the membrane using 10-12% dry fat free milk suspended in tris-buffer solution and 0.5 % solution of twin-20 for 60 minute at  $25^{\circ}\text{C}$  temperature, incubated with 1gm/mL anti ICP-5, anti ICP-6, anti ICP-8, anti-gb anti  $\beta$ -actin antibodies (control for protein loading). Later on washing membranes were incubated for 60-70 minute with secondary antibodies conjugated with horseradish peroxidase band of protein were appearent by ECL plus detection system by following instruction manualed by manufacturer and densitometry data for the blots were analysed with help of imagJ software.

## 5. Result and Discussion

The cytotoxicity of HDF was represented by dose response curve between cell viability percentage versus varied concentration of HDF. (Figure : 6)




**Fig.6:**Determination of CC<sub>50</sub> value of HDF.

During experiment, no substantial changes in cell count were observed up to 200 µg/mL<sup>-1</sup> concentration of HDF which evidenced by trypan blue staining. HDF start to show a toxic effect on the viability of Vero cells after 300 µg/mL<sup>-1</sup> till the highest dose used in experiment.


Curve in Figure-7 shows the half maximum effective concentration value (EC<sub>50</sub> value) of HDF against HSV-1 strain. Such Graph indicates at 3.69 µg/mL<sup>-1</sup> concentration of HDF, the percentage of plaque formation start to reduce.

For the reduction of 50% of plaques, the required concentration of compound was 25.98 µg/mL<sup>-1</sup>.

The efficacy of HDF as an antiviral compound can be expressed by the selectivity index value (S.I. value). The S.I. value of HDF was 11.54 which was greater than 10 which ensures that HDF can act as a potential antiviral agent.




**Fig. 7:**Determination of EC<sub>50</sub> value of HDF



**Fig.8:**Effect of HDF on viral plaque of HSV-1 (a) Viral plaques in absence of HDF (b) Viral plaques in presence of HDF


It was observed by synergistic action of HDF with Acyclovir that 0.5 µg/mL<sup>-1</sup> quantity of Acyclovir was enough to inhibit HSV-1 replications nearly 48 percent as compared with CTR and almost alike inhibition was

monitored in Vero cells treated with 2.5 µg/mL<sup>-1</sup> concentration of HDF. (Figure-9)




**Fig.9:**Synergistic antiviral activity of HDF and Standard drug Acyclovir

On observing the effectivity of HDF at different stages of HSV-1 induced infection, that HDF not show any inhibitory action on replication of virus during adsorption period meanwhile after the penetration of virus it show successive increment of extent of inhibition with time. Such result suggested that HDF can act as antiherpetic agent for HSV-1 induced infection at post infection. The inhibitory action of HDF was confirmed when infected Vero cells were treated with compound at different time of post infection (for 0-24 hour, 15 minute-24 hour, 45 minute-24 hour, 2-24 hour, 3-24 hour and 4-24 hour) (Figure 10 & 11)



**Fig.10:**Antiviral effect of HDF on Viral replication during adsorption period and at initial and different time of post infection (p.i.) at 0-1 hour, 1-3 hour, 4-6 hour and 6-8 hour



**Fig.11:**Antiviral effect of HDF on adsorption period and different time of post infection

The result demonstrated by western blot analysis (Figure-12) indicates that  $\beta$ -(ICP-6) and  $\gamma$ -(ICP-5 and gB) protein expression of HSV-1 was reduced



remarkably in Vero cells which were treated with HDF but  $\alpha$ -(ICP-8) show partial reduction.




Fig.12:western blot analysis of HDF

## 6. Conclusion

Our finding suggested that compound ((E)-N-(3,4-dimethylisoxazol-5-yl)-4-((4-hydroxy-2-oxo-2H-chromen-3-yl)diazetyl)benzenesulfonamide act as antiviral agent against *HSV-1* evidenced by its good selectivity index (SI value = 11.54) and lesser toxicity. This compound is less able to inhibit the viral replication during the adsorption period but it can inhibit such replication at post infection. In western blotting analysis the reduction of specific gene products  $\beta$ -(ICP-6) and  $\gamma$ -(ICP-5 and gB) groups strongly recommended that HDF can able to inhibit *HSV-1* replication in host cell by blocking virus DNA synthesis at late stage of infection.

## Acknowledgements

All the authors are highly thankful to Dr. R. G. Sharma, former Principal & Head, Department of Chemistry, N.A.S. College, Meerut (U.P.) India for the valuable suggestion during whole research work.

## Nomenclature

ACV	acyclovir
CC <sub>50</sub>	50% cytotoxic concentration
HSV-1	herpes simplex virus type-1
EC <sub>50</sub>	half minimum effective concentration
MOI	multiplicity of infection
Pfu	plaque forming unit
SI	selectivity index
CTR	untreated infected Vero cells

## References

- 1) Aldrin Man, Mark Slevin, Eugen Petcu & Cornel Fraefel, "The Cyclin-Dependent Kinase 5 Inhibitor Peptide Inhibits Herpes Simplex Virus Type 1

- Replication", *Scientific Reports*, **9** (1260) 1-8 (2019). doi.org/10.1038/s41598-018-37989-3
- 2) Pamela Chayavichitsilp, Joseph V Buckwalter, Andrew C Krakowski, "Herpes simplex", *Pediatrics in Review*, **30** (4)119-129(2009). doi: 10.1542/pir.30-4-119
- 3) Erik De Clercq, "Selective Antiherpes virus agents", *Antiviral Chem. & chemotherapy*, **22** 93-101 (2013). doi.org/10.3851/IMP2533
- 4) Scott H James and Mark N Prichard, "Current and future therapies for herpes simplex Virus infections: Mechanism of action and drug resistance", *Curr. Opin. Virology*, **8** 54-64 (2014). doi: 10.1016/J.coviro.2014.06.003
- 5) Angela Stefanachi, Francesco Leonetti, Leonardo Pisani, Marco Catto & Angelo Carotti, "Coumarin: A Natural, Privileged and Versatile Scaffold for Bioactive Compounds", *Molecules*, **23** (2) 250 (2018). doi: 10.3390/molecules23020250
- 6) F. Borgess, N Roleria, I Mulhazes Santana, E. Uriate, "Simple coumarins and analogues in medicinal chemistry, Occurance, synthesis and Biological activity", *Curr. Med. Chem.* **12** (8)887-916 (2005) doi: 10.2174/0929867053507315.
- 7) B. Serge Kirkiacharian, Erik De Clercq, Raffi Kurkjian, "New synthesis and anti-HIV and antiviral properties of 3-arylsulfonyl derivatives of 4-hydroxycoumarin and 4-hydroxyquinolone", *Pharm Chem J.*, **42**, 265 (2008). doi.org/10.1007/s11094-008-0103-0
- 8) SB Diwakar, B Govindh, Y Nagendra Sastry, "Synthesis, X-ray characterization and biological evaluation of some new 2-(4-methy-2-oxo-2H-chromen-7-yl)oxy)acetamide derivatives", *Med. Chem. Res.* **24**(4) 1546-1557 (2015). doi:10.1007/s00044-014-1230-7
- 9) O. Nagaraja, D. Bodke Yadav, PushPavathi Itte "Synthesis, characterization and biological investigations of potentially bioactive heterocyclic compounds containing 4-hydroxy coumarin", *Heliyon*, **6** (6)(2020) 1-11. doi: 10.1016/J.heliyon.2020.e04245
- 10) J. Sahoo, Makeap Kumar & Sudhir Kumar "Synthesis, spectral characterization of some new 3-heteroaryl azo-4-hydroxy coumarin derivatives and their antimicrobial evaluation", *J.of Taibah University for Science* **9** 187-195 (2015). doi.org/10.1016/J.jtusci.2014.08.001
- 11) Ahmed Sabt, O.M. Abdelhofer, R.S. El-Hagger "Novel coumarin-6-sulfonamides as apoptotic antiproliferative agents: synthesis, in vitro biological evaluation, and QSAR studies", *J. of Enzyme Inhibition & Medicinal Chem* **33**(1)1095-1107 (2018). doi.org/10.1080/14756366.2018.1477137
- 12) Mohd. Zaheen Hassan, Hasnati Osman and Mohd. Jawed Ahsan, "Therapeutic potential of coumarins

- as antiviral agents, *Eur. J. Med. Chem.* **123**(10) 236-255 (2016). doi: 10.1016/J.ejmech.2016.07.056
- 13) M.R. Yazdanbakhsh, A. Ghanadzadeh, E. Moradi, "Synthesis of some new azo dyes derived from 4-hydroxy coumarin and spectrometric determination of their acidic dissociation constants", *J molecular liquids* **136** 165-168 (2007). doi: 10.1016/J.molliq.2007.03.005
- 14) Davorka Završnik, Samija Muratović, Damjan Makuc, Janez Plavec, "bis-(4-Hydroxycoumarin) and Benzopyrano- Coumarin Derivatives: Synthesis, <sup>1</sup>H/ <sup>13</sup>C-NMR Conformational and X-ray Crystal Structure Studies and In Vitro Antiviral Activity Evaluations", *Molecules* **16**, 6023-6040 (2011). doi: 10.3390/molecules16076023
- 15) Neda mostajeran, Farzaneh Amin Arshad, Hamid Aliyan and Ahmad Reza Mass, "Solvent free synthesis and antibacterial evaluation of novel coumarin sulfonamides", *Pharmaceutical Chemistry Journal*, **52** (1) 1-6 (2018). doi: 10.1007/s11094-018-1756-y
- 16) Ito Kazuhide, Micro and Macro scale measurement of fungal growth under various temperature and humidity condition, *Evergreen*, **1** (1) 32-39 (2014)
- 17) M Wahid, M. Noor, H. Hara, Recombinant Moringa Oleifera Lectin produced in Pichia Pastoris is a Potential Natural Coagulant *Evergreen*, **3** 11-16 (2016)
- 18) L Boff, IT Silva, DF Argenta, LM Farias, "Strychnos pseudoquina A. St. Hil.: a Brazilian medicinal plant with promising in vitro antiherpes activity", *J Appl Microbiol*, **121** (6) 1519-1529 (2016). doi.org/10.1111/Jam.13279.
- 19) L Boff, J Munkert, FM Ottoni, NFZ Schneider, GS Ramos, "Potential anti-herpes and cytotoxic action of novel semisynthetic digitoxigenin-derivatives" *Eur J Med Chem* **167** 546-561 (2019).doi.org/10.1016/J.eJmech.2019.01.076
- 20) John F. Marcelletti, "Synergistic inhibition of herpesvirus replication by docosanol and antiviral nucleoside analogs", *Antiviral Res.* **56** (2) 153-166 (2002). doi.org/10.1016/S0166-3542(02)00105-5
- 21) Livia Civitelli, Simona Panella, Maria Elena Marcocci, Alberto De Petris "In vitro inhibition of herpes simplex virus type 1 replication by Mentha suaveolens essential oil and its main component piperitenone oxide", *Phytomedicine* **21** 857-865 (2014). doi: 10.1016/J.phymed.2014.01.013
- 22) ALF Larsson, "Mode of action, toxicity, pharmacokinetics and efficacy of some new antiherpesvirus Guanosine analogs related to Buciclovir", *Antimicrobial Agent & Chemotherapy*, **30** (4) 598-605 (1986)
- 23) Arhiro Kano, Moes Kibunja Kamita, Tauma Iwasaki, Mitsuru Shindo, "Bongkreic Acid Induces Selective cytotoxicity in tumor cells revealed by CCK-8", *Evergreen*, **2/3** 23-27 (2017)
- 24) M M Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding". *Anal Biochem.* **72** 248-254 (1986). doi.org/10.1016/0003-2697(76)90527-3
- 25) Laurita Boff, Naira Fernanda, Zanchett Schneider, Jennifer Munkert, Flaviano Melo Ottoni, "Elucidation of the mechanism of anti-herpes action of two novel semisynthetic cardenolide derivatives", *Archives of Virology* **165** 1385-1396 (2020).doi.org/10.1007/s00705-020-04562-1