Reliability and Maintainability of a Multicomponent Series-Parallel System with Simultaneous Failure under Preemptive Repeat Repair Discipline

児玉, 正憲

<https://doi.org/10.15017/4491667>

出版情報 : 経済學研究. 52 (6), pp.9-23, 1987-08-10. 九州大学経済学会
バージョン :
権利関係 :

RELIABILITY AND MAINTAINABILITY OF A MULTICOMPONENT SERIES-PARALLEL SYSTEM WITH SIMULTANEOUS FAILURE UNDER PREEMPTIVE REPEAT REPAIR DISCIPLINE

Masanori KODAMA

1. INTRODUCTION

In this paper we consider a multicomponent system composed of two subsystem S_0 and S_1 in series. Subsystem S_0 is K -out-of- N : G system (more than $N-K+1$ must fail for S_0 to fail), while subsystem S_1 consists of M different units connected in series (the failure of any one causes S_1 to fail).

The life-times of the active units depend on each other in having simultaneous failure of all the operating units and repair times are distributed quite generally. A failed unit is repaired at a single service channel. Kulshrestha [1,2] has studied the model for $K=1$ under the assumption that the repair of any unit is possible only when the system stops operating and the life times of the active units are mutually independent. Nakamichi et al. [3] and Kodama [4] have studied the model for $K=1$ with exponential failure distribution under the different repair policies. Kodama et al. [5] have investigated the system presented here under the assumption that the repair of any unit is possible only when the system stops operating. Assuming that the failed unit is repaired as soon as the unit fails and the system has priority in repair, Laplace transform of pointwise availability and reliability, mean time to the first system failure and steady state availability are derived. In practice, there are a lot of cases which the above conditions are satisfied (for example, an electronics and communication system, a space shuttle, . . .) and the above mentioned results are used for the reliability design of complex system, so it is important to study the above system. And also the theoretical interest in reliability analysis of the above system exists.

2. DEFINITION OF MODELS AND NOTATIONS

Independent Poisson processes $Z_1(t; \lambda_0), \dots, Z_N(t; \lambda_0), Z_1(t; \lambda_1), \dots, Z_M(t; \lambda_M)$ and $Z(t;$

λ^*) govern the occurrence of shock (By $Z(t; \lambda)$, we mean a Poisson process with a parameter λ). Events in the process $Z_i(t; \lambda_0)$ are shocks to unit i only in $S_0 (1 \leq i \leq N)$, and events in the process $Z_j(t; \lambda_j)$ are shocks to unit j only in $S_1 (1 \leq j \leq M)$, and events in the process $Z(t; \lambda^*)$ are shocks to all operating units.

Each unit results in failure after receiving only one “shock”. The system breaks down if more than $(N - K + 1)$ units in S_0 are simultaneously in a failed state or if any failure in S_1 occurs or if simultaneous failure occurs. A failed unit is repaired at a single service channel, and the repaired unit is assumed to behave like new after repair. Repair times are distributed quite generally. When the system is in a failed state, the residual good units do not fail.

While the system is operating, whenever any unit in S_0 fails, it is sent to the repair station at once where it is repaired on a first-come-first-served basis.

A repaired unit is put in operation immediately. When the system stops operating by stop of S_0 , as soon as the unit in S_0 being served is repaired, it is reentered and then the system operates again. When the system stops operating by the stop of S_1 , as soon as the failed unit of S_1 is repaired the system operates again. If a unit in S_1 fails when a unit in S_0 is being repaired, it is repaired without delay by interrupting the repair of unit from S_0 ; the preempted unit from S_0 repeat service from the beginning on its reentry (preemptive repeat repair discipline). Through this model, m units in $S_0 (K \leq m \leq N)$ and all the units in S_1 are repaired when the system stops operating by the simultaneous failure. If the simultaneous failure occurs when a unit in S_0 is being repaired, all unit in S_1 and m units in $S_0 (K \leq m \leq N - 1)$ are repaired without delay by interrupting the repair of the unit from S_0 , if any; the preempted unit from S_0 obeys the ‘repeat rule’ so that upon reentry its repair is to be started from the beginning. Similarly, we can discuss for the case of preemptive resumption priority (see reference [6]).

The system is always in one of the following states: E_N -all the unit in S_0 and S_1 are operative; E_i - i units in S_0 and all the units in S_1 are operative, and a failed unit is undergoing repair, $K \leq i \leq N - 1$; F_{K-1} - $(N - K + 1)$ units in S_0 have failed and the system is undergoing repair; F_{ij} - i units in S_0 are operative and the system stops operating by the failure of the j th unit in S_1 and the system is undergoing repair, $K \leq i \leq N, 1 \leq j \leq M$; F_0 -all the units in S_0 and S_1 have failed and the system is undergoing repair. We note that E . means that the system is operating and F . means that the system is failed. Initially, all the units are operating.

Through this paper we use the following notations;

$$\lambda = \sum_{j=1}^M \lambda_j;$$

$f_0(t)$ is the density function of repair time of a unit in S_0 ,

$$f_0(t) = \mu_0(t) \exp \left[- \int_0^t \mu_0(x) dx \right];$$

$f_j(t)$ is the density function of repair time of j th unit in S_1 ,

$$f_j(t) = \mu_j(t) \exp \left[- \int_0^t \mu_j(x) dx \right], \quad 1 \leq j \leq M, \quad f(t) = \sum_{j=1}^M \lambda_j f_j(t) / \lambda;$$

$g_{i+M}^*(t)$ is the density function of repair time of i units in S_0 and M units in S_1 when the system stops operating by the simultaneous failure,

$$g_{i+M}^*(t) = \mu_{i+M}^*(t) \exp \left[- \int_0^t \mu_{i+M}^*(x) dx \right], \quad K \leq i \leq N;$$

K_0 , K_j and K_{i+M}^* are the first moment (mean repair time) of $f_0(t)$, $f_j(t)$ and $g_{i+M}^*(t)$ respectively;
 δ is the Kronecker delta;

$\binom{m}{i}$ is the binomial coefficient that is assumed to be zero if $m < i$; $P_i(t) = P$ {the system is in state E_i at time t }, $K \leq i \leq N$;

$P_i(t, x) dx = P$ {the system is in state E_i at time t and elapsed repair time lies between x and $x + dx$ },

$$P_i(t) = \int_0^t P_i(t, x) dx, \quad K \leq i \leq N-1;$$

$P_{K-1}(t) = P$ {the system is in state F_{K-1} at time t }

$P_{K-1}(t, x) dt = P$ {the system is in state F_{K-1} at time t and elapsed repair time lies between x and $x + dx$ },

$$P_{K-1}(t) = \int_0^t P_{K-1}(t, x) dx;$$

$P_{ij}(t) = P$ {the system is in state F_{ij} at time t }

$P_{ij}(t, x) dx = P$ {the system is in state F_{ij} at time t and elapsed repair time lies between x and $x + dx$ }

$$P_{ij}(t) = \int_0^t P_{ij}(t, x) dx, \quad K \leq i \leq N, \quad 1 \leq j \leq N;$$

$Q_0(t) = P$ {the system is in state F_0 at time t }

$Q_0(t, x) dt = P$ {the system is in state F_0 at time t and elapsed repair time lies between x and $x + dx$ }

$$Q_0(t) = \int_0^t Q_0(t, x) dx$$

$P_A(t) = P$ {the system is functioning at time t } = $\sum_{i=K}^N P_i(t)$

(pointwise availability function)

$R(t) = P$ {the system does not break down during $[0, t]$ }

(reliability function)

MTSF is the mean time to the first system failure $\bar{M}(s)$ is the Laplace transform of $M(t)$, $\bar{M}(s)$

$$= \int_0^\infty e^{-st} M(t) dt;$$

$$P_i(x) = \lim_{t \rightarrow \infty} P_i(t, x), \quad K-1 \leq i \leq N-1;$$

$$P_i = \int_0^\infty P_i(x) dx, \quad K-1 \leq i \leq N-1;$$

$$P_N = \lim_{t \rightarrow \infty} P_N(t);$$

$$P_{ij}(x) = \lim_{t \rightarrow \infty} P_{ij}(t, x), \quad K \leq i \leq N, \quad 1 \leq j \leq M;$$

$$P_{ji} = \int_0^\infty P_{ij}(x) dx, \quad K \leq i \leq N, \quad 1 \leq j \leq M;$$

$$Q_0(x) = \lim_{t \rightarrow \infty} Q_0(t, x);$$

$$Q_0 = \int_0^\infty Q_0(x) dx;$$

$$P_A = \sum_{i=K}^N P_i \text{ (steady state availability of the system);}$$

$$\alpha(s, i, \lambda^*) = s + i\lambda_0 + \lambda(1 - \bar{f}(s)) + \lambda^*, \quad K \leq i \leq N;$$

$$V_i^{(1)}(s) = \begin{cases} \prod_{r=K+1}^i \{[\bar{f}_0(s + r\lambda_0 + \lambda + \lambda^*) - \lambda \bar{f}(s)(1 - \bar{f}_0(s + r\lambda_0 + \lambda + \lambda^*))]/ \\ (s + r\lambda_0 + \lambda + \lambda^*)\} / \bar{f}_0(s + r\lambda_0 + \lambda + \lambda^*)\}, & K+1 \leq i \leq N-1, \\ 1, & i = K \end{cases}$$

$$\sum_{n=i}^j h(n) = 0, \quad i > j \text{ (} h(n) \text{ is a function defined for } n = i, \dots, j \text{)}$$

$$C_{i,j}^{(1)}(s) = \sum_{n=i}^j \binom{N}{n} \alpha(s, n, \lambda^*) V_{n-1}^{(1)}(s) / \alpha(s, N, \lambda^*);$$

$$E_{i,j}^{(1)}(s) = \sum_{n=i}^j \binom{N}{n} (N-n)\lambda_0 V_{n-1}^{(1)}(s) / \alpha(s, N, \lambda^*);$$

$$E_{i,j}^{(1),m} = \sum_{n=i}^j \binom{m}{n} V_{n-1}^{(1)}(s);$$

$$a_i^{(1)}(s) = \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) C_{i+1,N}^{(1)}(s) / [\bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*) V_i^{(1)}(s) C_{K+1,N}^{(1)}(s)], \\ K+1 \leq i \leq N-1,$$

$$b_i^{(1)}(s) = [C_{K+1,N}^{(1)}(s) E_{K+1,i}^{(1),m}(s) - C_{K-1,i}^{(1)}(s) E_{K+1,N-1}^{(1),m}(s)] / \\ [\bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*) V_i^{(1)}(s) C_{K+1,N}^{(1)}(s)], \quad K+1 \leq i \leq N-1$$

$$b_i^{(1)*}(s) = [C_{K+1,N}^{(1)}(s) E_{K+1,i}^{(1)}(s) - C_{K+1,i}^{(1)}(s) E_{K+1,N-1}^{(1)}(s)] / \\ [\bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*) V_i^{(1)}(s) C_{K+1,N}^{(1)}(s)], \quad K+1 \leq i \leq N-1$$

$$k_1^{(1)}(s) = \{1 - \lambda^* \bar{g}_{m+M}^*(s) \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)}(s) / \\ \alpha(s, N, \lambda^*) - \lambda^* \bar{g}_{m+M}^*(s) \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} (1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) \\ \cdot b_n^{(1)}(s) / (s + n\lambda_0 + \lambda + \lambda^*)\}^{-1} \{ \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) a_{N-1}^{(1)}(s) / \alpha(s, N, \lambda^*) \\ + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} (1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) a_n^{(1)}(s) / (s + n\lambda_0 + \lambda + \lambda^*) \}$$

$$k_0^{(1)}(s) = \{1 - \lambda^* \bar{g}_{m+M}^*(s) \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)}(s) / \\ \alpha(s, N, \lambda^*) - \lambda^* \bar{g}_{m+M}^*(s) \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} (1 - \bar{f}_0(s + n\lambda_0 + \lambda^*)) \\ \cdot b_n^{(1)}(s) / (s + n\lambda_0 + \lambda + \lambda^*)\}^{-1} \{ [1 + \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)*}(s)] / \\ \alpha(s, N, \lambda^*) + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} (1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) b_n^{(1)*}(s) / \\ (s + n\lambda_0 + \lambda + \lambda^*) \}$$

$$a_i^{(1)'}(s) = a_i^{(1)}(s) + \lambda^* \bar{g}_{m+M}^*(s) k_1^{(1)}(s) b_i^{(1)}(s), \quad K+1 \leq i \leq N-1;$$

$$b_i^{(1)'}(s) = \lambda^* \bar{g}_{m+M}^*(s) k_0^{(1)}(s) b_i^{(1)}(s) + b_i^{(1)*}(s), \quad K+1 \leq i \leq N-1;$$

3. SOLUTIONS OF THE PROBLEM

The analyses are classified according to the nature of this system and the relations among m , K , and N and in this way three cases may occur: i) $1 \leq K \leq m \leq N-1$, ii) $1 \leq K \leq N-1$, $m = N$, iii) $2 \leq K = N$, $m = N$. The unified analysis of three cases is more difficult than a separate analysis. Therefore, we make the separate analysis of three cases. The analysis crucially depends on the method of supplementary variables, and the supplementary variable x denotes the elapsed time that a unit has been undergoing repair. Viewing the nature of this model, we obtain the following set of differential-difference equations:

(i) $1 \leq K \leq m \leq N-1$,

$$[d/dt + N\lambda_0 + \lambda + \lambda^*]P_N(t) = \int_0^t \mu_0(x)P_{N-1}(t, x)dx + \sum_{j=1}^M \int_0^t \mu_j(x)P_{Nj}(t, x)dx, \quad (3.1)$$

$$[\partial/\partial t + \partial/\partial x + i\lambda_0 + \lambda + \lambda^* + \mu_0(x)]P_i(t, x) = (1 - \delta_{i, N-1})(i+1)\lambda_0 P_{i+1}(t, x), \quad K \leq i \leq N-1, \quad (3.2)$$

$$[\partial/\partial t + \partial/\partial x + \mu_0(x)]P_{K-1}(t, x) = K\lambda_0 P_K(t, x), \quad (3.3)$$

$$[\partial/\partial t + \partial/\partial x + \mu_j(x)]P_{ij}(t, x) = 0, \quad K \leq i \leq N, \quad 1 \leq j \leq M, \quad (3.4)$$

$$[\partial/\partial t + \partial/\partial x + \mu_{m+M}^*(x)]Q_0(t, x) = 0, \quad (3.5)$$

$$P_i(t, 0) = \delta_{i, N-1}N\lambda_0 P_N(t) + \int_0^t \mu_0(x)P_{i-1}(t, x)dx + \sum_{j=1}^M \int_0^t \mu_j(x)P_{ij}(t, x)dx + \delta_{i, m} \int_0^t \mu_{m+M}^*(x)Q_0(t, x)dx, \quad K \leq i \leq N-1, \quad (3.6)$$

$$P_{K-1}(t, 0) = 0, \quad (3.7)$$

$$P_{ij}(t, 0) = \lambda_j P_i(t), \quad K \leq i \leq N, \quad 1 \leq j \leq M, \quad (3.8)$$

$$Q_0(t, 0) = \lambda^* \sum_{i=K}^N P_i(t) = \lambda^* P_N(t) + \lambda^* \sum_{i=K}^{N-1} \int_0^t P_i(t, x)dx, \quad (3.9)$$

$$P_N(0) = 1. \quad (3.10)$$

In order to solve these equations we introduce the discrete transform (see reference [10], p. 26-30).

$$U_i(t, x) = \sum_{n=i}^{N-1} \binom{n}{i} P_n(t, x), \quad K \leq i \leq N-1, \quad (3.11)$$

$$P_i(t, x) = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} U_n(t, x), \quad K \leq i \leq N-1 \quad (3.12)$$

Taking the Laplace transform of (3.1)-(3.9), (3.11) and (3.12) under the initial condition and using the results of Appendix 1, we obtain.

$$\bar{P}_{ij}(s) = \lambda_j \bar{P}_i(s)(1 - \bar{f}_j(s))/s, \quad K \leq i \leq N, \quad 1 \leq j \leq M, \quad (3.13)$$

$$\bar{Q}_0(s) = \lambda^* \sum_{i=K}^N \bar{P}_i(s)(1 - \bar{g}_{m+M}^*(s))/s, \quad (3.14)$$

$$\bar{P}_i(s) = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} \bar{U}_n(s, 0)[1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)]/(s + n\lambda_0 + \lambda + \lambda^*), \quad K \leq i \leq N-1, \quad (3.15)$$

$$\bar{P}_N(s) = [\bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*)\bar{U}_{N-1}(s, 0) + 1]/\alpha(s, N, \lambda^*), \quad (3.16)$$

$$\begin{aligned} \bar{P}_{K-1}(s) = & K\lambda_0 \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} [\bar{U}_n(s, 0) / (n\lambda_0 + \lambda + \lambda^*)] \\ & \cdot [(1 - \bar{f}_0(s)) / s - (1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*)] \end{aligned} \quad (3.17)$$

$$\bar{U}_i(s, 0) = a_i^{(1)'}(s) \bar{U}_K(s, 0) + b_i^{(1)'}(s), \quad K+1 \leq i \leq N-1, \quad (3.18)$$

$$\begin{aligned} \bar{U}_K(s, 0) = & \left\{ \binom{N}{K} [(N-K)\lambda_0 - \alpha(s, K, \lambda^*) \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)'}(s)] / \alpha(s, N, \lambda^*) \right. \\ & + \sum_{n=K}^{N-1} (-1)^{n-K} \left[\binom{n}{K-1} \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*) + \binom{n}{K} K\lambda_0 (\bar{f}_0(s) - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / \right. \\ & (n\lambda_0 + \lambda + \lambda^*) \left. \right] b_n^{(1)'}(s) + \binom{m}{K} \lambda^* \bar{g}_{m+M}^*(s) k_0^{(1)}(s) \left. \right\} \\ & \cdot \left\{ 1 - \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) - \lambda \bar{f}(s) [1 - \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*)] / (s + K\lambda_0 + \lambda + \lambda^*) \right. \\ & + \binom{N}{K} \alpha(s, K, \lambda^*) \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) a_{N-1}^{(1)'}(s) / \\ & \alpha(s, N, \lambda^*) - \sum_{n=K}^{N-1} (-1)^{n-K} \left[\binom{n}{K-1} \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*) + \binom{n}{K} K\lambda_0 (\bar{f}_0(s) \right. \\ & - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / \\ & (n\lambda_0 + \lambda + \lambda^*) \left. \right] a_n^{(1)'}(s) - \binom{m}{K} K\lambda^* \bar{g}_{m+M}^*(s) k_1^{(1)}(s) \left. \right\}^{-1}, \end{aligned} \quad (3.19)$$

$$\sum_{j=K}^N \bar{P}_j(s) = k_1^{(1)}(s) \bar{U}_K(s, 0) + k_0^{(1)}(s), \quad (3.20)$$

$P_{ij}(t)$, $Q_0(t)$, $P_{K-1}(t)$ and $P_i(t)$ are the probability that the system are in state F_{ij} , F_0 , F_{K-1} and E_i at time t respectively. Hence the Laplace transform of the system availability $\bar{P}_A(s)$ is given by (3.20).

The Laplace transform of system reliability $\bar{R}(s)$ can be obtain from $\bar{P}_A(s)$ by modifying the system so that a down state is an absorbing state, it can never go up. So we substitute $\bar{f}(s) = 0$, $(\bar{f}_0(s) - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) = 0$, $K \leq n \leq N-1$ and $\bar{g}_{m+M}^*(s) = 0$ in $\bar{P}_A(s)$. Moreover, if we set $s = 0$ in $\bar{R}(s)$, we obtain MTSF.

In order to derive the steady state availability we set up the following set of differential-difference equations:

$$[N\lambda_0 + \lambda + \lambda^*]P_N = \int_0^\infty \mu_0(x)P_{N-1}(x)dx + \sum_{j=1}^M \int_0^\infty \mu_j(x)P_{Nj}(x)dx, \quad (3.21)$$

$$[d/dx + i\lambda_0 + \lambda + \lambda^* + \mu_0(x)]P_i(x) = (1 - \delta_{i,N-1})(i+1)\lambda_0 P_{i+1}(x), \quad K \leq i \leq N-1, \quad (3.22)$$

$$[d/dx + \mu_0(x)]P_{K-1}(x) = K\lambda_0 P_K(x), \quad (3.23)$$

$$[d/dx + \mu_j(x)]P_{ij}(x) = 0, \quad K \leq i \leq N, \quad 1 \leq j \leq M, \quad (3.24)$$

$$[d/dx + \mu_{m+M}^*(x)]Q_0(x) = 0, \quad (3.25)$$

With the following boundary conditions and normalizing condition:

$$\begin{aligned} P_i(0) = & \delta_{i,N-1} N\lambda_0 P_N + \int_0^\infty \mu_0(x)P_{i-1}(x)dx + \sum_{j=1}^M \int_0^\infty \mu_j(x)P_{ij}(x)dx \\ & + \delta_{i,m} \int_0^\infty \mu_{m+M}^*(x)Q_0(x)dx, \quad K \leq i \leq N-1, \end{aligned} \quad (3.26)$$

$$P_{K-1}(0) = 0, \quad (3.27)$$

$$P_{Nj}(0) = \lambda_j P_N, \quad 1 \leq j \leq M, \quad (3.28)$$

$$P_{ij}(0) = \lambda_j \int_0^\infty P_i(x) dx = \lambda_j P_i, \quad K \leq i \leq N-1, \quad 1 \leq j \leq M, \quad (3.29)$$

$$Q_0(0) = \lambda^* P_N + \lambda^* \sum_{i=K}^{N-1} \int_0^\infty P_i(x) dx = \lambda^* \sum_{i=K}^N P_i, \quad (3.30)$$

$$P_N + \sum_{i=K}^{N-1} \int_0^\infty P_i(x) dx + \int_0^\infty P_{K-1}(x) dx + \sum_{j=1}^M \int_0^\infty P_{Nj}(x) dx + \sum_{i=K}^{N-1} \sum_{j=1}^M \int_0^\infty P_{ij}(x) dx + \int_0^\infty Q_0(x) dx = 1. \quad (3.31)$$

Solving the above equations, we obtain (see Appendix 2)

$$P_{ij} = \lambda_j K_j P_i, \quad (3.32)$$

$$Q_0 = \lambda^* K^*_{m+M} \sum_{i=K}^N P_i, \quad (3.33)$$

$$P_i = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} U_n(0) (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*), \quad K \leq i \leq N-1, \quad (3.34)$$

$$P_N = \bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) U_{N-1}(0) / (N\lambda_0 + \lambda^*), \quad (3.35)$$

$$P_{K-1} = \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} K \lambda_0 U_n(0) [K_0 - (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*)] / (n\lambda_0 + \lambda + \lambda^*), \quad (3.36)$$

$$U_i(0) = a_i^{(1)'}(0) U_K(0), \quad K+1 \leq i \leq N-1, \quad (3.37)$$

$$U_K(0) = \left\{ k_1^{(1)}(0) \left[1 + \lambda^* K^*_{m+M} + \sum_{j=1}^M \lambda_j K_j \right] + \sum_{n=K}^{N-1} (-1)^{n+K} \binom{n}{K} [K \lambda_0 / (n\lambda_0 + \lambda + \lambda^*)] \cdot [K_0 - (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*)] a_n^{(1)'}(0) \right\}^{-1}, \quad (3.38)$$

$$\sum_{n=K}^N P_n = k_1^{(1)}(0) U_K(0). \quad (3.39)$$

P_{ij} , Q_0 , P_{K-1} and P_i are the probability that the system are in state F_{ij} , F_o , F_{K-1} , and E_i in equilibrium respectively. Hence the stationary availability P_A is given by (3.39) explicitly in closed form.

Remark 1) $\lambda^*=0$, we have from (3.38) and (3.39)

$$U_K(0) = \left\{ \left(1 + \sum_{j=1}^M \lambda_j K_j \right) \left[\bar{f}_0((N-1)\lambda_0 + \lambda) a_{N-1}^{(1)}(0) / N\lambda_0 + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} (1 - \bar{f}_0(n\lambda_0 + \lambda)) a_n^{(1)}(0) / (n\lambda_0 + \lambda) \right] + \sum_{n=K}^{N-1} (-1)^{n-K} K \lambda_0 \binom{n}{K} [K_0 - (1 - \bar{f}_0(n\lambda_0 + \lambda)) / (n\lambda_0 + \lambda)] a_n^{(1)}(0) / (n\lambda_0 + \lambda) \right\}^{-1} \quad (3.40)$$

Equation (3.40) coincides with the formula (30)' of model 5 in M. Kodama [8].

Remark 2) $\lambda=0$, we have from (3.38) and (3.39)

$$P_A = U_K(0) \left\{ \bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) a_{N-1}^{(1)}(0) / (N\lambda_0 + \lambda^*) + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} [(1 - \bar{f}_0(n\lambda_0 + \lambda^*)) / (n\lambda_0 + \lambda^*)] a_n^{(1)'}(0) \right\}, \quad (3.41)$$

where

$$U_K(0) = \left\{ (1 + \lambda^* K^*_{m+M}) \bar{f}_0((N-1)\lambda_0 + \lambda^*) a_{N-1}^{(1)}(0) / (N\lambda_0 + \lambda^*) \right.$$

$$+ \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} a_n^{(1)'}(0) \cdot [n\lambda_0 K_0 / (n\lambda_0 + \lambda^*) + \lambda^* (1 - \bar{f}_0(n\lambda_0 + \lambda^*)) (K_{m+M}^* + 1 / (n\lambda_0 + \lambda^*)) / (n\lambda_0 + \lambda^*)] \}^{-1} . \quad (3.42)$$

Equations (3.41) and (3.42) coincide with the formulas (29) and (28) in M. Kodama et. al [9], respectively.

(ii) $1 \leq K \leq N-1$, $m=N$

(3.2), (3.3), (3.4), (3.7), (3.8), (3.9) and (3.10) remain the same. Three new equations are added.

$$[d/dt + N\lambda_0 + \lambda + \lambda^*]P_N(t) = \int_0^t \mu_0(x)P_{N-1}(t, x)dx + \sum_{j=1}^M \int_0^t \mu_j(x)P_{Nj}(t, x)dx + \int_0^t \mu_{N+M}^*(x)Q_0(t, x)dx , \quad (3.43)$$

$$[\partial/\partial t + \partial/\partial x + \mu_{N+M}^*(x)]Q_0(t, x) = 0 , \quad (3.44)$$

$$P_i(t, 0) = \delta_{i, N-1} N\lambda_0 P_N(t) + \int_0^t \mu_0(x)P_{i-1}(t, x)dx + \sum_{j=1}^M \int_0^t \mu_j(x)P_{ij}(t, x)dx , \quad (3.45)$$

By the similar method we obtain

$$\bar{P}_N(s) = [\bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*)U_{N-1}(s, 0) + 1 + \lambda^* \bar{g}_{N+M}^*(s) \sum_{i=K}^N \bar{P}_i(s)] / \alpha(s, N, \lambda^*) , \quad (3.46)$$

$$\bar{P}_i(s) = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} U_n(s, 0) [1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)] / (s + n\lambda_0 + \lambda + \lambda^*) , \quad 1 \leq i \leq N-1 , \quad (3.47)$$

$$U_n(s, 0) = a_n^{(2)'}(s)U_K(s, 0) + b_n^{(2)*'}(s) , \quad K+1 \leq n \leq N-1 , \quad (3.48)$$

$$\bar{P}_A(s) = \sum_{i=K}^N \bar{P}_i(s) = k_1^{(2)}(s)U_K(s, 0) + k_0^{(2)}(s) , \quad (3.49)$$

$$U_K(s, 0) = \left\{ \binom{N}{K} [(N-K)N\lambda_0(1 + \lambda^* \bar{g}_{N+M}^*(s)k_0^{(2)}(s)) - \alpha(s, K, \lambda^*) \cdot \bar{f}(s + (N-1)\lambda_0 + \lambda + \lambda^*)b_{N-1}^{(1)*'}(s)] / \alpha(s, N, \lambda^*) + \sum_{n=K}^{N-1} (-1)^{n-K} \left[\binom{n}{K-1} \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*) + \binom{n}{K} K\lambda_0(\bar{f}_0(s) - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*) \right] b_n^{(1)*'}(s) \right\} \left\{ 1 - \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) - \lambda \bar{f}(s)(1 - \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*)) / (s + K\lambda_0 + \lambda + \lambda^*) + \binom{N}{K} [\alpha(s, K, \lambda^*) \cdot \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*)a_{N-1}^{(1)'}(s) - (N-K)\lambda_0 \lambda^* \bar{g}_{N+M}^*(s)k_1^{(2)}(s)] / \alpha(s, N, \lambda^*) - \sum_{n=K}^{N-1} (-1)^{n-K} \left[\binom{n}{K-1} \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*) + \binom{n}{K} K\lambda_0 \cdot (\bar{f}_0(s) - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*) \right] a_n^{(1)'}(s) \right\}^{-1} , \quad (3.50)$$

$$a_n^{(2)'}(s) = a_n^{(1)'}(s) + \lambda^* \bar{g}_{N+M}^*(s)k_1^{(2)}(s)b_n^{(1)*}(s) , \quad K+1 \leq n \leq N-1 , \quad (3.51)$$

$$b_n^{(2)*'}(s) = (1 + \lambda^* \bar{g}_{N+M}^*(s)k_0^{(2)}(s))b_n^{(1)*}(s) , \quad K+1 \leq n \leq N-1 , \quad (3.52)$$

$$k_1^{(2)}(s) = \left\{ 1 - \lambda^* \bar{g}_{N+M}^*(s) \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} [(1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*)] \right.$$

$$\begin{aligned}
 & \cdot b_n^{(1)*}(s) \Big\}^{-1} \Big\{ \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) a_{N-1}^{(1)}(s) / \alpha(s, N, \lambda^*) \\
 & + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} [(1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*)] a_n^{(1)}(s) \Big\} , \\
 & \hspace{15cm} (3.53)
 \end{aligned}$$

$$\begin{aligned}
 k_0^{(2)}(s) = & \left\{ 1 - \lambda^* \bar{g}_{N+M}^*(s) [1 + \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)*}(s)] / \alpha(s, N, \lambda^*) \right. \\
 & - \lambda^* \bar{g}_{N+M}^*(s) \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} [(1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*)] \\
 & \cdot b_n^{(1)*}(s) \Big\}^{-1} \Big\{ [1 + f_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)*}(s)] / \alpha(s, N, \lambda^*) \\
 & + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} [(1 - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)) / (s + n\lambda_0 + \lambda + \lambda^*)] b_n^{(1)*}(s) \Big\} . \\
 & \hspace{15cm} (3.54)
 \end{aligned}$$

Stationary availability (3.22)-(3.24), (3.27)-(3.31) remain the same. Three new equations are added :

$$\begin{aligned}
 [N\lambda_0 + \lambda + \lambda^*] P_N = & \int_0^\infty \mu_0(x) P_{N-1}(x) dx + \sum_{j=1}^M \int_0^\infty \mu_j(x) P_{Nj}(x) dx \\
 & + \int_0^\infty \mu_{N+M}^*(x) Q_0(x) dx , \\
 & \hspace{15cm} (3.55)
 \end{aligned}$$

$$[d/dx + \mu_{N+M}^*(x)] Q_0(x) = 0 , \quad (3.56)$$

$$P_i(0) = \delta_{i, N-1} N \lambda_0 P_N + \int_0^\infty \mu_0(x) P_{i-1}(x) dx + \sum_{j=1}^M \int_0^\infty \mu_j(x) P_{ij}(x) dx . \quad (3.57)$$

By the similar method we obtain

$$P_N = \{ \bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) U_{N-1}(0) + \lambda^* \sum_{i=K}^N P_i \} / (N\lambda_0 + \lambda^*) , \quad (3.58)$$

$$P_i = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} U_i(0) (1 - f_0(i\lambda_0 + \lambda + \lambda^*)) / (i\lambda_0 + \lambda + \lambda^*) , \quad K \leq i \leq N-1 , \quad (3.59)$$

$$\begin{aligned}
 P_{K-1} = & \sum_{n=K}^N (-1)^{n-K} \\
 & \binom{n}{K} U_n(0) [K\lambda_0 / (n\lambda_0 + \lambda + \lambda^*)] [K_0 - (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*)] , \\
 & \hspace{15cm} (3.60)
 \end{aligned}$$

$$U_i(0) = a_i^{(2)'}(0) U_K(0) , \quad (3.61)$$

$$P_{ij} = \lambda_j K_j P_i , \quad K \leq i \leq N , \quad 1 \leq j \leq M , \quad (3.62)$$

$$Q_0 = \lambda^* K_{N+M}^* \sum_{i=K}^N P_i = \lambda^* K_{N+M}^* P_A , \quad (3.62)$$

$$P_A = \sum_{i=K}^N P_i = k_1^{(2)}(0) U_K(0) , \quad (3.64)$$

$$a_i^{(2)'}(0) = a_i^{(1)}(0) + \lambda^* k_1^{(2)}(0) b_i^{(1)*}(0) , \quad (3.65)$$

$$\begin{aligned}
 U_K(0) = & \left\{ k_1^{(2)}(0) \left[1 + \lambda^* K_{N+M}^* + \sum_{j=1}^M \lambda_j K_j \right] + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} [K\lambda_0 / (n\lambda_0 + \lambda + \lambda^*)] \right. \\
 & \cdot K_0 - (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*) \Big\} a_n^{(1)'}(0) \Big\}^{-1} , \\
 & \hspace{15cm} (3.66)
 \end{aligned}$$

(iii) $2 \leq K = N$, $m = N$ (3.10), (3.43) and (3.44) remain the same. Five new equations are added :

$$[\partial/\partial t + \partial/\partial x + \mu_0(x)] P_{K-1}(t, x) = 0 , \quad (3.67)$$

$$[\partial/\partial t + \partial/\partial x + \mu_j(x)] P_{Nj}(t, x) = 0 , \quad 1 \leq j \leq M , \quad (3.68)$$

$$P_{Nj}(t, 0) = \lambda_j P_N(t) , \quad 1 \leq j \leq M , \quad (3.69)$$

$$P_{K-1}(t) = N\lambda_0 P_N(t) , \quad (3.70)$$

$$Q_0(t, 0) = \lambda^* P_N(t) . \quad (3.71)$$

The Laplace transform of $P_A(t)$ and $R(t)$, MTSF and P_A are derived easily.

$$\bar{P}_A(s) = \bar{P}_N(s) = \{s + N\lambda_0(1 - \bar{f}_0(s)) + \lambda(1 - \bar{f}(s)) + \lambda^*(1 - \bar{g}_{N+M}^*(s))\}^{-1} , \quad (3.72)$$

$$\bar{P}_{K-1}(s) = N\lambda_0 \bar{P}_N(s)(1 - \bar{f}(s))/s , \quad (3.73)$$

$$\bar{P}_{Nj}(s) = \lambda_j \bar{P}_N(s)(1 - \bar{f}_j(s))/s , \quad 1 \leq j \leq M , \quad (3.74)$$

$$\bar{Q}_0(s) = \lambda^* \bar{P}_N(s)(1 - \bar{g}_{N+M}^*(s))/s , \quad (3.75)$$

$$P_A = P_N = \left(1 + N\lambda_0 K_0 + \sum_{j=1}^M \lambda_j K_j + \lambda^* K_{N+M}^*\right)^{-1} , \quad (3.76)$$

$$P_{K-1} = N\lambda_0 K_0 P_N , \quad (3.77)$$

$$P_{Nj} = \lambda_j K_j P_N , \quad 1 \leq j \leq M , \quad Q_0 = \lambda^* K_{N+M}^* P_N . \quad (3.78)$$

APPENDIXES

Appendix 1

Taking the Laplace transform of (1.1)-(1.9) under the initial condition (3.10), we have

$$[s + N\lambda_0 + \lambda + \lambda^*] \bar{P}_N(s) = \int_0^\infty \mu_0(x) \bar{P}_{N-1}(s, x) dx + \sum_{j=1}^M \int_0^\infty \mu_j(x) \bar{P}_{Nj}(s, x) dx , \quad (A.1)$$

$$[s + \partial/\partial x + i\lambda_0 + \lambda + \lambda^* + \mu_0(x)] \bar{P}_i(s, x) = (1 - \delta_{i, N-1})(i+1)\lambda_0 \bar{P}_{i+1}(s, x) , \quad K \leq i \leq N-1 , \quad (A.2)$$

$$[s + \partial/\partial x + \mu_0(x)] \bar{P}_{K-1}(s, x) = K\lambda_0 \bar{P}_K(s, x) , \quad (A.3)$$

$$[s + \partial/\partial x + \mu_j(x)] \bar{P}_{ij}(s, x) = 0 , \quad K \leq i \leq N-1 , \quad 1 \leq j \leq M , \quad (A.4)$$

$$[s + \partial/\partial x + \mu_{m+M}^*(x)] \bar{Q}_0(s, x) = 0 , \quad (A.5)$$

$$\begin{aligned} \bar{P}_i(s, 0) = & \delta_{i, N-1} N\lambda_0 \bar{P}_N(s) + \int_0^\infty \mu_0(x) \bar{P}_{i-1}(s, x) dx + \sum_{j=1}^M \int_0^\infty \mu_j(x) \bar{P}_{ij}(s, x) dx \\ & + \delta_{i, m} \int_0^\infty \mu_{m+M}^*(x) \bar{Q}_0(s, x) dx , \quad K \leq i \leq N-1 , \end{aligned} \quad (A.6)$$

$$\bar{P}_{K-1}(s, 0) = 0 , \quad (A.7)$$

$$P_{ij}(s, 0) = \lambda_j \bar{P}_i(s) , \quad K \leq i \leq N-1 , \quad 1 \leq j \leq M , \quad (A.8)$$

$$\bar{Q}_0(s, 0) = \lambda^* \sum_{i=K}^N \bar{P}_i(s) \quad (A.9)$$

From (A.4), (A.5), (A.8), (A.9) and (A.1) we obtain

$$\bar{P}_{ij}(s, x) = \lambda_j \bar{P}_i(s) \exp \left[-sx - \int_0^x \mu_j(t) dt \right] , \quad K \leq i \leq N-1 , \quad 1 \leq j \leq M , \quad (A.10)$$

$$\bar{Q}_0(s, x) = \lambda^* \sum_{i=K}^N \bar{P}_i(s) \exp \left[-sx - \int_0^x \mu_{m+M}^*(z) dz \right] \quad (A.11)$$

$$\alpha(s, N, \lambda^*) \bar{P}_N(s) = \int_0^\infty \mu_0(x) \bar{P}_{N-1}(s, x) dx + 1 . \quad (A.12)$$

Using (A.2) and Laplace transform of (3.11), we have

$$[s + \partial/\partial x + i\lambda_0 + \lambda + \lambda^* + \mu_0(x)] \bar{U}_i(s, x) = 0 , \quad K \leq i \leq N-1 , \quad (A.13)$$

which gives

$$\bar{U}_i(s, x) = \bar{U}_i(s, 0) \exp \left[-(s + i\lambda_0 + \lambda + \lambda^*)x - \int_0^x \mu_0(z) dz \right] , \quad (A.14)$$

From (3.12), (A.14) and (A.3) we obtain

$$\bar{P}_i(s, x) = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} \bar{U}_n(s, 0) \exp \left[-(s + n\lambda_0 + \lambda + \lambda^*)x - \int_0^x \mu_0(z) dz \right],$$

$$K \leq i \leq N-1, \quad (\text{A.15})$$

$$\begin{aligned} \bar{P}_{K-1}(s, x) &= \exp \left[- \int_0^x (s + \mu_0(z)) dz \right] \\ &\cdot \left\{ \int_0^x \exp \left[\int_0^x (s + \mu_0(z)) dz \right] K \lambda_0 \bar{P}_K(s, x) dx + \bar{P}_{K-1}(s, 0) \right\} \\ &= \exp \left[-sx - \int_0^x \mu_0(z) dz \right] \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} K \lambda_0 \bar{U}_n(s, 0) \\ &\cdot \left\{ 1 - \exp [-(n\lambda_0 + \lambda + \lambda^*)x] \right\} / [n\lambda_0 + \lambda + \lambda^*] \end{aligned} \quad (\text{A.16})$$

Using (A.10)-(A.12), (A.15) and (A.16) and nothing that $\bar{P}_i(s) = \int_0^\infty \bar{P}_i(s, x) dx$ ($K-1 \leq i \leq N-1$),

$\bar{P}_{ij}(s) = \int_0^\infty \bar{P}_{ij}(s, x) dx$ ($K \leq i \leq N-1, 1 \leq j \leq M$) and $\bar{Q}_0(s) = \int_0^\infty \bar{Q}_0(s, x) dx$, we obtain (3.13)-(3.

17). Next we will calculate $\bar{U}_K(s, 0), \dots, \bar{U}_{N-1}(s, 0)$ (see reference [6]). From (3.11), (A.6), (3.16), (A.11), (A.10) and intermediate manipulation we obtain

$$\begin{aligned} \bar{U}_i(s, 0) &= \sum_{n=i}^{N-1} \binom{n}{i} \bar{P}_n(s, 0) = \sum_{n=i}^{N-1} \binom{n}{i} \left\{ \delta_{n, N-1} N \lambda_0 \bar{P}_N(s) + \int_0^\infty \mu_0(x) \bar{P}_{n-1}(s, x) dx \right. \\ &\quad \left. + \delta_{n, m} \int_0^\infty \mu_{m+M}^*(x) \bar{Q}_0(s, x) dx + \sum_{j=1}^M \int_0^\infty \mu_j(x) \bar{P}_{nj}(s, x) dx \right\} \\ &= \binom{N-1}{i} N \lambda_0 \{ [\bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{N-1}(s, 0) + 1] / \alpha(s, N, \lambda^*) \} \\ &\quad + \int_0^\infty \mu_0(x) \bar{U}_{i-1}(s, x) dx + \int_0^\infty \mu_0(x) \bar{U}_i(s, x) dx - \binom{N}{i} \int_0^\infty \mu_0(x) \bar{U}_{N-1}(s, x) dx \\ &\quad + \binom{m}{i} \bar{g}_{m+M}^*(s) \lambda^* \sum_{j=K}^N \bar{P}_j(s) + \lambda \bar{f}(s) \sum_{n=i}^{N-1} \binom{n}{i} \bar{P}_n(s) \\ &= \binom{N}{i} \{ (N-i)\lambda_0 + [(N-i)\lambda_0 - \alpha(s, N, \lambda^*)] \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{N-1}(s, 0) \} \\ &\quad / \alpha(s, N, \lambda^*) + \bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*) \bar{U}_i(s, 0) \\ &\quad + \bar{f}_0(s + (i-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{i-1}(s, 0) + \lambda \bar{f}(s) [(1 - \bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*)) \\ &\quad / (s + i\lambda_0 + \lambda + \lambda^*)] \bar{U}_i(s, 0) + \binom{m}{i} \lambda^* \bar{g}_{m+M}^*(s) \sum_{j=K}^N \bar{P}_j(s), \end{aligned} \quad (\text{A.17})$$

which gives

$$\begin{aligned} &\{1 - \bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*) - \lambda \bar{f}(s) [(1 - \bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*)) / (s + i\lambda_0 + \lambda + \lambda^*)] \bar{U}_i(s, 0)\} \\ &= \bar{f}_0(s + (i-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{i-1}(s, 0) + \binom{m}{i} \lambda^* \bar{g}_{m+M}^*(s) \sum_{j=K}^N \bar{P}_j(s) \\ &\quad + \binom{N}{i} [(N-i)\lambda_0 - \alpha(s, i, \lambda^*) \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \\ &\quad \cdot \bar{U}_{N-1}(s, 0)] / \alpha(s, N, \lambda^*), \quad K+1 \leq i \leq N-1 \end{aligned} \quad (\text{A.18})$$

Multiplying (A.18) by $V_i^{(1)}(s)$ and adding for $i = K+1, K+2, \dots, i$, we obtain

$$\begin{aligned} &\bar{f}_0(s + i\lambda_0 + \lambda + \lambda^*) V_i^{(1)}(s) \bar{U}_i^{(1)}(s, 0) \\ &= \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) \bar{U}_K(s, 0) + \sum_{j=K+1}^i V_j^{(1)}(s) \left\{ \binom{m}{j} \lambda^* \right. \end{aligned}$$

$$\begin{aligned} & \cdot \bar{g}_{m+M}^*(s) \sum_{\gamma=K}^N \bar{P}_{\gamma}(s) + \binom{N}{j} (N-j)\lambda_0 - \alpha(s, j, \lambda^*) \\ & \cdot \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{N-1}(s, 0) / \alpha(s, N, \lambda^*) \} \end{aligned} \quad (A.19)$$

Setting $i = N-1$, we have

$$\begin{aligned} & \sum_{j=K+1}^N \binom{N}{j} [\alpha(s, j, \lambda^*) / \alpha(s, N, \lambda^*)] V_{j-1}^{(1)}(s) \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{N-1}(s, 0) \\ & = \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) \bar{U}_K(s, 0) + \sum_{j=K+1}^{N-1} \binom{N}{j} (N-j)\lambda_0 V_{j-1}^{(1)}(s) / \alpha(s, N, \lambda^*) \\ & + \lambda^* \bar{g}_{m+M}^*(s) \sum_{j=K}^N \bar{P}_j(s) \sum_{j=K+1}^{N-1} \binom{m}{j} V_{j-1}^{(1)}(s) \end{aligned} \quad (A.20)$$

which gives

$$\bar{U}_{N-1}(s, 0) = a_{N-1}^{(1)}(s) \bar{U}_K(s, 0) + \lambda^* \bar{g}_{m+M}^*(s) \sum_{j=K}^N \bar{P}_j(s) b_{N-1}^{(1)}(s) + b_{N-1}^{(1)*}(s) \quad (A.21)$$

Substituting (A.21) into (A.19), we obtain after some manipulation and simplification

$$\bar{U}_i(s, 0) = a_i^{(1)}(s) \bar{U}_K(s, 0) + \lambda^* \bar{g}_{m+M}^*(s) \sum_{j=K}^N \bar{P}_j(s) b_i^{(1)}(s) + b_i^{(1)*}(s), \quad K+1 \leq i \leq N-1, \quad (A.22)$$

Substituting (A.22) into (3.15) and (3.16) and arranging for $\sum_{j=K}^N \bar{P}_j(s)$ and nothing that

$$\sum_{i=K}^{N-1} \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} f_n(x) = \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} f_n(x), \quad (A.23)$$

, we obtain (3.20).

Substituting (3.20) into (A.22) we have (3.18).

Since

$$\sum_{n=i}^m (-1)^n \binom{m}{n} \binom{n}{i} = \begin{cases} (-1)^i, & m = i, \\ 0, & m \neq i, \end{cases} \quad (A.24)$$

we have

$$\begin{aligned} & \int_0^\infty \mu_0(x) \sum_{n=K}^{N-1} \binom{n}{K} \bar{P}_{n-1}(s, x) dx \\ & = \int_0^\infty \mu_0(x) \left[\bar{P}_{K-1}(s, x) + \sum_{n=K+1}^{N-1} \binom{n}{K} \bar{P}_{n-1}(s, x) \right] dx \\ & = \int_0^\infty \mu_0(x) \bar{P}_{K-1}(s, x) dx + \int_0^\infty \mu_0(x) \sum_{\gamma=K}^{N-1} \binom{\gamma+1}{K} \bar{P}_{\gamma}(s, x) dx \\ & \quad - \binom{N}{K} \int_0^\infty \mu_0(x) \bar{P}_{N-1}(s, x) dx \\ & = \int_0^\infty \mu_0(x) \bar{P}_{K-1}(s, x) dx + \int_0^\infty \mu_0(x) \bar{U}_K(s, x) dx \\ & \quad + \sum_{r=K}^{N-1} \binom{r}{K-1} \int_0^\infty \mu_0(x) \sum_{n=r}^{N-1} (-1)^{n-r} \binom{n}{r} \bar{U}_n(s, x) dx - \binom{N}{K} \int_0^\infty \mu_0(x) \bar{U}_{N-1}(s, x) dx \\ & = \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} K\lambda_0 \bar{U}_n(s, 0) [\bar{f}_0(s) - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)] / (n\lambda_0 + \lambda + \lambda^*) \end{aligned}$$

$$\begin{aligned}
 & + \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) \bar{U}_K(s, 0) + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K-1} \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*) U_n(s, 0) \\
 & - \binom{N}{K} \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{N-1}(s, 0) ,
 \end{aligned} \tag{A.25}$$

Noting that (3.11) and (3.12) do not define for $i = K-1$, we obtain by using arguments analogous to those in the calculation of (A.22),

$$\begin{aligned}
 \bar{U}_K(s, 0) &= \sum_{n=K}^{N-1} \binom{n}{K} \bar{P}_n(s, 0) \\
 &= \binom{N-1}{K} N\lambda_0 \bar{P}_N(s) + \int_0^\infty \mu_0(x) \sum_{n=K}^{N-1} \binom{n}{K} \bar{P}_{n-1}(s, x) dx \\
 &\quad + \sum_{j=1}^M \int_0^\infty \mu_j(x) \sum_{n=K}^{N-1} \binom{n}{K} \bar{P}_{n,j}(s, x) dx + \binom{m}{K} \int_0^\infty \mu_{m+M}^*(x) \bar{Q}_0^*(s, x) dx \\
 &= \binom{N-1}{K} N\lambda_0 \{ \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) \bar{U}_{N-1}(s, 0) + 1 \} / a(s, N, \lambda^*) \\
 &\quad + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} K\lambda_0 \bar{U}_n(s, 0) \\
 &\quad \cdot [\bar{f}_0(s) - \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*)] / (n\lambda_0 + \lambda + \lambda^*) + \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*) \bar{U}_K(s, 0) \\
 &\quad + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K-1} \bar{f}_0(s + n\lambda_0 + \lambda + \lambda^*) \bar{U}_n(s, 0) \\
 &\quad - \binom{N}{K} \bar{f}_0(s + (N-1)\lambda_0 + \lambda + \lambda^*) U_{N-1}(s, 0) \\
 &\quad + \lambda \bar{f}(s) \bar{U}_K(s, 0) (1 - \bar{f}_0(s + K\lambda_0 + \lambda + \lambda^*)) / (s + K\lambda_0 + \lambda + \lambda^*) \\
 &\quad + \binom{m}{K} \lambda^* \bar{g}_{m+M}^*(s) \sum_{j=K}^N \bar{P}_j(s) .
 \end{aligned} \tag{A.26}$$

Substituting (3.20) and (3.18) into (A.26), we obtain (3.19) after some manipulation and simplification.

Appendix 2

From (3.24), (3.25), (3.28)-(3.30) and (3.21) we obtain

$$P_{ij}(x) = \lambda_j P_i \exp \left[- \int_0^x \mu_j(y) dy \right] , \tag{A.27}$$

$$Q_0(x) = \lambda^* \sum_{j=K}^N P_j \exp \left[- \int_0^x \mu_{m+M}^*(y) dy \right] , \tag{A.28}$$

$$[N\lambda_0 + \lambda^*] P_N = \int_0^\infty \mu_0(x) P_{N-1}(x) dx \tag{A.29}$$

Since the similar discrete transform

$$U_i(x) = \sum_{n=i}^{N-1} \binom{n}{i} P_n(x) , \quad K \leq i \leq N-1 , \tag{A.30}$$

$$P_i(x) = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} U_n(x) , \quad K \leq i \leq N-1 , \tag{A.31}$$

by (3.22) we get

$$[d/dx + i\lambda_0 + \lambda + \lambda^* + \mu_0(x)] U_i(x) = 0 , \quad K \leq i \leq N-1 \tag{A.32}$$

which gives

$$U_i(x) = U_i(0) \exp \left[-(i\lambda_0 + \lambda + \lambda^*)x - \int_0^x \mu_0(y) dy \right]. \quad (\text{A.33})$$

From (A.31), (A.33), (3.23) and (3.27) we have

$$P_i(x) = \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} U_n(0) \exp \left[-(n\lambda_0 + \lambda + \lambda^*)x - \int_0^x \mu_0(y) dy \right], \quad K \leq i \leq N-1, \quad (\text{A.34})$$

$$P_{K-1}(x) = \exp \left[- \int_0^x \mu_0(y) dy \right] \sum_{n=K}^{N-1} \binom{n}{K} K\lambda_0 U_n(0) \{1 - \exp [-(n\lambda_0 + \lambda + \lambda^*)x]\} / (n\lambda_0 + \lambda + \lambda^*) \quad (\text{A.35})$$

Using (A.27)-(A.29), (A.34) and (A.35) and noting that $P_{ij} = \int_0^\infty P_{ij}(x) dx$, ($K \leq i \leq N$, $1 \leq j \leq M$),

$Q_0 = \int_0^\infty Q_0(x) dx$, and $P_i = \int_0^\infty P_i(x) dx$, $K-1 \leq i \leq N-1$, we obtain (3.32)-(3.36).

Next we will calculate $U_K(0), \dots, U_{N-1}(0)$ (see reference[6]). By the similar argument to the case of (A.18) and using (A.30) and (3.26) we get

$$\begin{aligned} & \{1 - \bar{f}_0(i\lambda_0 + \lambda + \lambda^*) - \lambda[1 - \bar{f}_0(i\lambda_0 + \lambda + \lambda^*)] / (i\lambda_0 + \lambda + \lambda^*)\} U_i(0) \\ &= \bar{f}_0((i-1)\lambda_0 + \lambda + \lambda^*) U_{i-1}(0) \\ & \quad - (i\lambda_0 + \lambda^*) \binom{N}{i} \bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) U_{N-1}(0) / (N\lambda_0 + \lambda^*) + \lambda^* \binom{m}{i} \sum_{j=K}^N P_j, \\ & \quad K+1 \leq i \leq N-1 \end{aligned} \quad (\text{A.36})$$

Multiplying (A.36) by $V_i^{(1)}(0)$ and adding for $i=K+1, K+2, \dots, i$, we obtain

$$\begin{aligned} & \bar{f}_0(i\lambda_0 + \lambda + \lambda^*) V_i^{(1)}(0) U_i(0) = \bar{f}_0(K\lambda_0 + \lambda + \lambda^*) U_K(0) + \lambda^* \sum_{n=K}^N P_n \sum_{j=K+1}^i \binom{m}{j} V_j^{(1)}(0) \\ & \quad - [\bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) U_{N-1}(0) / (N\lambda_0 + \lambda^*)] \sum_{j=K+1}^i \binom{N}{j} (j\lambda_0 + \lambda^*) V_j^{(1)}(0), \end{aligned} \quad (\text{A.37})$$

and setting $i=N-1$ we obtain after some manipulation

$$U_{N-1}(0) = a_{N-1}^{(1)}(0) U_K(0) + \lambda^* \sum_{j=K}^N P_j b_{N-1}^{(1)}(0) \quad (\text{A.38})$$

Substituting (A.38) into (A.37), we have

$$U_i(0) = a_i^{(1)}(0) U_K(0) + \lambda^* \sum_{j=K}^N P_j b_i^{(1)}(0), \quad K \leq i \leq N-1 \quad (\text{A.39})$$

Using (3.34), (3.35), (A.39) and (A.23) we obtain

$$\begin{aligned} \sum_{j=K}^N P_i &= \bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) \left[a_{N-1}^{(1)}(0) U_K(0) + \lambda^* b_{N-1}^{(1)}(0) \sum_{i=K}^N P_i \right] / (N\lambda_0 + \lambda^*) \\ & \quad + \sum_{i=K}^{N-1} \sum_{n=i}^{N-1} (-1)^{n-i} \binom{n}{i} \left[a_n^{(1)}(0) U_K(0) \right. \\ & \quad \left. + \lambda^* b_n^{(1)}(0) \sum_{i=K}^N P_i \right] [1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)] / (n\lambda_0 + \lambda + \lambda^*) \\ &= \left[\bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) a_{N-1}^{(1)}(0) / (N\lambda_0 + \lambda^*) \right. \\ & \quad \left. + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} a_n^{(1)}(0) (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*) \right] U_K(0) \\ & \quad + \sum_{i=K}^N P_i \left[\lambda^* \bar{f}_0((N-1)\lambda_0 + \lambda + \lambda^*) b_{N-1}^{(1)}(0) / (N\lambda_0 + \lambda^*) \right] \end{aligned}$$

$$+ \lambda^* \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n-1}{K-1} b_n^{(1)}(0) (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*) \Big]$$

which gives (3.39).

Substituting (3.39) into (A. 39) we have (3.27). From (3.81)-(3.36), (3.37) and (3.39) we get

$$\begin{aligned} 1 &= \sum_{n=K}^N P_n + P_{K-1} + P_N \sum_{j=1}^M \lambda_j K_j + \sum_{j=1}^M \lambda_j K_j \sum_{i=K}^{N-1} P_i + \lambda^* K_{m+M}^* \sum_{i=K}^N P_i \\ &= k_1^{(1)}(0) \left[1 + \lambda^* K_{m+M}^* + \sum_{j=1}^M \lambda_j K_j \right] U_K(0) + \sum_{n=K}^{N-1} (-1)^{n-K} \binom{n}{K} [K\lambda_0 / (n\lambda_0 + \lambda + \lambda^*)] \\ &\quad \cdot [K_0 - (1 - \bar{f}_0(n\lambda_0 + \lambda + \lambda^*)) / (n\lambda_0 + \lambda + \lambda^*)] a_n^{(1)'}(0) U_K(0) \end{aligned}$$

which gives (3.38).

REFERENCES

- [1] D. K. Kulshrestha, "Reliability of a parallel redundant complex system," *Opns. Res.* 16, No. 1, 1968.
- [2] D. K. Kulshrestha, "Reliability of a repairable multicomponent system with redundancy in parallel," *IEEE Trans. on Reliability*, 19, No. 2, 1970.
- [3] H. Nakamichi, J. Fukuta, S. Takamatsu, and M. Kodama, "Reliability consideration on a repairable multicomponent system with redundancy in parallel," *J. Opns. Res. Soc. of Japan*, 17, No. 1, 1974.
- [4] M. Kodama, "Probabilistic analysis of a multicomponent seriesparallel system under preemptive repeat repair discipline," *Opns. Res.* 24, No. 23, 1976.
- [5] M. Kodama and I. Sawa, "Reliability and maintainability of a multicomponent series-parallel system under several repair disciplines," *Microelectron. Reliab.* 22, No. 6, 1982.
- [6] M. Kodama and I. Sawa, "Reliability and maintainability of a multicomponent series-parallel system with simultaneous failure and repair priorities," to be published in *Microelectron. Reliab.*
- [7] M. Kodama, "Reliability and maintainability of a multicomponent series-parallel system I," *Society of Political Economy*, Kyushu University, Vol 46, Nos. 1-2, 1981 (in Japanese).
- [8] M. Kodama, "Reliability and maintainability of a multicomponent series-parallel system II," *Society of Political Economy*, Kyushu University, Vol. 46, Nos. 4-5, 1981 (in Japanese).
- [9] M. Kodama and K. Adachi, "On the steady state availability of a single-sever k-out-of-n: G system with simultaneous failure and general repair distribution," *Bulletin of Nagoya Institute of Technology*, Vol. 30, 1978.
- [10] N. K. Jaiswal, *Priority Queues*, Academic Press, New York, 1968.