

Tongue Microbiota Composition and Dental Caries Experience in Primary School Children

張, 代熙

<https://hdl.handle.net/2324/4496013>

出版情報 : Kyushu University, 2021, 博士 (歯学), 課程博士
バージョン :

権利関係 : (c) 2021 Zhang et al. This is an openaccess article distributed under the terms of the Creative Commons Attribution 4.0 International license.



Tongue Microbiota Composition and Dental Caries Experience in Primary School Children

Daixi Zhang,^a Toru Takeshita,^{a,b} Michiko Furuta,^a  Shinya Kageyama,^a Mikari Asakawa,^a Koki Nambu,^a Yoshihisa Yamashita^a

^aSection of Preventive and Public Health Dentistry, Division of Oral Health, Growth and Development, Faculty of Dental Science, Kyushu University, Fukuoka, Japan

^bOBT Research Center, Faculty of Dental Science, Kyushu University, Fukuoka, Japan

Daixi Zhang and Toru Takeshita contributed equally to this work. Author order was determined in order of increasing seniority.

ABSTRACT The tongue microbiota of elderly adults is composed of two cohabiting commensal groups and their ratios are related to the number of teeth with dental caries experience. In this study, the variation in the tongue microbiota of primary school children and its relationship with the dental caries experience were investigated. We examined the tongue microbiota of 138 children aged 6 to 7 years and 11 to 12 years (61 and 77 children, respectively) who underwent annual dental examinations. The bacterial composition was determined by sequencing the V1-V2 region of the 16S rRNA gene. Cooccurrence network analysis indicated two groups of cohabiting predominant commensals in the tongue microbiota of children. The microbiota in children without a history of dental caries showed significantly higher relative abundances of one of the cohabiting groups, primarily composed of *Neisseria subflava*, *Porphyromonas pasteri*, and *Fusobacterium periodonticum*, compared to that in children with a history of dental caries, which is consistent with that of elderly adults with fewer teeth with dental caries experience. Linear discriminant analysis effect size (LEfSe) further identified *Streptococcus oralis* subsp. *dentisani*, belonging to the aforementioned commensal group, as a discriminant species in children without dental caries experience aged 6 to 7 years and 11 to 12 years. Our results describe the tongue microbiota composition of primary school children without history of dental caries and support the possibility that dental caries experience is accompanied by a shift in the tongue microbiota.

IMPORTANCE Dental caries is now considered to be caused by acids produced by the overall dental plaque microbiota rather than by specific pathogens. This study focused on the relationship between dental caries experience and the variations in tongue microbiota, which is adjacent but separate from the dental plaque microbiota. Our results demonstrated that the tongue microbiota of primary school children with no history of dental caries experience was composed of predominant commensals with different relative abundances compared to those present in children with dental caries experience, suggesting that dental caries experience is accompanied by a shift in the tongue microbiota. The maintenance of a healthy tongue microbiota may indirectly contribute to the prevention of dental caries.

KEYWORDS 16S rRNA, children, dental caries, tongue microbiota

Dental caries is a common chronic oral disease characterized by the demineralization of tooth tissue and decomposition of organic matter, which leads to cavitation or tooth decay (1). The etiology of this disease involves acid production by the dental plaque microbiota, which accumulates on the tooth surface. Of the bacterial members, mutans streptococci were implicated as the single causal agent of the disease in earlier studies, and their virulence has been studied extensively (2, 3). However,

Citation Zhang D, Takeshita T, Furuta M, Kageyama S, Asakawa M, Nambu K, Yamashita Y. 2021. Tongue microbiota composition and dental caries experience in primary school children. *mSphere* 6:e01252-20. <https://doi.org/10.1128/mSphere.01252-20>.

Editor Garret Suen, University of Wisconsin-Madison

Copyright © 2021 Zhang et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Yoshihisa Yamashita, yoshi@dent.kyushu-u.ac.jp.

Received 7 December 2020

Accepted 19 March 2021

Published 28 April 2021

even though the microorganism is undoubtedly associated with the initiation of dental caries (2, 4), its presence is not indispensable for dental caries development. Currently, the prevailing “ecological hypothesis” proposes that changes in the oral environment alter the dental plaque microbiota into an aciduric one, composed of diverse bacteria with acidogenic potential, which leads to the development and progression of dental caries (5, 6).

The tongue microbiota is another complex oral microbial community that is adjacent but unattached to the plaque microbiota, and has a community composition distinct from that of the dental plaque microbiota (7–9). Interestingly, a population-based study in elderly adults showed a significant relationship between the ratio of two groups of cohabiting predominant commensals in the tongue microbiota and the number of teeth with dental caries experience (10). A similar result was observed in another population-based study in middle-aged and elderly adults analyzing salivary microbial populations (11), which showed dominant tongue microbes (7–9). The findings from these studies indicate that dental caries development might be accompanied by dysbiotic microbiota on decayed teeth as well as elsewhere in the oral cavity. However, there remains a possibility that the shift in the tongue microbiota in middle-aged and elderly adults is caused by other changes occurring during their life course, including periodontal diseases and tooth loss.

Here, we studied the tongue microbiota of primary school children aged 6 to 7 years and 11 to 12 years with recent complaints of dental caries. We determined the tongue microbiota composition and analyzed the variation in the composition and its relationship with dental caries.

RESULTS

In this study, the tongue microbiota composition of 138 primary school children aged 6 to 7 years and 11 to 12 years (61 and 77 children, respectively) was studied based on the sequencing of the 16S rRNA gene using a next-generation sequencer, the Ion PGM system (Thermo Fisher Scientific, Waltham, MA, USA). The samples provided 3,478,851 quality reads corresponding to the V1-V2 regions of the 16S rRNA gene sequences, which were subsequently assigned to 302 species-level operational taxonomic units (OTUs) based on 97% sequence similarity. Of these, 21 OTUs corresponding to typical oral commensals were the predominant taxa present in the tongue microbiota of the children (Table 1), and their mean relative abundance in each individual was $79.0 \pm 6.4\%$ (mean \pm standard deviation [SD]).

Cooccurrence network analysis using FastSpar (12) suggested that the predominant commensals were a part of two distinct cohabiting groups in the tongue microbiota (Fig. 1A). The members of each cohabiting group are listed in Table S1 in the supplemental material. The first group was composed of bacterial taxa such as *Prevotella histicola*, *Veillonella parvula*, *Streptococcus salivarius*, *Prevotella pallens*, and *Prevotella melaninogenica* (commensal group I, Fig. 1B), whereas the second group was composed of *Neisseria subflava*, *Porphyromonas pasteri*, *Fusobacterium periodonticum*, *Haemophilus parainfluenzae*, and *S. oralis* subsp. *dentisani* (commensal group II, Fig. 1C). There was an inverse relationship between the relative abundances of the two commensal groups (Spearman’s correlation coefficient = -0.91) (Fig. S1).

The relative abundances of the commensal groups in the tongue microbiota varied according to the dental health conditions, whereas no notable differences were observed between children of different genders, ages, and dentition stages (Fig. 2). The commensal group II was significantly more predominant in children without dental caries experience and those with better dental hygiene (lower debris index-simplified [DI-S]) (13). A higher relative abundance of this commensal group was also observed in the microbiota of children with lower mutans streptococci levels in their saliva (lower Dentocult SM score). Alternatively, commensal group I was more predominant in the microbiota of children with higher Dentocult SM and DI-S scores. The ratios of group II commensals to group I commensals were significantly higher in children without dental caries experience and those with lower Dentocult SM and DI-S scores.

TABLE 1 Twenty-one predominant operational taxonomic units (OTUs) with mean relative abundances of >1% in the tongue microbiota of 138 primary school children

OTU no.	Bacterial taxa corresponding to each OTU ^a	% Mean relative abundance \pm SD	% Detection rate
OTU4	<i>Neisseria subflava</i> (476)	12.1 \pm 9.2	100
OTU3	<i>Prevotella melaninogenica</i> (469)	12.0 \pm 7.3	100
OTU7	<i>Streptococcus salivarius</i> (755)	7.4 \pm 6.0	100
OTU10	<i>Veillonella parvula</i> (161)	6.8 \pm 3.6	100
OTU2	<i>Rothia mucilaginosa</i> (681)	5.1 \pm 4.0	99.3
OTU1	<i>Fusobacterium periodonticum</i> (201)	4.5 \pm 3.5	100
OTU6	<i>Haemophilus parainfluenzae</i> (718)	4.4 \pm 4.5	100
OTU9	<i>Granulicatella adiacens</i> (534)	3.9 \pm 2.1	100
OTU8	<i>Porphyromonas pasteri</i> (279)	3.8 \pm 4.0	97.1
OTU5	<i>Prevotella histicola</i> (298)	3.5 \pm 4.5	87.7
OTU30	<i>Streptococcus infantis</i> (431)	2.2 \pm 2.1	100
OTU25	<i>Actinomyces graevenitzi</i> (866)	2.2 \pm 2.9	94.9
OTU13	<i>Prevotella pallens</i> (714)	1.6 \pm 1.3	94.9
OTU11	<i>Alloprevotella</i> sp. (308)	1.5 \pm 1.8	95.7
OTU22	<i>Actinomyces odontolyticus</i> (701)	1.3 \pm 0.9	99.3
OTU19	Genus <i>Streptococcus</i> ^b	1.3 \pm 1.5	98.6
OTU170	<i>Rothia mucilaginosa</i> (681)	1.1 \pm 0.9	99.3
OTU15	<i>Leptotrichia</i> sp. (417)	1.1 \pm 2.3	86.2
OTU46	<i>Streptococcus</i> sp. (064)	1.1 \pm 0.8	100
OTU12	<i>Solobacterium moorei</i> (678)	1.1 \pm 0.9	97.8
OTU133	<i>Prevotella</i> sp. (313)	1.0 \pm 1.7	79.0

^aOral taxon IDs in eHOMD are given in parentheses following bacterial names.

^bNo blast hit with \geq 98.5% identity was found in the expanded Human Oral Microbiome database (eHOMD).

No notable relationship was observed between these clinical parameters and the alpha diversity indices of the tongue microbiota, except for a higher microbial richness observed in children with a higher DI-S score (Fig. S2). In contrast, the principal-coordinate analysis (PCoA) plot based on unweighted UniFrac metrics showed differences between children with and without dental caries experience in the direction of principal coordinate 1 and differences between children aged 6 to 7 years and 11 to 12 years in the direction of principal coordinate 2 (Fig. 3). The significant relationship with the overall composition of the tongue microbiota was confirmed by permutational multivariate analysis of variance (PERMANOVA $F = 2.92$ and $P = 0.001$; $F = 2.04$ and $P = 0.02$, respectively), whereas there was no significant relationship with gender, dentition stage, DI-S, or a presence of decayed teeth. Significant differences were observed in the relative abundances of *Streptococcus* and *Neisseria* among the four groups categorized by age and dental caries experience (Fig. S3). The children could also be differentiated according to the mutans streptococci levels in the saliva in the direction of principal coordinate 1 in the UniFrac plot (PERMANOVA $F = 2.72$ and $P < 0.001$) (Fig. S4).

While the overall tongue microbiota composition was different between children aged 6 to 7 years and 11 to 12 years, the commensal group II was significantly more predominant in caries-free children than in children with dental caries experience from either age group (6 to 7 years and 11 to 12 years) (Fig. 4). Linear discriminant analysis effect size (LEfSe) further revealed ten and six taxa associated with the presence or absence of dental caries experience in children aged 6 to 7 years and 11 to 12 years, respectively (Tables 2 and 3). Of these, *S. oralis* subsp. *dentisani*, which belongs to commensal group II, was identified as a discriminant taxon associated with the absence of dental caries experience in both age groups. In contrast, *Streptococcus parasanguinis*, a member of the commensal group I, was detected in the tongue microbiota of children with dental caries experience in both age groups.

DISCUSSION

This study demonstrated that the tongue microbiota of primary school children is dominated by common oral commensals, including *N. subflava* and *P. melaninogenica*




FIG 1 Cooccurrence network in the tongue microbiota of 138 children built from SparCC correlation coefficients between sequence abundances. Each node corresponds to a distinct operational taxonomic unit (OTU) and correlations with values greater than 0.5 and P values less than 0.001 are represented as edges. The size of each node indicates the mean relative abundance of each OTU. (A) All OTUs are shown as nodes in the diagram. The OTUs belonging to two major networks (commensal group I and II) are colored red and blue, respectively. (B) Commensal group I. The bacterial name corresponding to each OTU is described at each node. Oral taxon IDs from the expanded Human Oral Microbiome database are provided in parentheses following the bacterial name. The bacterial names corresponding to minor OTUs (relative abundance < 0.5%) are indicated using numbers and described in Table S1 in the supplemental material. (C) Commensal group II. The bacterial name corresponding to each OTU is described at each node. Oral taxon IDs in the expanded Human Oral Microbiome database are provided in parentheses following the bacterial name.

(Table 1), and is composed of two competing cohabiting groups (Fig. 1). The two cohabiting groups were mostly consistent with those found in the tongue microbiota of elderly adults in a previous study (10); the findings suggest that the tongue microbiota, composed of specific indigenous taxa, has already been established in childhood during the transitional stage between primary and permanent dentition.

This study also characterized the tongue microbiota of primary school children with no history of dental caries. One of the two cohabiting predominant commensal groups, primarily composed of *N. subflava*, *H. parainfluenzae*, *P. pasteri*, and *F. periodonticum* (commensal group II), was significantly more predominant in the microbiota of children without dental caries experience (Fig. 2), which is consistent with the microbiota composition of elderly adults with less dental caries experience (10). Significant differences in the relative abundances of the commensal group were observed in the microbiota of children aged 6 to 7 years and 11 to 12 years who were affected by dental caries more recently (Fig. 2, Fig. 4). These results support the possibility that dental caries experience is accompanied by a shift in the tongue microbiota.

The genera *Neisseria*, *Haemophilus*, and *Rothia* were associated with good oral health, whereas *Prevotella* species (predominant in commensal group I) and *Streptococcus mutans*




FIG 2 Relative abundances of two cohabiting commensal groups in the tongue microbiota and the ratios of group II commensals to group I commensals (log₂-transformed) of primary school children with different conditions. The dots indicate the mean, and the error bars indicate the 95% confidence interval (CI). ***, $P < 0.001$; *, $P < 0.05$ by Student's t test (two categories) or analysis of variance (three categories).

were implicated in dental caries in a recent study using a shotgun metagenomics approach (14). Another metagenomic study demonstrated that the genera *Prevotella*, *Veillonella*, and *Atopobium* (belonging to commensal group I in this study) were more abundant in the supragingival microbiota of children with dental caries than in those without dental caries (15). In addition, considering that oral *Prevotella* species have been implicated as dental caries-associated taxa in several studies previously (16–19), our results are consistent with previous evidence. However, a large-scale twin study showed that *Prevotella pallens* and *Veillonella* spp. are heritable taxa associated with the absence of dental caries, which is discordant with our findings (20). Further careful investigation is needed to elucidate the relationship between the ratios of the abundances of predominant oral commensals and dental caries.

This cross-sectional study was unable to clarify whether the shifted tongue microbiota is responsible for dental caries or is a result of the development of dental caries. Nevertheless, considering that salivary levels of mutans streptococci were clearly correlated with the ratio of the two cohabiting commensals in the tongue microbiota (Fig. 2), it is unlikely that dental restorations, such as resin and metal, result in a shift in the tongue microbiota. The shift in the tongue microbiota coincides with the overgrowth of mutans streptococci in the plaque microbiota induced by an oral environment susceptible to dental caries. Future longitudinal and microbiological studies will help clarify the relationship between the tongue microbiota and susceptibility to dental caries.

Of the members of the commensal group II, *S. oralis* subsp. *dentisani* was identified as a key discriminant species for children with no history of dental caries experience from both age groups (6 to 7 years and 11 to 12 years) (Tables 2 and 3). *S. oralis* subsp. *dentisani* was first isolated from the dental plaque of caries-free individuals (21). This species was further reported to inhibit the growth of oral pathogens, including mutans streptococci, through the production of potent bacteriocins, as well as through its pH-buffering capacity via an arginolytic pathway (22), which makes it promising as an oral




FIG 3 A principal coordinate analysis plot showing the similarity relationship among the tongue microbiota of 138 primary school children using an unweighted UniFrac distance metric. The points corresponding to the children with different categories of dental caries experience (left panel) and age groups (right panel) are depicted using different colors. The two axes explain 20.7% and 10.6% of the variance, respectively.

probiotic with anti-caries potential (23). Our results further support the potential of this taxon as a species beneficial for oral health, although its role in lowering susceptibility to dental caries remains unclear.

Another noteworthy species in this study was *S. parasanguinis*, which was a discriminant taxon for children with dental caries experience from either age group (Tables 2 and 3) and was previously implicated in the dental plaque microbiota of children with




FIG 4 Relative abundances of commensal group II in the tongue microbiota of primary school children aged 6 to 7 years and 11 to 12 years with and without dental caries experience; *, $P < 0.05$, Student's *t* test.

TABLE 2 Bacterial taxa corresponding to differentially abundant OTUs in the tongue microbiota of children aged 6 to 7 years with and without dental caries experience

OTU no.	Bacterial taxa corresponding to each OTU ^a	% Relative abundance \pm SD		LDA score ^c
		Without previous dental caries (n = 33)	With previous dental caries (n = 28)	
Differentially abundant in children with dental caries experience				
OTU10	<i>Veillonella parvula</i> (161)	6.30 \pm 3.15	8.53 \pm 4.12	4.09
OTU38	<i>Streptococcus parasanguinis</i> (411)	0.66 \pm 0.76	1.44 \pm 1.66	3.62
OTU151	Genus <i>Streptococcus</i> ^b	0.1 \pm 0.16	0.34 \pm 0.43	3.13
OTU136	<i>Leptotrichia</i> sp. (498)	0 \pm 0	0.02 \pm 0.05	3.07
OTU188	Genus <i>Prevotella</i> ^b	0 \pm 0	0.01 \pm 0.02	3.04
OTU168	Genus <i>Fusobacterium</i> ^b	0 \pm 0	0.01 \pm 0.01	3.02
Differentially abundant in caries-free children				
OTU103	Genus <i>Fusobacterium</i> ^b	0.01 \pm 0.01	0.00 \pm 0.01	3.10
OTU66	<i>Streptococcus australis</i> (073)	0.58 \pm 0.73	0.39 \pm 0.68	3.13
OTU129	Order <i>Clostridiales</i> ^b	0.01 \pm 0.01	0 \pm 0	3.48
OTU34	<i>Streptococcus oralis</i> subsp. <i>dentisani</i> (058)	1.49 \pm 1.47	0.55 \pm 0.87	3.71

^aOral taxon IDs in eHOMD are given in parentheses following bacterial names.

^bNo blast hit with \geq 98.5% identity was found in the expanded Human Oral Microbiome database (eHOMD).

^cLinear discriminant analysis (LDA) effect size (LEfSe) was conducted, and OTUs with a high LDA score ($>$ 3.0) are shown.

severe early childhood dental caries (18). However, these commensal streptococci were also reported to restrict the growth and biofilm formation of *S. mutans* in the presence of nitrite (24). Therefore, careful attention should be paid to the association of *S. parasanguinis* with dental caries.

This study assessed the tongue microbiota composition in primary school children with no previous experience of dental caries based on the results of annual dental examination throughout their school years. However, decayed deciduous teeth lost prior to the first dental examination were not accounted for. Nevertheless, all children were younger than 7 years and 3 months in age at their first dental examination and had almost complete primary dentition except for the occasional loss of some incisors. The details of dental caries experience, including that in deciduous teeth, were mostly covered in this study. In contrast, the absence of dental caries diagnosis based on the observation of enamel caries is a major limitation of this study. Dental caries is evaluated based on visual inspection during dental examination conducted in schools; therefore, only obvious tooth decay without air drying is considered in dental caries diagnosis. A more rigorous diagnosis, such as that based on the International Caries Detection and

TABLE 3 Bacterial taxa corresponding to differentially abundant OTUs in the tongue microbiota of children aged 11 to 12 years with and without dental caries experience

OTU no.	Bacterial taxa corresponding to each OTU ^a	% Relative abundance \pm SD		LDA score ^b
		Without previous dental caries (n = 16)	With previous dental caries (n = 61)	
Differentially abundant in children with dental caries experience				
OTU5	<i>Prevotella histicola</i> (298)	1.92 \pm 2.81	3.93 \pm 5.09	4.03
OTU25	<i>Actinomyces graevenitzi</i> (866)	1.33 \pm 2.11	2.36 \pm 3.33	3.74
OTU26	<i>Lachnospiraceae</i> bacterium (096)	0.08 \pm 0.23	0.62 \pm 1.51	3.42
OTU38	<i>Streptococcus parasanguinis</i> (411)	0.67 \pm 0.78	1.05 \pm 0.86	3.40
Differentially abundant in caries-free children				
OTU161	<i>Actinomyces johnsonii</i> (849)	0.00 \pm 0.01	0 \pm 0	3.45
OTU34	<i>Streptococcus oralis</i> subsp. <i>dentisani</i> (058)	1.80 \pm 1.73	0.7 \pm 0.86	3.70

^aOral taxon IDs in eHOMD are given in parentheses following bacterial names.

^bLinear discriminant analysis (LDA) effect size (LEfSe) was conducted, and OTUs with a high LDA score ($>$ 3.0) are shown.

Assessment System (ICDAS) criteria (25), would be required for the accurate evaluation of dental caries experience. Another limitation is this study was the inability to prevent the children from cleaning their tongues before sample collection. Although the tongue microbiota composition is relatively stable over time (26), it is possible that the tongue microbiota of some children differs from the usual composition.

Our results suggest that a shifted tongue microbiota is present in the background of dental caries. Although dental caries is undoubtedly caused by dysbiosis in the dental plaque microbiota, a broader view of the entire oral cavity as a microbial consortium might add a fresh perspective in defining the etiology of dental caries.

MATERIALS AND METHODS

Ethics approval. All parents or guardians of child participants understood the nature of the study and provided written informed consent. The Ethics Committee of Kyushu University approved the study design, as well as the procedure for obtaining informed consent (reference no. 30-303). All experiments were performed in accordance with the approved guidelines.

Study population, sample collection, and dental examination. We enrolled 173 children aged 6 to 7 years and 11 to 12 years in two primary schools in Hisayama town, Fukuoka prefecture, Japan, who underwent annual dental examinations. Prior to the dental examination, the salivary levels of mutans streptococci were evaluated using a Dentocult SM strip (Orion Diagnostica, Espoo, Finland), and a sample of the tongue coating was collected by swabbing the center of the tongue dorsum five times using a HydraFlock swab (Puritan Medical Products, Guilford, ME, USA). The collected tongue coating samples were stored in 200 μ l of lysis buffer containing 10 mM Tris-HCl, 1 mM EDTA, and 1% sodium dodecyl sulfate and transferred to the laboratory on ice. In addition to determining the number of present, decayed, or filled teeth, the oral hygiene status was evaluated using the DI-S score of the simplified oral hygiene index (OHI-S) (13) for six selected teeth (positions 16, 11, 26, 36, 41, and 46) based on visual inspection during the dental examination. The target teeth were modified to six deciduous teeth (positions 55, 51, 65, 75, 81, and 85) in children aged 6 to 7 years because they had primary dentition. For children aged 6 to 7 years, the absence of decayed or filled deciduous and permanent teeth at this dental examination was considered the absence of dental caries experience. However, for children aged 11 to 12 years, the absence of dental caries experience was defined based on the continued absence of decayed and filled teeth in the dental examination records throughout their school years, including the record for the current year. Three children without decayed and filled teeth in this examination were excluded from the analysis owing to incomplete dental examination data recorded during their school years. The number of decayed and filled deciduous teeth in children aged 6 to 7 years and 11 to 12 years ranged from 0 to 10 (median, 0) and 0 to 6 (median, 0), respectively. The number of decayed and filled permanent teeth in children aged 6 to 7 years and 11 to 12 years ranged from 0 to 3 (median, 0) and 0 to 3 (median, 0), respectively.

Bacterial community analysis based on 16S rRNA gene sequencing. DNA was extracted from the tongue coating samples using the bead-beating method, as described previously (27). Thirty children were excluded from the analysis as insufficient DNA was extracted due to a technical failure in the process. The V1-V2 region of the 16S rRNA gene from each sample was amplified using the following primers: 8F (5'-AGA GTT TGA TYM TGG CTC AG-3') with the Ion Torrent adaptor A and a sample-specific 8-base tag sequence, and 338R (5'-TGC TGC CTC CCG TAG GAG T-3') with the Ion Torrent trP1 adaptor sequence. The V1-V2 region can be used to discriminate among most *Streptococcus* species, which is the most predominant and clinically important oral taxon at the species level, and its use is currently recommended in studies on the oral microbiota (28). PCR amplification, purification, and quantification were performed as described previously (11). The indexed PCR amplicons were normalized and pooled to prepare a sequencing library. Emulsion PCR and library enrichment were performed using the Ion One Touch 2 system (Thermo Fisher Scientific) with an Ion PGM Hi-Q View OT2 kit (Thermo Fisher Scientific). Sequencing was conducted using an Ion PGM system (Thermo Fisher Scientific) with an Ion PGM Hi-Q View sequencing kit (Thermo Fisher Scientific).

Data processing. The obtained sequence reads were filtered based on their quality using a script written in R. The reads were excluded from the analysis if they contained ≤ 240 bases, had an average quality score of ≤ 25 , did not include the correct forward primer sequence, did not include the correct reverse primer sequence (one mismatch was allowed), or had > 7 nucleotide homopolymer. The remaining reads were assigned to the appropriate sample by examining the tag sequences. Operational taxonomic unit (OTU) clustering was conducted using UPARSE (29), with a minimum pairwise identity of 97%. The taxonomy of representative sequences of each OTU was determined using BLAST against 998 16S rRNA gene sequences (eHOMD 16S rRNA RefSeq, version 15.1) deposited in the expanded Human Oral Microbiome Database (30). Nearest-neighbor species with 98.5% identity were identified as candidate taxa for each OTU. The taxonomy of sequences with no appropriate hits was determined using the RDP classifier (31) with a minimum support threshold of 80% up to the genus level. The alpha diversity index and the relative abundance of each OTU were calculated following rarefaction with a depth of 5,000 reads per sample using R. The rarefaction curves are shown in Fig. S5. Two children were excluded from the analysis as the number of obtained reads was insufficient.

Statistical analysis. All statistical analyses were conducted using R version 3.5.2. Cooccurrence network analysis was performed for 283 OTUs detected from ≥ 2 individuals using FastSpar (12), which is a

fast and parallelizable implementation of the SparCC algorithm (32), with 5,000 bootstraps. Cooccurrence relationships with correlation values of >0.5 and P values of <0.001 were represented as edges, and displayed using the gplot function in the sna library. The relative abundances of each bacterial group were compared using Student's t test between two categories and analysis of variance (ANOVA) among three or more categories. The similarity relationship of overall tongue microbiota structure, assessed using the UniFrac metric (33), was presented in a principal-coordinate analysis (PCoA) plot. Permutational multivariate analysis of variance (PERMANOVA) was used to assess the effects of general and dental conditions on tongue microbiota composition based on 9,999 permutations. Discriminant OTUs of children with or without dental caries experience were explored using the LEfSe approach (34).

Data availability. The obtained sequence data were deposited in the DDBJ Sequence Read Archive under accession no. DRA011686.

SUPPLEMENTAL MATERIAL

Supplemental material is available online only.

FIG S1, PDF file, 0.2 MB.

FIG S2, PDF file, 0.2 MB.

FIG S3, PDF file, 0.2 MB.

FIG S4, PDF file, 0.3 MB.

FIG S5, PDF file, 0.3 MB.

TABLE S1, DOCX file, 0.02 MB.

ACKNOWLEDGMENTS

This study was supported by Grants-in-Aid by JSPS KAKENHI grant numbers JP19H03863 (T.T.), JP19K22722 (T.T.), JP20H03901 (Y.Y.), and JP20K21682 (Y.Y.) from the Japan Society for the Promotion of Science.

We thank the staff of the Division of Health and Welfare of the Hisayama Town Office for their cooperation in this study.

REFERENCES

- Pitts NB, Zero DT, Marsh PD, Ekstrand K, Weintraub JA, Ramos-Gomez F, Tagami J, Twetman S, Tsakos G, Ismail A. 2017. Dental caries. *Nat Rev Dis Primers* 3:17030. <https://doi.org/10.1038/nrdp.2017.30>.
- Loesche WJ. 1986. Role of *Streptococcus mutans* in human dental decay. *Microbiol Rev* 50:353–380. <https://doi.org/10.1128/MR.50.4.353-380.1986>.
- Kuramitsu HK, He X, Lux R, Anderson MH, Shi W. 2007. Interspecies interactions within oral microbial communities. *Microbiol Mol Biol Rev* 71:653–670. <https://doi.org/10.1128/MMBR.00024-07>.
- Bratthall D. 1970. Demonstration of five serological groups of streptococcal strains resembling *Streptococcus mutans*. *Odontol Revy* 21:143–152.
- Marsh PD. 2006. Dental plaque as a biofilm and a microbial community—implications for health and disease. *BMC Oral Health* 6 Suppl 1:S14. <https://doi.org/10.1186/1472-6831-6-S1-S14>.
- Takahashi N, Nyvad B. 2011. The role of bacteria in the caries process: ecological perspectives. *J Dent Res* 90:294–303. <https://doi.org/10.1177/0022034510379602>.
- Mager DL, Ximenez-Fyvie LA, Haffajee AD, Socransky SS. 2003. Distribution of selected bacterial species on intraoral surfaces. *J Clin Periodontol* 30:644–654. <https://doi.org/10.1034/j.1600-051x.2003.00376.x>.
- Kageyama S, Takeshita T, Asakawa M, Shibata Y, Takeuchi K, Yamanaka W, Yamashita Y. 2017. Relative abundance of total subgingival plaque-specific bacteria in salivary microbiota reflects the overall periodontal condition in patients with periodontitis. *PLoS One* 12:e0174782. <https://doi.org/10.1371/journal.pone.0174782>.
- Segata N, Haake SK, Mannon P, Lemon KP, Waldron L, Gevers D, Huttenhower C, Izard J. 2012. Composition of the adult digestive tract bacterial microbiome based on seven mouth surfaces, tonsils, throat and stool samples. *Genome Biol* 13:R42. <https://doi.org/10.1186/gb-2012-13-6-r42>.
- Asakawa M, Takeshita T, Furuta M, Kageyama S, Takeuchi K, Hata J, Ninomiya T, Yamashita Y. 2018. Tongue microbiota and oral health status in community-dwelling elderly adults. *mSphere* 3:e00332-18. <https://doi.org/10.1128/mSphere.00332-18>.
- Takeshita T, Kageyama S, Furuta M, Tsuboi H, Takeuchi K, Shibata Y, Shimazaki Y, Akifusa S, Ninomiya T, Kiyohara Y, Yamashita Y. 2016. Bacterial diversity in saliva and oral health-related conditions: the Hisayama Study. *Sci Rep* 6:22164. <https://doi.org/10.1038/srep22164>.
- Watts SC, Ritchie SC, Inouye M, Holt KE. 2019. FastSpar: rapid and scalable correlation estimation for compositional data. *Bioinformatics* 35:1064–1066. <https://doi.org/10.1093/bioinformatics/bty734>.
- Greene JC, Vermillion JR. 1964. The simplified oral hygiene index. *J Am Dent Assoc* 68:7–13. <https://doi.org/10.14219/jada.archive.1964.0034>.
- Baker JL, Morton JT, Dinis M, Alvarez R, Tran NC, Knight R, Edlund A. 2021. Deep metagenomics examines the oral microbiome during dental caries, revealing novel taxa and co-occurrences with host molecules. *Genome Res* 31:64–74. <https://doi.org/10.1101/gr.265645.120>.
- Al-Hebshi NN, Baraniya D, Chen T, Hill J, Puri S, Tellez M, Hasan NA, Colwell RR, Ismail A. 2019. Metagenome sequencing-based strain-level and functional characterization of supragingival microbiome associated with dental caries in children. *J Oral Microbiol* 11:1557986. <https://doi.org/10.1080/20002297.2018.1557986>.
- Nadkarni MA, Caldon CE, Chhour KL, Fisher IP, Martin FE, Jacques NA, Hunter N. 2004. Carious dentine provides a habitat for a complex array of novel *Prevotella*-like bacteria. *J Clin Microbiol* 42:5238–5244. <https://doi.org/10.1128/JCM.42.11.5238-5244.2004>.
- Teng F, Yang F, Huang S, Bo C, Xu ZZ, Amir A, Knight R, Ling J, Xu J. 2015. Prediction of early childhood caries via spatial-temporal variations of oral microbiota. *Cell Host Microbe* 18:296–306. <https://doi.org/10.1016/j.chom.2015.08.005>.
- Tanner AC, Kent RL, Jr., Holgerson PL, Hughes CV, Loo CY, Kanasi E, Chalmers NI, Johansson I. 2011. Microbiota of severe early childhood caries before and after therapy. *J Dent Res* 90:1298–1305. <https://doi.org/10.1177/0022034511421201>.
- Hurley E, Barrett MPJ, Kinirons M, Whelton H, Ryan CA, Stanton C, Harris HMB, O'Toole PW. 2019. Comparison of the salivary and dentinal microbiome of children with severe-early childhood caries to the salivary microbiome of caries-free children. *BMC Oral Health* 19:13. <https://doi.org/10.1186/s12903-018-0693-1>.
- Gomez A, Espinoza JL, Harkins DM, Leong P, Saffery R, Bockmann M, Torralba M, Kuelbs C, Kodukula R, Inman J, Hughes T, Craig JM, Highlander SK, Jones MB, Dupont CL, Nelson KE. 2017. Host genetic control of the oral microbiome in health and disease. *Cell Host Microbe* 22:269–278. <https://doi.org/10.1016/j.chom.2017.08.013>.

21. Camelo-Castillo A, Benitez-Paez A, Belda-Ferre P, Cabrera-Rubio R, Mira A. 2014. *Streptococcus dentisani* sp. nov., a novel member of the mitis group. *Int J Syst Evol Microbiol* 64:60–65. <https://doi.org/10.1099/ijs.0.054098-0>.
22. Lopez-Lopez A, Camelo-Castillo A, Ferrer MD, Simon-Soro A, Mira A. 2017. Health-associated niche inhabitants as oral probiotics: the case of *Streptococcus dentisani*. *Front Microbiol* 8:379. <https://doi.org/10.3389/fmicb.2017.00379>.
23. Ferrer MD, Lopez-Lopez A, Nicolescu T, Perez-Vilaplana S, Boix-Amoros A, Dzidic M, Garcia S, Artacho A, Llena C, Mira A. 2020. Topic application of the probiotic *Streptococcus dentisani* improves clinical and microbiological parameters associated with oral health. *Front Cell Infect Microbiol* 10:465. <https://doi.org/10.3389/fcimb.2020.00465>.
24. Scofield J, Michalek S, Harber G, Eipers P, Morrow C, Wu H. 2019. Dietary nitrite drives disease outcomes in oral polymicrobial infections. *J Dent Res* 98:1020–1026. <https://doi.org/10.1177/0022034519855348>.
25. Ismail AI, Sohn W, Tellez M, Amaya A, Sen A, Hasson H, Pitts NB. 2007. The International Caries Detection and Assessment System (ICDAS): an integrated system for measuring dental caries. *Community Dent Oral Epidemiol* 35:170–178. <https://doi.org/10.1111/j.1600-0528.2007.00347.x>.
26. Zhou Y, Gao H, Mihindukulasuriya KA, La Rosa PS, Wylie KM, Vishnivetskaya T, Podar M, Warner B, Tarr PI, Nelson DE, Fortenberry JD, Holland MJ, Burr SE, Shannon WD, Sodergren E, Weinstock GM. 2013. Biogeography of the ecosystems of the healthy human body. *Genome Biol* 14:R1. <https://doi.org/10.1186/gb-2013-14-1-r1>.
27. Takeshita T, Nakano Y, Kumagai T, Yasui M, Kamio N, Shibata Y, Shiota S, Yamashita Y. 2009. The ecological proportion of indigenous bacterial populations in saliva is correlated with oral health status. *ISME J* 3:65–78. <https://doi.org/10.1038/ismej.2008.91>.
28. Wade WG, Prosdociemi EM. 2020. Profiling of oral bacterial communities. *J Dent Res* 99:621–629. <https://doi.org/10.1177/0022034520914594>.
29. Edgar RC. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nat Methods* 10:996–998. <https://doi.org/10.1038/nmeth.2604>.
30. Escapa IF, Chen T, Huang Y, Gajare P, Dewhirst FE, Lemon KP. 2018. New insights into human nostril microbiome from the expanded human oral microbiome database (eHOMD): a resource for the microbiome of the human aerodigestive tract. *mSystems* 3:e00187-18. <https://doi.org/10.1128/mSystems.00187-18>.
31. Cole JR, Wang Q, Cardenas E, Fish J, Chai B, Farris RJ, Kulam-Syed-Mohideen AS, McGarrell DM, Marsh T, Garrity GM, Tiedje JM. 2009. The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res* 37:D141–5. <https://doi.org/10.1093/nar/gkn879>.
32. Friedman J, Alm EJ. 2012. Inferring correlation networks from genomic survey data. *PLoS Comput Biol* 8:e1002687. <https://doi.org/10.1371/journal.pcbi.1002687>.
33. Lozupone C, Knight R. 2005. UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microbiol* 71:8228–8235. <https://doi.org/10.1128/AEM.71.12.8228-8235.2005>.
34. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, Huttenhower C. 2011. Metagenomic biomarker discovery and explanation. *Genome Biol* 12:R60. <https://doi.org/10.1186/gb-2011-12-6-r60>.