

## A Stereographic Representation of Euler's Formula to Show Spirals and Topological Similarities to Micro-Structures in Ruminants and Forages

Shimojo, Masataka

Laboratory of Animal Feed Science, Division of Animal Science, Department of Animal and Marine Bioresource Sciences, Faculty of Agriculture, Kyushu University

Ikeda, Kentaro

Research Fellow, Faculty of Agriculture, Kyushu University

Asano, Yoki

Research Fellow, Faculty of Agriculture, Kyushu University

Ishiwaka, Reiko

他

<https://doi.org/10.5109/4528>

---

出版情報：九州大学大学院農学研究院紀要. 48 (1/2), pp.71-75, 2003-10-01. 九州大学大学院農学研究  
院

バージョン：

権利関係：



## A Stereographic Representation of Euler's Formula to Show Spirals and Topological Similarities to Micro-Structures in Ruminants and Forages

Masataka SHIMOJO<sup>1</sup>, Yoki ASANO<sup>1)</sup>, Kentaro IKEDA<sup>1)</sup>, Reiko ISHIWAKA<sup>1)</sup>,  
Tao SHAO<sup>2)</sup>, Hiroyuki SATO<sup>3)</sup>, Manabu TOBISA, Yutaka NAKANO<sup>4)</sup>,  
Noriko OHBA<sup>5)</sup> Yasukatsu YANO<sup>6)</sup> and Yasuhisa MASUDA

Laboratory of Animal Feed Science, Division of Animal Science, Department of  
Animal and Marine Bioresource Sciences, Faculty of Agriculture,  
Kyushu University, Fukuoka 812–8581, Japan

(Received June 30, 2003 and accepted July 15, 2003)

This study suggests that the stereographic representation of Euler's formula and its conjugate complex in the combination with or without their phase shifts gives spirals showing topological similarities to micro-structures in ruminants and forages.

### INTRODUCTION

We suggested that some mathematical treatments of Euler's formula gave a symbolic description of field–forage–ruminant relationships (Shimojo *et al.*, 2003a, b), and in addition, exponential functions with base  $e$  used for the growth analysis of ruminants and forages (Shimojo *et al.*, 2003c). These show an aspect of macro-structures in ruminant agriculture.

It is reported by Yoshida (2000) that a stereographic representation of Euler's formula gives spirals. If precise shapes are not considered, spirals suggested from Euler's formula might be expected to show topological similarities to helical structures in living things (Watson *et al.*, 1987; Calladine and Drew, 1992; Voet *et al.*, 1999, for example).

The present study was designed to investigate spirals suggested from Euler's formula and their topological similarities to helical structures in ruminants and forages.

### EULER'S FORMULA AND ITS STEREOGRAPHIC DESCRIPTIONS

#### Euler's formula

Euler's formula is given by

---

<sup>1)</sup> Research Fellow, Faculty of Agriculture, Kyushu University

<sup>2)</sup> Visiting Research Scientist from Lanzhou Institute of Animal Science and Veterinary Medicine of CAAS, Lanzhou Gansu Province of the People's Republic of China

<sup>3)</sup> Laboratory of Animal Feed Science, Division of Animal Science, Department of Animal and Marine Bioresource Sciences, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

<sup>4)</sup> Kyushu University Farm, Fukuoka 811–2307

<sup>5)</sup> Research Student, School of Agriculture, Kyushu University

<sup>6)</sup> Former Technical Specialist

\* Corresponding author (E-mail: mshimojo@agr.kyushu-u.ac.jp)

$$\exp(i\beta) = \cos\beta + i\sin\beta, \quad (1)$$

where  $i$ =imaginary unit,  $\beta$ =real variable.

### Stereographic description of Euler's formula gives a right-handed spiral

There are the following two sequences on the complex plane (Shimojo *et al.*, 2003a); a series of anti-clockwise  $\pi/2$  rotations [ $\times i$ ],


$$(\cos\beta + i\sin\beta) \xrightarrow{\times i} (-\sin\beta + i\cos\beta) \xrightarrow{\times i} (-\cos\beta - i\sin\beta) \xrightarrow{\times i} (\sin\beta - i\cos\beta), \quad (2)$$

and a series of clockwise  $\pi/2$  rotations [ $\times (-i)$ ],

$$(\sin\beta + i\cos\beta) \xrightarrow{\times (-i)} (\cos\beta - i\sin\beta) \xrightarrow{\times (-i)} (-\sin\beta - i\cos\beta) \xrightarrow{\times (-i)} (-\cos\beta + i\sin\beta). \quad (3)$$

In the present study,  $\cos\beta + i\sin\beta$  in (2),  $\cos\beta - i\sin\beta$  in (3) and their phase shifts will be described stereographically to show spirals.

Yoshida (2000) shows that the stereographic description of Euler's formula using coordinates  $(\beta, \cos\beta, i\sin\beta)$  gives a right-handed spiral (Fig. 1a). Its planar projection is also shown in Fig. 1b. In the present study coordinate axes are eliminated in order to show the structure of spirals only, otherwise there will occur interferences of waves when spirals are combined.



**Fig. 1.** A stereographic representation of Euler's formula [ $\exp(i\beta) = \cos\beta + i\sin\beta$ ] using coordinates  $(\beta, \cos\beta, i\sin\beta)$ . (a) A right-handed spiral. (b) A planar projection of the right-handed spiral. (c) The planar projection of  $\exp(i\beta)$  in the combination with  $\exp\{i(\beta + 2\pi/3)\}$ . In each case the right-hand end of spirals is the starting point.

**Phase shifts of Euler's formula**

Let us take some phase shifts for  $\exp(i\beta)$ , for example  $2\pi/3$ ,  $4\pi/3$  and  $\pi$ . Thus,

$$\exp\left(i\frac{2\pi}{3}\right) \cdot \exp(i\beta) = \exp\left\{i\left(\beta + \frac{2\pi}{3}\right)\right\}, \quad (4)$$

$$\exp\left(i\frac{4\pi}{3}\right) \cdot \exp(i\beta) = \exp\left\{i\left(\beta + \frac{4\pi}{3}\right)\right\}, \quad (5)$$

$$\exp(i\pi) \cdot \exp(i\beta) = \exp\{i(\beta + \pi)\}, \quad (6)$$

It is shown in (4)~(6) that the form of Euler's formula is kept invariant when given any phase shift.

**Combining Euler's formula and its phase shifts in the stereographic description**

(A) Combination of (1) and (4)


Combining  $\exp(i\beta)$  and  $\exp\{i(\beta + 2\pi/3)\}$  gives a pair of right-handed spirals (Fig. 1c).

(B) Combination of (1), (4) and (5)

The combination of  $\exp(i\beta)$ ,  $\exp\{i(\beta + 2\pi/3)\}$  and  $\exp\{i(\beta + 4\pi/3)\}$  gives right-handed three spirals (Fig. 2a).

(C) Combination of (1) and (6)

Combining  $\exp(i\beta)$  and  $\exp\{i(\beta + \pi)\}$  gives a pair of right-handed spirals (Fig. 2b).



**Fig. 2.** The planar projection of the stereographic representation of Euler's formula [ $\exp(i\beta)$ ] in the combination with  $\exp\{i(\beta+2\pi/3)\}$  and  $\exp\{i(\beta+4\pi/3)\}$  in (a) and in the combination with  $\exp\{i(\beta+\pi)\}$  in (b). In each case the right-hand end of spirals is the starting point.

### The conjugate complex to Euler's formula gives a left-handed spiral

As shown in (3), the conjugate complex to Euler's formula is given by

$$\exp(-i\beta) = \cos\beta - i\sin\beta. \quad (7)$$


The conjugate complex to Euler's formula gives a left-handed spiral when described stereographically using coordinates  $(\beta, \cos\beta, -i\sin\beta)$ , as shown in Fig. 3a. Its planar projection is also shown in Fig. 3b.

The  $-\pi$  phase shift of  $\exp(-i\beta)$  is given by

$$\exp(-i\pi) \cdot \exp(-i\beta) = \exp\{-i(\beta + \pi)\}. \quad (8)$$

It is shown in (8) that the form of conjugate complex to Euler's formula is kept invariant when given a phase shift of  $-\pi$ .

Combining (7) and (8), namely  $\exp(-i\beta)$  and  $\exp\{-i(\beta + \pi)\}$ , gives a pair of left-handed spirals (Fig. 3c).



**Fig. 3.** A stereographic representation of the conjugate complex to Euler's formula [ $\exp(-i\beta) = \cos\beta - i\sin\beta$ ] using coordinates  $(\beta, \cos\beta, -i\sin\beta)$ . (a) A left-handed spiral. (b) A planar projection of the left-handed spiral. (c) The planar projection of  $\exp(-i\beta)$  in the combination with  $\exp\{-i(\beta + \pi)\}$ . In each case the right-hand end of spirals is the starting point.

### Topological similarities to micro-structures in ruminants and forages

Sine curve and cosine curve are used for the analysis of helical structures in living things (Calladine and Drew, 1992). It is suggested from the present study that stereographic descriptions of Euler's formula and its conjugate complex in the combination with or without their phase shifts show topological similarities to the following helical structures: a right-handed helix [(1)], a left-handed helix [(7)], a right-handed double helix of

two kinds [combining (1) and (4), combining (1) and (6)], a left-handed double helix [combining (7) and (8)], and a right-handed triple helix [combining (1), (4) and (5)]. A feature of the present approach is as follows. (a) Right-handed spirals and left-handed spirals are related to Euler's formula and its conjugate complex, respectively. (b) Double and triple spirals are given by Euler's formula or its conjugate complex in the combination with corresponding phase shifts. (c) The complementary property in the form of double spiral is related to keeping the form of Euler's formula or its conjugate complex invariant with respect to phase shifts. Therefore, the use of stereographic representation of Euler's formula and its conjugate complex in the combination with or without their phase shifts might give a rough image of helical structures in ruminants and forages.

## REFERENCES

- Calladine, C. R. and H. R. Drew 1992 Understanding DNA—The Molecule and How It Works—, Academic Press Limited, London. (Translated into Japanese by Watt, F. and Y. Nishimura, Kyouritsu Publishing Corp., Tokyo, 1996)
- Shimojo, M., K. Ikeda, Y. Asano, R. Ishiwaka, T. Shao, H. Sato, M. Tobisa, Y. Nakano, N. Ohba, Y. Yano and Y. Masuda 2003a A symbolic representation of field–forage–ruminant relationships using polar form on the complex plane. *J. Fac. Agr., Kyushu Univ.*, **47**: 359–366
- Shimojo, M., Y. Asano, K. Ikeda, R. Ishiwaka, T. Shao, H. Sato, M. Tobisa, Y. Nakano, N. Ohba, Y. Yano and Y. Masuda 2003b Complex representation of field–forage–ruminant relationships using symmetric properties of Euler's formula. *J. Fac. Agr., Kyushu Univ.*, **47**: 367–372
- Shimojo, M., K. Ikeda, Y. Asano, R. Ishiwaka, T. Shao, H. Sato, M. Tobisa, Y. Nakano, N. Ohba, Y. Yano and Y. Masuda 2003c Exponential functions with base  $e$  in growth analysis and deriving them from rotations of axes of time described using Euler's formula. *J. Fac. Agr., Kyushu Univ.*, **48**: 65–69
- Voet, D., J. G. Voet and C. W. Pratt 1999 Fundamentals of Biochemistry. John Wiley & Sons, Inc., Publishing Co., Inc., California. (Translated into Japanese by Tamiya, N., M. Muramatsu, T. Yagi and T. Endo, Tokyo Kagaku Dojin, Tokyo, 2001)
- Watson, J. D., N. H. Hopkins, J. W. Roberts, J. A. Steitz and A. M. Weiner 1987 Molecular Biology of the Gene (4th ed.), The Benjamin/Cummings Publishing Co., Inc., California. (Translated into Japanese under the supervision of Matsubara, K., K. Nakamura and K. Miura, Tokyo Denki University Press, Tokyo, 2001)
- Yoshida, T. 2000 Emotion for imaginary number. Tokai University Press, Tokyo. (Written in Japanese)