

## Introducing Complex Numbers into Basic Growth Functions (4) : Hypothetic Appearance of Exponential Function with Base $e$ from the Complex Representation of ' $0+(-1)+1$ '

Shimojo, Masataka

Laboratory of Animal Feed Science, Division of Animal Science, Department of Animal and Marine Bioresource Sciences, Faculty of Agriculture, Kyushu University

Ikeda, Kentarou

Research Fellow, Faculty of Agriculture, Kyushu University

Ishiwaka, Reiko

Research Fellow, Faculty of Agriculture, Kyushu University

Sato, Hiroyuki

Laboratory of Animal Feed Science, Division of Animal Science, Department of Animal and Marine Bioresource Sciences, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

他

<https://doi.org/10.5109/4596>

---

出版情報 : 九州大学大学院農学研究院紀要. 49 (2), pp.349-353, 2004-10-01. Faculty of Agriculture, Kyushu University

バージョン :

権利関係 :



## Introducing Complex Numbers into Basic Growth Functions

### – (IV) Hypothetic Appearance of Exponential Function with Base $e$ from the Complex Representation of ' $0=(-1)+1$ ' –

Masataka SHIMOJO\*, Kentaro IKEDA<sup>1</sup>, Reiko ISHIWAKA<sup>1</sup>, Hiroyuki SATO<sup>2</sup>,  
Yoki ASANO<sup>3</sup>, Manabu TOBISA<sup>4</sup>, Yutaka NAKANO<sup>5</sup>, Noriko OHBA<sup>6</sup>,  
Minako EGUCHI<sup>7</sup> and Yasuhisa MASUDA

Laboratory of Animal Feed Science, Division of Animal Science, Department of Animal  
and Marine Bioresource Sciences, Faculty of Agriculture,  
Kyushu University, Fukuoka 812–8581, Japan  
(Received June 30, 2004 and accepted July 13, 2004)

This study was designed to investigate the hypothetic appearance of exponential function with base  $e$  from the complex representation of ' $0=(-1)+1$ ' and phase shifts. The results obtained were as follows. There was a hypothetic appearance of  $\exp(i\theta)$  from the complex representation of ' $-1$ ' in ' $0=(-1)+1$ ' when subjected to the hypothetic breakdown of multiplication form connecting complex numbers constructing ' $0=(-1)+1$ '. The clockwise  $\pi/2$  rotation of  $\theta$  in  $\exp(i\theta)$  led to  $\exp(\theta)$ . If there were many different complex numbers with their replicas to be left, then this would be given by an equation with continuous phase shift of  $\theta$  in  $\exp(i\theta)$ . The hypothetic inheritance of phase shift of  $\theta$  in  $\exp(i\theta)$  to the change in  $\theta$  in  $\exp(\theta)$  might lead to the calculation of definite integral of  $\exp(\theta)$ . Might the change in  $\theta$  in  $\exp(i\theta)$  and/or  $\exp(\theta)$  be related to the passage of time ( $t$ ) in  $\exp(t)$ ?

## INTRODUCTION

The exponential function with base  $e$  is a function of importance to the basic growth analysis of ruminants (Brody, 1945) and forages (Watson, 1952; Radford, 1967; Hunt, 1990). It was shown hypothetically by Shimojo *et al.* (2003) that the exponential function with base  $e$  was obtained by  $\pm \pi/2$  rotation of axes of variable described using Euler's formula. Shimojo *et al.* (2004a, b, c) reported that ' $0=(-1)+1$ ' was described using the product of complex numbers and the hypothetic breakdown of multiplication form caused incomplete pair disappearances of them to leave some sorts of complex numbers. These two hypothetic phenomena might be related to suggest the appearance of exponential

<sup>1</sup> Research Fellow, Faculty of Agriculture, Kyushu University

<sup>2</sup> Laboratory of Animal Feed Science, Division of Animal Science, Department of Animal and Marine Bioresource Sciences, Graduate School of Bioresource and Bioenvironmental Sciences, Kyushu University

<sup>3</sup> Employed Research Scientist, Miyazaki University, Miyazaki 889–2192, Japan

<sup>4</sup> Faculty of Agriculture, Miyazaki University, Miyazaki 889–2192, Japan

<sup>5</sup> University Farm, Faculty of Agriculture, Kyushu University

<sup>6</sup> Research Student, School of Agriculture, Kyushu University

<sup>7</sup> Technical Official, School of Agriculture, Kyushu University

\* Corresponding author (E-mail: mshimojo@agr.kyushu-u.ac.jp)

function with base  $e$  from ' $0 = (-1) + 1$ '.

This fourth study was designed to investigate the hypothetic appearance of exponential function with base  $e$  from the complex representation of ' $0 = (-1) + 1$ ' and phase shifts.

### COMPLEX REPRESENTATION OF ' $0 = (-1) + 1$ '

#### Complex representation of '-1' to leave complex numbers

The following equation is taken up as an example of complex representation of ' $0 = (-1) + 1$ ' according to reports by Shimojo *et al.* (2004a, b, c). Thus,

$$\begin{aligned}
 0 &= (-1) + 1 \\
 &= [\exp(i\theta) \cdot i\exp(-i\theta) \cdot i\exp(i\theta) \cdot (-\exp(-i\theta)) \\
 &\quad \cdot \{-(-\exp(i\theta))\} \cdot (-i\exp(-i\theta)) \cdot (-i\exp(i\theta)) \cdot \exp(-i\theta)] \\
 &\quad + [\exp(i\theta) \cdot i\exp(-i\theta) \cdot i\exp(i\theta) \cdot (-\exp(-i\theta)) \\
 &\quad \cdot (-\exp(i\theta)) \cdot (-i\exp(-i\theta)) \cdot (-i\exp(i\theta)) \cdot \exp(-i\theta)]. \tag{1}
 \end{aligned}$$

The hypothetic breakdown of multiplication form in (1) leaves a sort of complex number,

$$\begin{aligned}
 &[\exp(i\theta) + i\exp(-i\theta) + i\exp(i\theta) + (-\exp(-i\theta)) \\
 &\quad + \{-(-\exp(i\theta))\} + (-i\exp(-i\theta)) + (-i\exp(i\theta)) + \exp(-i\theta)] \\
 &\quad + [\exp(i\theta) + i\exp(-i\theta) + i\exp(i\theta) + (-\exp(-i\theta)) \\
 &\quad + (-\exp(i\theta)) + (-i\exp(-i\theta)) + (-i\exp(i\theta)) + \exp(-i\theta)] \\
 &= 2\exp(i\theta), \tag{2}
 \end{aligned}$$

where the complex representation of '1' vanishes due to complete pair disappearances of complex numbers constructing '1'.

Each of the other seven complex numbers will be left according to other ways of breakdown, as shown by Shimojo *et al.* (2004c).

If there are many different complex numbers with their replicas to be left, then the following equation will be given after the hypothetic breakdown of multiplication form connecting complex numbers constructing '-1',

$$\begin{aligned}
 &\sum_{h=1}^m \left[ \sum_{k=0}^n \{ \exp(i(\theta_h + 2k\pi)) + i\exp(-i(\theta_h + 2k\pi)) + i\exp(i(\theta_h + 2k\pi)) \right. \\
 &\quad + (-\exp(-i(\theta_h + 2k\pi))) + (-(-\exp(i(\theta_h + 2k\pi)))) + (-i\exp(-i(\theta_h + 2k\pi))) \\
 &\quad \left. + (-i\exp(i(\theta_h + 2k\pi))) + \exp(-i(\theta_h + 2k\pi)) \} \right] \\
 &= 2 \sum_{h=1}^m \left[ \sum_{k=0}^n \exp(i(\theta_h + 2k\pi)) \right], \tag{3}
 \end{aligned}$$

where  $0 < \theta_1 < \theta_2 < \dots < \theta_h < \dots < \theta_m < \pi/2$ .

Thus, there is a hypothetic appearance of  $\exp(i\theta)$  from the complex representation of

'0=(-1)+1', namely from '-1'. It is suggested hypothetically from equation (3) that  $\exp(i\theta_1)$  is the starting point and the phase shift of  $\theta$  will lead to the other complex numbers,  $\exp(i(\theta_m+2n\pi))$  as the last point. If  $m$  and  $n$  are extremely large, then there will be the continuous phase shift of a micro-scale. Broadly speaking at the risk of making mistakes, equation (3) suggests that  $\exp(i\theta)$  with continuous micro-scale phase shift appears hypothetically from the complex representation of '0=(-1)+1'.

### Hypothetic conversion of $\exp(i\theta)$ into $\exp(\theta)$

As shown in the preceding section, there was a product of  $\pm i$  and  $\exp(\pm i\theta)$  in the complex representation of '0=(-1)+1'. However, it was reported by Shimojo *et al.* (2003) that  $\exp(\theta)$  was given by the product of  $-i$  and variable in  $\exp(i\theta)$ . Thus,

$$\exp((-i) \cdot (i\theta)) = \exp(\theta). \quad (4)$$

This is a kind of conversion from the world of complex numbers [ $\exp(i\theta)$ ] into the world of real numbers [ $\exp(\theta)$ ].

If the phase shift of  $\theta$  in  $\exp(i\theta)$  is inherited hypothetically to the change in  $\theta$  in  $\exp(\theta)$ , then the calculation of definite integral of  $\exp(\theta)$  will be allowed.

### Definite integral of $\exp(\theta)$ expanded into infinite series

The definite integral of  $\exp(\theta)$  expanded into infinite series is given by

$$\begin{aligned} \int_{\theta_1}^{\theta_2} \exp(\theta) d\theta &= \int_{\theta_1}^{\theta_2} \left( 1 + \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right) d\theta \\ &= \left[ \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right]_{\theta_1}^{\theta_2} \\ &= \left[ \left( 1 + \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right) + (-1) \right]_{\theta_1}^{\theta_2} \\ &= [\exp(\theta) + (-1)]_{\theta_1}^{\theta_2} \\ &= \exp(\theta_2) - \exp(\theta_1). \end{aligned} \quad (5)$$

Another hypothetic description of definite integral of  $\exp(\theta)$  suggested by Shimojo *et al.* (2004b, c) is

$$\begin{aligned} \int_{\theta_1}^{\theta_2} \exp(\theta) d\theta &= \int_{\theta_1}^{\theta_2} \left( 1 + \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right) d\theta \\ &= \left[ \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right]_{\theta_1}^{\theta_2} + \{(-1)+1\} - \{(-1)+1\}. \end{aligned} \quad (6)$$

If the complex representation of two sets of '(-1)+1' is followed by the hypothetic breakdown of multiplication form connecting complex numbers constructing '(-1)+1', then the following will be given (Shimojo *et al.*, 2004b, c)

$$\begin{aligned}
\int_{\theta_1}^{\theta_2} \exp(\theta) d\theta &= \left[ \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right]_{\theta_1}^{\theta_2} + \{(-1)+1\} - \{(-1)+1\} \\
&= \left[ \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right]_{\theta_1}^{\theta_2} \\
&\quad + [\exp(i\theta) + i\exp(-i\theta) + i\exp(i\theta) + (-\exp(-i\theta)) \\
&\quad + \{-(-\exp(i\theta))\} + (-i\exp(-i\theta)) + (-i\exp(i\theta)) + \exp(-i\theta)] \\
&\quad - [\exp(i\theta) + i\exp(-i\theta) + i\exp(i\theta) + (-\exp(-i\theta)) \\
&\quad + \{-(-\exp(i\theta))\} + (-i\exp(-i\theta)) + (-i\exp(i\theta)) + \exp(-i\theta)] \\
&= \left[ \frac{\theta}{1!} + \frac{\theta^2}{2!} + \cdots + \frac{\theta^n}{n!} + \cdots \right]_{\theta_1}^{\theta_2} + 2\exp(i\theta) - 2\exp(i\theta) \\
&= \exp(\theta_2) - \exp(\theta_1), \tag{7}
\end{aligned}$$

where the complex representation of '1' is not shown due to complete pair disappearances of complex numbers with their opposites.

In equation (7) there are two types of changes in  $\theta$ ; (i) the change from  $\theta_1$  to  $\theta_2$  for the definite integral of  $\exp(\theta)$ , (ii) a series of  $\pm \pi/2$  phase shifts of  $\theta$  in  $\exp(\pm i\theta)$  occurring due to the hypothetic breakdown of multiplication form connecting complex numbers to construct '-1'.

### Conclusions and suggestions from the present study

It is shown hypothetically from equations (1)~(7) that  $\exp(i\theta)$  appears from '-1' in ' $0 = (-1) + 1$ ' when described using complex numbers and subjected to the hypothetic breakdown of multiplication form connecting them to construct ' $0 = (-1) + 1$ '. The clockwise  $\pi/2$  rotation of the variable in  $\exp(i\theta)$  gives  $\exp(\theta)$ . The hypothetic inheritance of phase shift in  $\exp(i\theta)$  to  $\exp(\theta)$  might lead to the calculation of definite integral of  $\exp(\theta)$  under the change in  $\theta$ . Might the change in  $\theta$  in  $\exp(i\theta)$  and/or  $\exp(\theta)$  be related to the passage of time ( $t$ ) in  $\exp(t)$ ? However, what causes these hypothetic phenomena to occur is not known.

### REFERENCES

- Brody, S. 1945 Time relations of growth of individuals and populations. In "Bioenergetics and growth", Reinhold Publishing Corporation, New York, pp. 484-574
- Hunt, R. 1990 Basic Growth Analysis. Unwin Hyman Ltd., London.
- Radford, P. J. 1967 Growth analysis formulae—their use and abuse. *Crop Sci.*, **7**: 171-175
- Shimojo, M., K. Ikeda, Y. Asano, R. Ishiwaka, T. Shao, H. Sato, M. Tobisa, Y. Nakano, N. Ohba, Y. Yano and Y. Masuda 2003 Exponential functions with base  $e$  in growth analysis and deriving them from rotations of axes of time described using Euler's formula. *J. Fac. Agr., Kyushu Univ.*, **48**: 65-69
- Shimojo, M., K. Ikeda, Y. Asano, R. Ishiwaka, T. Shao, H. Sato, M. Tobisa, Y. Nakano, N. Ohba, Y. Yano and Y. Masuda 2004a Hypothetic incomplete pair disappearances of complex numbers forming ' $0 = (-1) + 1$ ' in the application to complex representation of some aspects of ruminant agriculture. *J.*

- Fac. Agr., Kyushu Univ.*, **49**: 61–67
- Shimojo, M., K. Ikeda, R. Ishiwaka, H. Sato, Y. Asano, M. Tobisa, Y. Nakano, N. Ohba, M. Eguchi and Y. Masuda 2004b Introducing complex numbers into basic growth functions – (II) Applying complex representation of  $'(-1)+1'$  to definite integral of exponential function with base  $e$  expanded into infinite series –. *J. Fac. Agr., Kyushu Univ.*, **49**: 337–341
- Shimojo, M., K. Ikeda, R. Ishiwaka, H. Sato, Y. Asano, M. Tobisa, Y. Nakano, N. Ohba, M. Eguchi and Y. Masuda 2004c Introducing complex numbers into basic growth functions – (III) Symmetry breakdown in complex representation of  $'0=(-1)+1'$  in definite integral of exponential function with base  $e$  expanded into infinite series –. *J. Fac. Agr., Kyushu Univ.*, **49**: 343–348
- Watson, D. J. 1952 The physiological basis of variation in yield. *Adv. Agron.*, **4**: 101–145