

Electro Protoplast Fusion between *Solanum sisymbriifolium* and Other *Solanum* Species

Oda, Naoki
Faculty of Agriculture, Kyushu University

Isshiki, Shiro
Faculty of Agriculture, Kyushu University

Sadohara, Takeshi
Faculty of Agriculture, Kyushu University

Ozaki, Yukio
Faculty of Agriculture, Kyushu University

他

<https://doi.org/10.5109/4712>

出版情報：九州大学大学院農学研究院紀要. 51 (1), pp.67-72, 2006-02-01. 九州大学大学院農学研究院
バージョン：
権利関係：



Electro Protoplast Fusion between *Solanum sisymbriifolium* and Other *Solanum* Species

Naoki ODA¹, Shiro ISSHIKI², Takeshi SADOHARA¹, Yukio OZAKI^{3*}
and Hiroshi OKUBO

Laboratory of Horticultural Science, Division of Agricultural Botany, Department of Plant Resources,
Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan
(Received October 28, 2005 and accepted November 16, 2005)

Symmetric or asymmetric protoplast fusions between *Solanum sisymbriifolium* and other *Solanum* species (*S. integrifolium* and *S. toxicarium*) were performed to obtain the perfectly soil-borne disease resistant rootstocks for eggplant cultivation. Somatic hybrid was not obtained from symmetric fusion in selective media, where one parent's protoplasts formed colonies and the other did not. Combining of iodoacetamide-induced inactivation of *S. integrifolium* and no cell division ability of *S. sisymbriifolium* on a selective medium in asymmetric fusion, the hybridity of two calli was elucidated by analyzing IDH, PGM and SKDH isozymes. Plant regeneration from the somatic hybrid calli was also successful, but the regenerated plants did not show the hybridity. The result may be caused by the elimination of *S. integrifolium* chromosomes during the process of shoot regeneration from calli.

INTRODUCTION

Somatic hybridization is a technique that allows the manipulation of cellular genomes by protoplast fusion. Its major contribution to plant breeding is in overcoming common crossing barriers among plant species and in organelle genetics and breeding.

Solanum sisymbriifolium Lam., a wild American species, shows resistance to root-knot nematodes (Fassuliotis and Bhatt, 1982), carmine spider mites (Schalk *et al.*, 1975) and a few races of bacterial wilt (Ali *et al.*, 1990). In addition, this species is not susceptible to low temperature, and fruit yield of eggplant grafted on this species is higher than that on eggplant (Ali, 1991). It is, however, not easy to prepare this species as a rootstock because of existing spines on its stems and leaf veins.

Solanum integrifolium is a source of genes for resistance to *Fusarium* wilt (Yamakawa and Mochizuki, 1979) and mite *Tetranychus urticae* (Rotino *et al.*, 1992). According to our previous experiment, this species showed rapid growth than other wild *Solanum* species such as *S. indicum* L., *S. torvum* Swartz, *S. sisymbriifolium* and *S. toxicarium*. Although it is also currently used as rootstock in eggplant cultivation in Japan (Okimura *et al.*, 1986), it is susceptible to root-knot nematodes and a few races of bacterial wilt (Mochizuki and Yamakawa, 1979; Ali *et al.*, 1990).

Solanum toxicarium has perfect resistance to all above-mentioned diseases (Yamawaka and Mochizuki, 1979; Ali *et al.*, 1990; Ali, 1991). But in eggplant on this species, the growth is very slow and the fruit quality is very poor. Moreover, this species is susceptible to low temperature (Ali, 1991).

Although interspecific cross seems to be the most appropriate method to combine resistant genes of wild species, three wild *Solanum* species, *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium*, are perfectly cross incompatible to each other (Nishio *et al.*, 1984; Ali and Fujieda, 1990; Ali, 1991).

Establishment of the protoplast culture system of the fusion parents is required for the production of somatic hybrids. Methods for protoplast culture and plant regeneration in *S. integrifolium* and *S. toxicarium* have been established (Sadohara, 1993; Sadohara *et al.*, 1993) and those in *S. sisymbriifolium* (Oda *et al.*, 2006) were reported in the previous paper. This study was undertaken to obtain the perfectly desirable characters, e.g., multiple resistance to all the diseases, spinelessness, tolerance to low temperature and high quality and yield, by using somatic hybridization between *S. sisymbriifolium* and *S. integrifolium* and between *S. sisymbriifolium* and *S. toxicarium*.

MATERIALS AND METHODS

Symmetric protoplast fusion between *S. sisymbriifolium* and other *Solanum* spp.

Plant materials

Seeds of *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium* were surface sterilized with sodium hypochlorite solution (2% active chlorine) for 20 min and washed three times with sterilized water. They were aseptically sown in 200 ml conical flasks containing 40 ml of MS medium (Murashige and Skoog, 1962) supplemented with 1% sucrose and 0.25% gellan gum, and incubated in the growth cabinet (30 μ E/m²·s, 16 h/day at

¹ Laboratory of Horticultural Science, Department of Agronomy, Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan

² Laboratory of Biotechnology and Plant Breeding, Department of Applied Biological Sciences, Faculty of Agriculture, Saga University, Saga 840–8502, Japan

³ Laboratory of Agricultural Ecology, Division of Agricultural Ecology, Department of Plant Resources, Faculty of Agriculture, Kyushu University, Fukuoka 811–2307, Japan

* Corresponding author (E-mail: ozaki@farm.kyushu-u.ac.jp)

Table 1. Components of media used for protoplast culture.

Medium	Mineral salts	Organic components	NAA (mg/l)	2,4-D (mg/l)	BA (mg/l)	Kinetin (mg/l)
A	Kao	Kao	5.0	1.0	1.0	0
B	1/2MS	MS	1.0	0	0	1.0
C	1/2MS	MS	0.1	0	0	0.1

25 °C). Plants of *S. toxicarium* were subcultured by leafy node cutting on the fresh medium of the same composition with three weeks intervals. Leaves of *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium* were taken from 3–4, 2 and 6 week-old explants, respectively.

Protoplast fusion

Protoplasts were isolated as the same method in the previous report (Oda *et al.*, 2006). The isolated protoplasts were suspended in fusion solution containing 0.225 M glucose, 0.225 M mannitol and 2.5 mM CaCl₂. The density was adjusted to 2.0×10^5 protoplasts/ml. The protoplast suspensions of *S. sisymbriifolium* and *S. integrifolium*, or those of *S. sisymbriifolium* and *S. toxicarium* were mixed at a ratio of 1:1 (v/v). The protoplast mixture were fused by means of electro fusion using a Somatic Hybridizer SSH-1 (Shimadzu Corp., Japan). A 0.8 ml aliquot of mixed protoplast suspension was introduced into the fusion chamber FTC-03 (Shimadzu Corp., Japan), aligned with a voltage altering current of 40 V, frequency of 1.0 MHz, voltage direct current of 150 V (750 V/cm field strength), pulse width of 400 μ s and pulse of 1–2 times. Following the fusion treatment, the protoplast suspension was transferred to 60 \times 15 mm petri dish and an equal volume of two-fold strength culture medium was added.

Protoplast culture

Fused protoplasts of *S. sisymbriifolium* and *S. integrifolium* were cultured in Kao medium containing 0.03 M sucrose, 0.4 M mannitol, 5.0 mg/l NAA, 1.0 mg/l 2,4-D and 1.0 mg/l BA (medium A; Table 1) or 1/2 MS medium containing 0.03 M sucrose, 0.4 M mannitol, 1.0 mg/l NAA and 1.0 mg/l kinetin (medium B) (Sadohara, 1993), and those of *S. sisymbriifolium* and *S. toxicarium* were cultured in medium A or a half strength MS medium containing 0.03 M sucrose, 0.4 M mannitol, 0.1 mg/l NAA and 0.1 mg/l kinetin (medium C) (Sadohara, 1993; Sadohara *et al.*, 1993). Protoplasts of both the fusion pairs were cultured in the dark at 30 °C for the first seven days and thereafter placed at 25 °C in the dark. Two ml of fresh media (same as each initial medium but modified with 0.2 M mannitol) were added at an interval of 14 days.

Plant regeneration

Protoplast-derived calli of 1–3 mm in diameter (eight weeks after culture) were transferred to 100 ml conical flasks containing 30 ml of MS medium supplemented with 30 g/l sucrose, 7.0 g/l agar, 0.1 mg/l IAA and 1.0 mg/l zeatin for shoot regeneration. This medium has been shown to be adequate for the regeneration of protoplast-derived calli of *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium* (Sadohara, 1993; Oda *et al.*,

2006). After four weeks of culture, regenerated plants were transferred for rooting on a half strength MS medium with 5.0 g/l sucrose and 2.5 g/l gellan gum.

Isozyme analysis

At the leaf regeneration stage, isozymes of the leaf of each plant were examined by starch gel electrophoresis. The gels were stained for isocitrate dehydrogenase (IDH; EC 1.1.1.41) and phosphoglucose dehydrogenase (PGD; 1.1.1.43) by means of the methods of Wendel and Parks (1982) and Wendel (1983).

Asymmetric protoplast fusion between *S. sisymbriifolium* and other *Solanum* spp.

Plant materials

Plant materials and the methods of protoplast isolation were identical to the previous experiment.

Iodoacetamide treatment

Protoplasts were treated with four levels of concentrations (5.0, 7.5, 10.0 and 12.5 mM) of iodoacetamide (IOA) solution containing 0.4 M mannitol and 0.03 M sucrose (pH 5.5) at 5 °C for 10 min. Then they were washed three times by centrifugation (100 \times g, 3 min) with culture medium. IOA-treated protoplasts of *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium* were cultured in media A, B and C, respectively. Frequency of cell division was examined after ten days of protoplast culture.

Protoplast fusion, culture and plant regeneration

Protoplast fusion and regeneration were carried out by the same methods in the previous experiment. Selection of the fused heterokaryon protoplast from homokaryon or non-fused protoplast was performed by means of the combination of IOA inactivation and the difference in cell division ability in the selective media.

Isozyme analysis

Samples were extracted from calli and leaves of regenerated plants. Isocitrate dehydrogenase (IDH), phosphoglucose mutase (PGM; EC 2.7.5.1) and shikimate dehydrogenase (SKDH; EC 1.1.1.25) isozymes were stained according to the same methods in the previous experiment.

RESULTS

Symmetric protoplast fusion between *S. sisymbriifolium* and other *Solanum* spp.

The process of electro fusion took less than 2 min per dish. Pearl chains of 6–7 protoplasts formed within 50 seconds of an A.C. field application at 1.0 MHz and 750 V/cm. Fusion between protoplasts occurred when D.C. square pulses were applied. The exposure of samples of protoplast mixture to a train of 2–3 D.C. pulses of

Table 2. Effect of media on colony formation from protoplasts of *S. sisymbriifolium* and *S. integrifolium* and from fused protoplast mixture^z.

Species	Medium	Volume of protoplast suspension ^y	No. of colonies
<i>S. sisymbriifolium</i>	A ^x	2.0	7
	B	2.0	0
<i>S. integrifolium</i>	A	2.0	0
	B	2.0	211
Fused protoplast mixture	A	6.0	13
	B	6.0	280

^z Protoplasts obtained after electro protoplast fusion between *S. sisymbriifolium* and *S. integrifolium*.

^y Protoplast density = 5.0×10^4 /ml.

^x See Table 1.

400 μ s, each 750 V/cm, resulted in a fusion rate of 40–50%. At least 20% of the fusion products were in the 1:1 (binary) category. The frequency of heterokaryons was not determined, since it was unable to distinguish protoplasts of one parent from those of the other parent.

Solanum sisymbriifolium (+) *S. integrifolium*

Effect of media on colony formation from protoplasts in *S. sisymbriifolium*, *S. integrifolium* and fused protoplast mixture is shown in Table 2. Protoplast division was found only in *S. integrifolium* in medium B, whereas in medium A, it was observed only in *S. sisymbriifolium*. Colony formation of the fused protoplasts was observed in both media A and B.

The rates of shoot and root formation from the protoplast-derived calli in the fusion plates were similar to those of the control plates of *S. sisymbriifolium* or *S. integrifolium* parental protoplasts (Table 3).

Two hundred thirty eight shoots transferred to rooting medium were used for the analysis of isozyme patterns of IDH. The isozyme pattern of 28 shoots regenerated in medium A was all identical and it was the same as that of *S. sisymbriifolium*. All the regenerated shoots in medium B exhibited the same isozyme pattern as the shoots of *S. integrifolium* (data not shown).

Solanum sisymbriifolium (+) *S. toxicarium*

Cell division and colony formation of the fused protoplasts were observed in both media A and C, whereas those of the both parents were not (Table 4). Compared

Table 4. Effect of media on colony formation from protoplasts of *S. sisymbriifolium* and *S. toxicarium* and from fused protoplast mixture^z.

Species	Medium	Volume of protoplast suspension ^y	No. of colonies
<i>S. sisymbriifolium</i>	A ^x	2.0	6
	C	2.0	0
<i>S. toxicarium</i>	A	2.0	0
	C	2.0	11
Fused protoplast mixture	A	6.0	18
	C	6.0	13

^z Protoplasts obtained after electro protoplast fusion between *S. sisymbriifolium* and *S. toxicarium*.

^y Protoplast density = 5.0×10^4 /ml.

^x See Table 1.

the control plates of *S. sisymbriifolium* or *S. toxicarium* to the fusion plates, the rates of shoot and root formation of the fused protoplast-derived calli were similar to those of the parents (Table 5).

Isozyme pattern of PGD in 37 shoots forming roots was analyzed. Twenty four shoots regenerated from the calli cultured in medium A and 13 shoots in medium C showed the same isozyme patterns of the shoots of *S. sisymbriifolium* and *S. toxicarium*, respectively.

Asymmetric protoplast fusion between *S. sisymbriifolium* and other *Solanum* spp.

Cell division of *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium* protoplasts were completely inhibited by IOA of 7.5, 12.5 and 10.0 mM, respectively (Fig. 1).

The only fusion pair of *S. sisymbriifolium* and IOA treated *S. integrifolium* protoplasts resulted in the formation of five colonies, whereas the other pairs did not. Five calli (Table 6) were transferred to the regeneration medium to reduce shoot formation.

Isozymes for IDH, PGM and SKDH in the parentages of these five calli were analyzed before the shoot regeneration. Two calli were somatic cell hybrids (Fig. 2; data of PGM and SKDH not shown) while the others showed the same isozyme banding patterns as *S. integrifolium*. The shoots regenerated from the two somatic hybrid calli, however, showed one isozyme pattern, and it was

Table 3. Comparison of shoot formation from protoplast-derived calli and rooting of the shoots in *S. sisymbriifolium*, *S. integrifolium* and fused protoplast mixture^z.

Species	Colony-induction medium	No. of calli		No. of shoots	
		Transferred	Forming shoots	Transferred	Forming roots
<i>S. sisymbriifolium</i>	A ^w	52	27 (51.9) ^y	50	43 (86.0) ^x
<i>S. integrifolium</i>	B	52	23 (44.2)	48	39 (81.3)
Fused protoplast mixture	A	13	6 (46.2)	28	20 (71.4)
	B	280	98 (35.0)	210	173 (82.4)

^z Protoplasts obtained after electro protoplast fusion between *S. sisymbriifolium* and *S. integrifolium*.

^y Percent calli forming shoots.

^x Percent shoots forming roots.

^w See Table 1.

Table 5. Comparison of shoot formation from protoplast-derived calli and rooting of the shoots in *S. sisymbriifolium*, *S. toxicarium* and fused protoplast mixture^z


Species	Colony-induction medium	No. of calli		No. of shoots	
		Transferred	Forming shoots	Transferred	Forming roots
<i>S. sisymbriifolium</i>	A ^w	52	25 (48.1) ^y	46	41 (89.1) ^x
<i>S. integrifolium</i>	C	52	16 (30.8)	30	30 (100)
Fused protoplast mixture	A	18	11 (61.1)	29	24 (82.8)
	C	13	5 (38.5)	14	13 (92.9)

^z Protoplasts obtained after electro protoplast fusion between *S. sisymbriifolium* and *S. toxicarium*.

^y Percent calli forming shoots.

^x Percent shoots forming roots.


^w See Table 1.

**Fig. 1.** Effect of IOA on the cell division of *S. sisymbriifolium*, *S. integrifolium*, and *S. toxicarium*.

the same as the pattern of *S. sisymbriifolium* (data not shown). The plants also had the similar morphology as *S. sisymbriifolium*, and died about four weeks after the shoot regeneration.

DISCUSSION

The present results indicate that the fusion rates of

**Fig. 2.** Electrophoretic banding patterns of IDH isozymes, in leaf tissue of *S. sisymbriifolium* (SS), *S. integrifolium* (SI) and their mixture (Mix), and in calli of somatic hybrids (1 and 2). Arrows indicate heterodimer bands.**Table 6.** Effect of media on colony formation from protoplast mixture fused between *S. sisymbriifolium* and *S. integrifolium*, and between *S. sisymbriifolium* and *S. toxicarium*.

Species	Medium	Cell division ^z	No. of colonies
<i>S. sisymbriifolium</i> * + <i>S. integrifolium</i>	A ^y	-	-
<i>S. sisymbriifolium</i> * + <i>S. integrifolium</i> *	B	+(4) ^x	5
<i>S. sisymbriifolium</i> * + <i>S. toxicarium</i>	A	-	-
<i>S. sisymbriifolium</i> * + <i>S. toxicarium</i> *	C	-	-

^z Volume of protoplast suspension = 6.0 ml, protoplast density = 5.0×10^4 /ml, +: Cell division, -: No cell division.

^y See Table 1.

^x Days to first cell division.

* IOA treatment.

40–50% were obtained in the first experiment, and they were similar to those in the fusion between *S. integrifolium* and *S. toxicarium* (Sadohara, 1993). Heterokaryon frequency was not determined since the protoplasts of the fusion partners were of the same type. At least 20% of the fusion products were estimated to be binary fusions. In a previous fusion experiment using the same electric apparatus (unpublished results) with

green mesophyll protoplasts of *S. integrifolium* or *S. toxicarium* and colorless cell suspension protoplasts of *S. sisymbriifolium*, heterokaryons formed one quarter of the fusion products (50% fusion rate overall). At least 10% of the fusion products were, therefore, estimated to be heterokaryon. This frequency was 5–6% higher than that reported by Sadohara (1993) in the electro fusion between *S. integrifolium* and *S. toxicarium* protoplasts.

The selection of somatic hybrids by combination of media has been reported (Carson *et al.*, 1972). In the present experiments, however, the medium for selection of only hybrid could not be found. But we found the media such as A, B and C where the protoplasts of one parent could form colonies and those of the other parent could not. It is unable to determine in these media whether the calli were from one parent or somatic hybrid but is possible to do that they were not from the other parent.

Isozyme banding patterns of the shoots regenerated from the calli cultured in the media A, B and C were identical to those of *S. sisymbriifolium*, *S. integrifolium* and *S. toxicarium*, respectively, in symmetric protoplast fusion treatments. There are two possible reasons that the somatic hybrids could not be obtained. One is that the only one parent protoplasts divided but heterokaryons did not divide or not form colonies after division. The other is that the one parent's chromosome(s) in hybrid cell disappeared during mitosis at cell division or regenerating stage.

IOA treatment of the protoplast of one parental species (*S. integrifolium*) was successfully combined with another untreated parent (*S. sisymbriifolium*) in the medium B. The IOA-treated protoplasts of *S. integrifolium* occasionally divided when they were cocultured with untreated protoplast of *S. sisymbriifolium*, even at the high IOA concentration enough to kill all the *S. integrifolium* protoplasts when cultured alone. This "nurse effect" of untreated protoplasts explains the appearance of an escaped callus of *S. integrifolium* even at the 12.5 mM concentration of IOA (Fig. 1).

Inability of cell division of one parental species has been utilized for selection of somatic hybrids (Kameya *et al.*, 1990). In this study, two calli that formed shoots were somatic hybrids, but the shoots regenerated from them showed the same IDH, PGM and SKDH isozyme patterns and morphology as *S. sisymbriifolium* did. This results may be caused by further elimination of *S. integrifolium* chromosomes in the plants regenerated from the hybrid calli, i.e., *S. integrifolium* chromosomes in hybrid cells disappeared during mitotic division at regenerating stage, because the parents' chromosomes were unbalanced. There are numerous reports that interfamilial or intergeneric somatic hybrids were genetically unstable and morphologically abnormal and showed the elimination of specific chromosome(s) and the disappearance of specific isozyme bands of a parent (Krumbiegel and Schieder, 1979; Gleba and Hoffman, 1979, 1980; Chien *et al.*, 1982; Niizeki *et al.*, 1985; Gleba *et al.*, 1988). It was considered that the similar phenom-

enon occurred in this study.

Although the hybridity of the regenerated plants was not confirmed, somatic hybrid calli were produced. This suggests that the possibility of obtaining somatic hybrids between *S. sisymbriifolium* and other *Solanum* species, particularly *S. integrifolium*, by asymmetric protoplast fusion when subculturing system of asymmetric fused protoplast is established.

ACKNOWLEDGMENTS

The authors wish to express our sincere appreciation to Prof. Emeritus Dr. Kunimitsu Fujieda, Kyushu University for the valuable advice during this experiment.

REFERENCES

- Ali, M. 1991 Breeding Eggplant Rootstock for Multiple Disease Resistance. Ph. D. Thesis, Kyushu University, Fukuoka, Japan
- Ali, M. and K. Fujieda 1990 Cross compatibility between eggplant (*Solanum melongena* L.) and wild relatives. *J. Japan. Soc. Hort. Sci.*, **58**: 977–984
- Ali, M., M. A. Quadir, H. Okubo and K. Fujieda 1990 Resistance of eggplant, its wild relatives and their hybrids to different strains of *Pseudomonas solanacearum*. *Scientia Hort.*, **45**: 1–9
- Carson, P. S., H. H. Smith and R. D. Dearing 1972 Parasexual interspecific plant hybridization. *Proc. Natl. Acad. Sci. USA*, **69**: 2292–2294
- Chien, Y. C., K. N. Kao and L. R. Wetter 1982 Chromosomal and isozyme studies of *Nicotiana tabacum* – *Glycine max* hybrid cell lines. *Theor. Appl. Genet.*, **62**: 301–304
- Fassuliotis, G. and D. P. Bhatt 1982 Potential of tissue culture for breeding root-knot nematode resistance into vegetables. *J. Nematol.*, **14**: 10–14
- Gleba, Y. Y. and F. Hoffmann 1979 "Arabidobrassica": Plant-genome engineering by protoplast fusion. *Naturwissenschaften*, **66**: 547–554
- Gleba, Y. Y. and F. Hoffmann 1980 "Arabidobrassica": A novel plant obtained by protoplast fusion. *Planta*, **149**: 112–117
- Gleba, Y. Y., S. Hinnisdaels, V. A. Sidorov, V. A. Kaleda, A. S. Parokony, N. V. Boryshuk, N. N. Cherep, I. Negrutiu and M. Jacobs 1988 Intergeneric asymmetric hybrids between *Nicotiana plumbaginifolia* and *Atropa belladonna* obtained by "gamma-fusion". *Theor. Appl. Genet.*, **76**: 760–766
- Kameya, T., N. Miyazawa and S. Toki 1990 Production of somatic hybrids between *Solanum melongena* L. and *S. integrifolium* Poir. *Japan. J. Breed.*, **40**: 429–434
- Krumbiegel, G. and O. Schieder 1979 Selection of somatic hybrids after fusion of protoplasts from *Datura innoxia* Mill. and *Atropa belladonna* L. *Planta*, **145**: 371–375
- Mochizuki, H. and H. Yamakawa 1979 Potential utilization of bacterial-wilt resistant *Solanum* species as rootstock for commercial eggplant production. *Bull. Veg. & Orn. Crops Res. Stn. Japan*, **A6**: 11–18 (In Japanese with English summary)
- Murashige, T. and F. Skoog 1962 A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.*, **15**: 473–497
- Niizeki, M., M. Tanaka, S. Akada, A. Hirai and K. Saito 1985 Callus formation of somatic hybrid of rice and soybean and characteristics of the hybrid callus. *Japan. J. Genet.*, **60**: 81–92
- Nishio, T., H. Mochizuki and K. Yamakawa 1984 Interspecific cross of eggplants and related species. *Bull. Veg. & Orn. Crops Res. Stn. Japan*, **A12**: 57–64 (In Japanese with English summary)

- Oda, N., S. Isshiki, T. Sadohara, Y. Ozaki and H. Okubo 2006 Establishment of protoplast culture of *Solanum sisymbriifolium*. *J. Fac. Agr., Kyushu Univ.*, **51**(1): 63–66. (In press)
- Okimura, M., S. Matsuo, K. Arai and S. Okitsu 1986 Influence of soil temperature on the growth of fruit vegetable grafted on different stocks. *Bull. Veg. & Ornam. Crops Res. Stn. Japan*, **C9**: 43–58 (In Japanese with English summary)
- Rotino G. L., D. Perrone, P. Ajmone-Marsan and E. Lupotto 1992 Transformation of *Solanum integrifolium* Poir via *Agrobacterium tumefaciens*: Plant regeneration and progeny analysis. *Plant Cell Rep.*, **11**: 11–15
- Sadohara, T. 1993 Studies on the breeding of *Solanum* species by protoplast fusion. Thesis of the Master Degree, Kyushu Univ., Fukuoka, Japan (In Japanese)
- Sadohara, T., H. Okubo and K. Fujieda 1993 Plant regeneration from mesophyll protoplasts of *Solanum toxicarium*. *Sci. Bull. Fac. Agr., Kyushu Univ.*, **48**: 13–19 (In Japanese with English summary)
- Schalk, J. M., A. K. Stoner, R. E. Webb and H. F. Winters 1975 Resistance in eggplant, *Solanum melongena* L. and nontuber-bearing *Solanum* species to carmin spider mite. *J. Amer. Soc. Hort. Sci.*, **100**: 479–481
- Wendel, J. F. 1983 Electrophoretic analysis of genetic variation in wild and cultivated *Camellia japonica* L. Ph. D. Thesis, University of North Carolina, Chapel Hill, North Carolina
- Wendel, J. F. and C. R. Parks 1982 Genetic control of isozyme variation in *Camellia japonica* L. *J. Hered.*, **73**: 197–204
- Yamakawa, K. and H. Mochizuki 1979 Nature and inheritance of Fusarium-wilt resistance in eggplant cultivars and related wild *Solanum* species. *Bull. Veg. & Ornam. Crops Res. Stn. Japan*, **A6**: 19–27 (In Japanese with English summary)