Enhancement of lepton flavor violation in a model with bimaximal mixing at the grand unification scale

Kanemura, Shinya Department of Physics, Osaka University

Matsuda, Koichi Department of Physics, Osaka University

Ota, Toshihiko Department of Physics, Osaka University

Shindou, Tetsuo Scuola Internazionale Superiore di Studi Avanzati

他

https://hdl.handle.net/2324/4795141

出版情報:PHYSICAL REVIEW D. 72, pp.093004-, 2005-11-11. American Physical Society バージョン: 権利関係:

Enhancement of lepton flavor violation in a model with bi-maximal mixing at the grand unification scale

Shinya Kanemura, $^{1,\,*}$ Koichi Matsuda, $^{1,\,\dagger}$ Toshihiko Ota, $^{1,\,\ddagger}$

Tetsuo Shindou,^{2, 3, §} Eiichi Takasugi,^{1, ¶} and Koji Tsumura^{1, **}

¹Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan

²Scuola Internazionale Superiore di Studi Avanzati, I-34014, Trieste, Italy ³Theory group, KEK, Tsukuba 305-0801, Japan

Abstract

We study phenomenological predictions in the scenario with the quasi-degenerate relation among neutrino Dirac masses, $m_{D1} \simeq m_{D2} < m_{D3}$, assuming the bi-maximal mixing at the grand unification scale in supersymmetric standard models with right-handed neutrinos. A sufficient lepton number asymmetry can be produced for successful leptogenesis. The lepton flavor violating process $\mu \rightarrow e\gamma$ can be enhanced due to the Majorana phase, so that it can be detectable at forthcoming experiments. The processes $\tau \rightarrow e\gamma$ and $\tau \rightarrow \mu\gamma$ are suppressed because of the structure of neutrino Dirac masses, and their branching ratios are smaller than that of $\mu \rightarrow e\gamma$.

PACS numbers: 11.30.Hv, 14.60.Pq, 14.60.St

Keywords: Neutrino mixing, Lepton flavor violation, Leptogenesis

^{*}Electronic address: kanemu@het.phys.sci.osaka-u.ac.jp

 $^{^{\}dagger} Electronic address: matsuda@het.phys.sci.osaka-u.ac.jp$

[‡]Electronic address: toshi@het.phys.sci.osaka-u.ac.jp

[§]Electronic address: shindou@sissa.it

 $[\]P Electronic address: takasugi@het.phys.sci.osaka-u.ac.jp$

^{**}Electronic address: ko2@het.phys.sci.osaka-u.ac.jp

1. INTRODUCTION

Neutrinos can be useful as a probe of physics at high energies such as grand unified theories (GUT). Supersymmetry (SUSY) may be introduced to avoid problems due to large hierarchy between the weak scale and the GUT scale. Tiny neutrino masses and observed mixing angles may be explained by assuming the existence of right-handed neutrinos with large Majorana masses[1]. They are determined from the high energy structure of the model by using renormalization group equations (RGEs). The resulting mass spectrum and mixing angles depend on the Majorana mass matrix of right-handed neutrinos and the neutrino Yukawa interaction. It would be possible to consider phenomenology of the model by putting additional assumptions in the high-energy structure of neutrino sector.

In this paper, we consider the minimal supersymmetric standard model with right-handed neutrinos (MSSMRN), in which the bi-maximal solution for the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix^[2] is assumed to be realized at the GUT scale. This solution is predicted in several GUT models^[3]. In the model with the bi-maximal mixing solution, the 1-3 element of the PMNS matrix is zero at the GUT scale, while there are two Majorana CP phases [4]. A non-zero value of the 1-3 element with the CP violating Dirac phase can be induced at the low energy through RGEs[5, 6]. The observed value θ_{\odot} (tan² $\theta_{\odot} \simeq 0.4[7, 8, 9]$) for the solar neutrino angle at low energy is clearly different from the maximal mixing $\pi/4$ at the GUT scale. This difference can be explained by taking into account the running effect due to the neutrino Yukawa couplings between the two scales [5, 6, 10, 11]. When masses of neutrinos corresponding to the solar neutrino data are relatively larger such as 0.05eV, the running effect becomes significant so that the value of θ_{\odot} can be reproduced at the low energy scale[6, 10]. On the other hand, when the mass scale of neutrino is larger than 0.15eV, the atmospheric neutrino mixing angle is so instable that the bi-maximal mixing model cannot explain the experimental result of atmospheric neutrino oscillation [6, 10]. Therefore, we here consider the case in which masses of neutrinos are in the range between 0.05eV and 0.15 eV. Then the solar neutrino data prefer two cases for the pattern of the eigenvalues of the neutrino Dirac mass matrix; (i) hierarchical case $(m_{D1} < m_{D2} < m_{D3})$ and (ii) quasidegenerate case $(m_{D1} \simeq m_{D2} < m_{D3})$. The case (i) has been studied in Ref. [12], and it has been found that lepton flavor violating processes are not significant. In the present paper, we study the case (ii) and investigate its low energy phenomenology.

We shall show that our scenario is compatible with the low energy neutrino data, and that sufficient amount of lepton number asymmetry can be produced for successful leptogenesis[13, 14, 15, 16, 17]. Furthermore, we find that the lepton flavor violating process $\mu \to e\gamma$ can be enhanced by the Majorana phase effect to be as large as the experimental reach at MEG[18]. We also find that the branching ratios of $\tau \to e\gamma$ and $\tau \to \mu\gamma$ are smaller than that of $\mu \to e\gamma$. These are striking features of the quasi-degenerate scenario.

In Sec.2, the quasi-degenerate scenario with the bi-maximal mixing solution is defined in the MSSMRN. In Sec.3, we discuss phenomenological results of our scenario, especially on leptogenesis and lepton flavor violation. Comments and conclusions are given in Sec.4. Some derivations are given in Appendices.

2. THE QUASI-DEGENERATE SCENARIO

We consider the neutrino Yukawa couplings in the quasi-degenerate scenario in the MSSMRN. The Lagrangian relevant to right-handed neutrinos is given by

$$\mathcal{L}_{Y+M} = \overline{N}_R \phi_u^0 Y_\nu \nu_L - \frac{1}{2} \overline{N}_R^c M_R N_R + \text{h.c.}, \qquad (1)$$

where N_R is the right-handed neutrino with the 3 × 3 Majorana mass matrix M_R , ν_L is the left-handed neutrino, ϕ_u^0 (ϕ_d^0) is the neutral component of the Higgs doublet with the hypercharge -1/2 (+1/2), and Y_{ν} is the 3 × 3 Yukawa matrix for the neutrinos. The lefthanded neutrino mass matrix is expressed at each scale as[1]

$$m_{\nu} = \frac{v^2 \sin^2 \beta}{2} Y_{\nu}^T M_R^{-1} Y_{\nu}, \qquad (2)$$

where the vacuum expectation values $\langle \phi_u^0 \rangle$ and $\langle \phi_d^0 \rangle$ satisfy $v = \sqrt{2}\sqrt{\langle \phi_u^0 \rangle^2 + \langle \phi_d^0 \rangle^2} \simeq 246$ GeV and $\tan \beta = \langle \phi_u^0 \rangle / \langle \phi_d^0 \rangle$.

Let us consider the neutrino mass matrix at the GUT scale, M_X , which is much higher than that of the Majorana masses of right-handed neutrinos. We take the basis such that the mass matrix of right-handed neutrinos is diagonal as $M_R = D_R \equiv \text{diag}(M_1, M_2, M_3)$, where M_i are real positive eigenvalues ($M_1 \leq M_2 \leq M_3$), and that the mass matrix of the charged leptons is also diagonal. The neutrino Dirac mass matrix m_D is diagonalized as

$$m_D \equiv Y_\nu \frac{v \sin \beta}{\sqrt{2}} = V_R^{\dagger} D_D V_L , \qquad (3)$$

where D_D is a diagonal matrix $D_D \equiv \text{diag}(m_{D1}, m_{D2}, m_{D3})$ with real positive eigenvalues m_{Di} $(m_{D1} \leq m_{D2} \leq m_{D3})$, and V_R and V_L are unitary matrices. As an important assumption of our model, we suppose that the neutrino mass matrix satisfies the bi-maximal mixing solution at M_X ; i.e.,

$$m_{\nu}(M_X) = O_B D_{\nu} O_B^T , \qquad (4)$$

where O_B is given by

$$O_B \equiv \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0\\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{\sqrt{2}}\\ \frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix} , \qquad (5)$$

and D_{ν} is a diagonal matrix,

$$D_{\nu} \equiv \operatorname{diag}(m_1, m_2 e^{i\alpha_0}, m_3 e^{i\beta_0}), \tag{6}$$

with α_0 and β_0 being the Majorana phases and m_i being real positive[4].

It is known that when the scale of neutrino masses are so large as $0.05 \text{eV} < m_1 \sim m_2 \equiv m$, the running effect on the neutrino mass matrix between the weak scale m_Z and M_X becomes large due to the neutrino Yukawa interaction[6, 10, 19]. The 1-3 element V_{13} of the PMNS matrix is also induced at the low energy scale, which is found to be proportional to m_1m_3 [6]. The element $|V_{13}|$ can be sizable when both m_1 and m_3 are sufficiently large¹. In order to reproduce the solar neutrino data from the bi-maximal solution at M_X with $Y_{\nu}^{\dagger}Y_{\nu}$ to be diagonal, there are two possibilities for the pattern of the neutrino Dirac masses, i.e., hierarchical case $m_{D1} < m_{D2} < m_{D3}$ and quasi-degenerate case $m_{D1} \simeq m_{D2} < m_{D3}$ with

$$V_L = P_{ex} \equiv \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ -1 & 0 & 0 \end{pmatrix} .$$
 (7)

The detailed discussion appears in Appendix A. In this paper, we concentrate on the latter case, namely the quasi-degenerate case. The hierarchical case has been studied in Ref. [12].

¹ There is also a chance to appear large running effect in the case where the neutrino mass spectrum is inverse hierarchical, i.e., $m_3 \ll m_1 < m_2$. In this case, different prediction for the 1-3 element of the PMNS matrix is obtained at the low energy scale.

From Eqs. (2), (3), and (4), we obtain

$$\widetilde{M}_{R}^{-1} \equiv (V_{R}^{*} D_{R}^{-1} V_{R}^{\dagger}) = D_{D}^{-1} (P_{ex} O_{B}) D_{\nu} (P_{ex} O_{B})^{T} D_{D}^{-1} .$$
(8)

The unitary matrix V_R as well as the eigenvalues M_i are obtained by diagonalizing \widetilde{M}_R^{-1} . Consequently, we find

$$V_R = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \cos\frac{\zeta}{2} & -\frac{i}{\sqrt{2}} \sin\frac{\zeta}{2} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \cos\frac{\zeta}{2} & -\frac{i}{\sqrt{2}} \sin\frac{\zeta}{2} \\ 0 & -i \sin\frac{\zeta}{2} & \cos\frac{\zeta}{2} \end{pmatrix} \begin{pmatrix} e^{-\frac{i}{2}\beta_0} & & \\ & e^{-\frac{i}{4}\alpha_0} & \\ & & e^{-\frac{i}{4}\alpha_0} \end{pmatrix} ,$$
(9)

where

$$\tan \zeta \equiv \frac{2r}{1+r^2} \tan \frac{\alpha_0}{2}, \quad \text{with} \quad r \equiv m_{D1}/m_{D3} < 1.$$
(10)

As seen in Eq. (9), some off-diagonal elements of V_R are of order one. This is the striking feature of the quasi-degenerate case in contrast with the hierarchical case in which V_R is approximately the unit matrix[12]. For the masses of the right-handed neutrinos, we obtain

$$M_{1} = \frac{m_{D1}^{2}}{m} ,$$

$$M_{2} = \frac{m_{D1}^{2}}{m} \frac{2}{\sqrt{[1-r^{2}]^{2} \cos^{2} \frac{\alpha_{0}}{2} + 4r^{2} + [1-r^{2}] \cos \frac{\alpha_{0}}{2}}},$$

$$M_{3} = \frac{m_{D1}^{2}}{m} \frac{2}{\sqrt{[1-r^{2}]^{2} \cos^{2} \frac{\alpha_{0}}{2} + 4r^{2} - [1-r^{2}] \cos \frac{\alpha_{0}}{2}}}.$$
(11)

The derivation of Eqs. (9) and (11) is shown in Appendix B.

3. THE PHENOMENOLOGY

In this section, phenomenological consequences of our scenario are studied. The parameters of the neutrino sector at M_X are related to the low energy observables by the RGEs (see Appendix A). They are constrained from the data of solar and atmospheric neutrino experiments. We here take the values $\Delta m_{\odot}^2 \equiv 8.3 \times 10^{-5} \text{ eV}^2$ and $\tan^2 \theta_{\odot} \equiv 0.4$ as the solar neutrino results[9], and $\Delta m_{\text{atm}}^2 \equiv 2.5 \times 10^{-3} \text{ eV}^2$ and $\sin^2 2\theta_{\text{atm}} \equiv 1.0$ as the atmospheric neutrino results[20]. Throughout this paper, M_X is assumed to be 2×10^{16} GeV. In the following, after the discussion on basic properties, we examine the consistency with leptogenesis, and predict lepton flavor violation (LFV).



FIG. 1: (a) The ratio r as a function of α_0 and m_{D1} for $\tan \beta = 5$ and m = 0.1 eV. The shaded region corresponds to $\cos 2\theta_{\odot} \leq |\cos(\alpha_0/2)| \leq 1$. (b) The eigenvalues M_i as a function of $|\cos(\alpha_0/2)|$ for $m_{D1} = 10$ GeV (black curves) and $m_{D1} = 50$ GeV (gray curves) for $\tan \beta = 5$ and m = 0.1 eV.

The experimental value θ_{\odot} can be reproduced in our scenario with the bi-maximal solution at M_X . As we take the degenerate mass of neutrinos, the running effect on the solar neutrino angle can be large between M_X and m_Z . On the other hand, the angle θ_{atm} can be explained by assuming the masses of neutrinos are less than 0.15 eV so that the running effect is small[6, 10]. In our scenario, the running effects on the neutrino mass matrix, which are parametrized by ϵ_e and ϵ_{τ} , can be expressed as

$$\epsilon_e = -\frac{1}{8\pi^2} \frac{m_{D1}^2}{(v\sin\beta)^2} \left(\frac{1}{r^2} - 1\right) \ln\frac{\overline{M}_R}{M_X} ,$$

$$\epsilon_\tau \simeq -\frac{1}{8\pi^2} \frac{m_\tau^2}{(v\cos\beta)^2} \ln\frac{m_Z}{M_X} .$$
(12)

where \overline{M}_R is the typical mass scale for right-handed neutrinos and m_{τ} is the mass of the tau lepton: see Appendix A. From Eqs. (12) and (A.4) with the experimental data for angles and mass differences, we obtain the relation among α_0 , m_{D1} and r. In Fig. 1-(a), we show the ratio r as a function of m_{D1} for each value of α_0 in the case of m = 0.1 eV and $\tan \beta = 5$. We find that r is insensitive to α_0 . In Fig. 1-(b), M_i are shown as a function of $|\cos(\alpha_0/2)|$ for $m_{D1} = 10$ GeV and $m_{D1} = 50$ GeV, which are determined by m, α_0 , and r through Eq. (11). Notice that M_1 and M_2 are coincident when $\alpha_0 \to 0$ up to $\mathcal{O}(\Delta m_{\text{atm}}^2/m^2)$. The scale \overline{M}_R takes the value between 10^{12} and 10^{14} GeV.

A. Leptogenesis

In models with the heavy Majorana neutrinos, it is possible to consider leptogenesis[13] in order to explain a baryon asymmetry of the universe. In leptogenesis, the out of equilibrium decays of heavy Majorana neutrinos produce a lepton number asymmetry which is converted to the baryon number asymmetry through the sphaleron processes. It is known that leptogenesis is successful to explain the baryon number of the universe when $m_3 < 0.15$ eV and $M_1 > 2 \times 10^7 \text{GeV}[14]$, and mass parameters of our model can be in this allowed range. On the other hand, a constraint from gravitino overproduction can be a serious problem for our model. In order to allow the reheating temperature $T_R > 10^{12} \text{GeV}$ which is required for case $M_1 > 10^{12} \text{GeV}$, the gravitino mass should be heavier than $10 \text{TeV}[21]^2$.

The lepton number asymmetry is produced in the decay of heavy Majorana neutrinos, which can be expressed as[13]

$$\epsilon_i \simeq -\frac{1}{8\pi} \sum_{k \neq i} f\left(\frac{M_k^2}{M_i^2}\right) \frac{\operatorname{Im}[(Y_\nu Y_\nu^\dagger)_{ik}^2]}{(Y_\nu Y_\nu^\dagger)_{ii}},\tag{13}$$

where f(x) is given in the MSSMRN as

$$f(x) = \sqrt{x} \left[\frac{2}{1-x} + \ln\left(\frac{1+x}{x}\right) \right] . \tag{14}$$

In our scenario, $(Y_{\nu}Y_{\nu}^{\dagger})_{ij}$ are calculated at the leading order as

$$(Y_{\nu}Y_{\nu}^{\dagger})_{ij} = \frac{2}{v^{2}\sin^{2}\beta} (V_{R}^{\dagger}D_{D}^{2}V_{R})_{ij}$$

$$\simeq \frac{2m_{D1}^{2}}{v^{2}\sin^{2}\beta} \times$$

$$\begin{pmatrix} 1 & -\frac{1}{2}\delta_{1}\cos\frac{\zeta}{2}e^{\frac{i}{4}(2\beta_{0}-\alpha_{0})} & \frac{i}{2}\delta_{1}\sin\frac{\zeta}{2}e^{\frac{i}{4}(2\beta_{0}-\alpha_{0})} \\ -\frac{1}{2}\delta_{1}\cos\frac{\zeta}{2}e^{-\frac{i}{4}(2\beta_{0}-\alpha_{0})} & \frac{1}{2}(1+1/r^{2}+(1-1/r^{2})\cos\zeta) & \frac{-i}{2}(1-1/r^{2})\sin\zeta \\ \frac{-i}{2}\delta_{1}\sin\frac{\zeta}{2}e^{-\frac{i}{4}(2\beta_{0}-\alpha_{0})} & \frac{i}{2}(1-1/r^{2})\sin\zeta & \frac{1}{2}(1+1/r^{2}-(1-1/r^{2})\cos\zeta) \end{pmatrix}, \quad (15)$$

where $\delta_1 \equiv m_{D2}^2/m_{D1}^2 - 1 \; (\ll 1)$. The asymmetry ϵ_3 is negligibly smaller than $\epsilon_{1,2}$ because of $M_1 \sim M_2 \ll M_3$.³ In the same reason, the elements including $(Y_{\nu}Y_{\nu}^{\dagger})_{12,21}$ are dominant in

² In various SUSY models such as the minimal supergravity model, the gravitino mass are related to the soft SUSY parameters. In our paper, we don't touch origin of soft SUSY breaking terms and we can take the gravitino mass as an independent parameter[22], though assume the universal soft SUSY parameters.

³ In the limit of $x \to 1$, there is an enhancement effect in the lepton number asymmetry[15, 16]. The enhancement is smeared by the following two reasons; i.e., (i) the effect of the decay width for N_i and (ii) the effect of the small mass difference between M_1 and $M_2 \sim \mathcal{O}(\Delta m_{\rm atm}^2/m^2)$.



FIG. 2: The baryon number asymmetry η_{B_0} as a function of $|\cos(\alpha_0/2)|$ for $\beta_0 = 0$, $\pi/2$ and π . The other parameters are assumed to be m = 0.1 eV, $m_{D1} = 50$ GeV, $\delta_1 = 0.05$ and $\tan \beta = 5$.

 $\epsilon_{1,2}$. Therefore $\epsilon_{1,2}$ are approximately proportional to δ_1^2 , which means that finite $\epsilon_{1,2}$ appear only when there is a deviation from the exact mass degeneracy $M_1 = M_2$.

In leptogenesis, the baryon number asymmetry η_{B_0} of the Universe is explained by using the lepton number asymmetry[17];

$$\eta_{B_0} \simeq -10^{-2} \kappa_0 \sum_i \epsilon_i , \qquad (16)$$

where $\kappa_0 \simeq 0.3/\{K(\ln K)^{3/5}\}$ with $K \simeq 170(m/[\text{eV}])$. The numerical result of η_{B_0} is given in Fig. 2. The experimental value $\eta_{B_0} \sim 6.5 \times 10^{-10}$ [23] can be realized in our scenario.

B. LFV processes

In the MSSMRN, the slepton mixing can be a source of LFV[24, 25]. Assuming the universal soft-breaking parameters at M_X , mixing among left-handed sleptons is induced by the renormalization group effects due to neutrino Yukawa couplings between M_X and \overline{M}_R ,

even when there is no mixing at M_X . The induced off-diagonal elements of the slepton mass matrix are approximately expressed as [24, 25]

$$(m_{\tilde{L}}^2)_{ij} \simeq \frac{6m_0^2 + 2|A_0|^2}{16\pi^2} (Y_{\nu}^{\dagger} \Omega Y_{\nu})_{ij} \quad (i \neq j) , \qquad (17)$$

where m_0 and A_0 are the universal SUSY breaking parameters, and

$$\Omega \equiv \operatorname{diag}\left(\ln\frac{M_1}{M_X}, \ln\frac{M_2}{M_X}, \ln\frac{M_3}{M_X}\right).$$
(18)

These off-diagonal elements contribute to LFV processes such as $\ell_i \to \ell_j \gamma$ $(i \neq j)$. The decay widths are given by

$$\Gamma(\ell_i \to \ell_j \gamma) \simeq \frac{\alpha^3 m_{\ell_i}^5}{192\pi^3} \frac{|(m_{\widetilde{L}}^2)_{ij}|^2}{m_S^8} \tan^2 \beta, \tag{19}$$

where α is the fine structure constant, and m_S denotes the typical mass scale of SUSY particles. In the case of universal soft terms, m_S is approximately evaluated in Ref. [26].

Let us consider Eq. (17) in our scenario. We can express Ω as

$$Y_{\nu}^{\dagger}\Omega Y_{\nu} = Y_{\nu}^{\dagger} \left\{ \ln \frac{M_X}{M_3} \mathbf{1} - \operatorname{diag}\left(\ln \frac{M_1}{M_3}, \ln \frac{M_2}{M_3}, 0 \right) \right\} Y_{\nu} , \qquad (20)$$

where the second term of RHS in Eq. (20) corresponds to the threshold effect of right-handed neutrinos. It has been often considered the case in which the first term gives dominant contributions to $\ell_i \to \ell_j \gamma$ processes. However, $Y_{\nu}^{\dagger} Y_{\nu}$ is diagonal at M_X so that the first term does not contribute to LFV. Therefore, remaining sources for LFV are in the second term of Eq. (20). The off-diagonal elements of $(Y_{\nu}^{\dagger} \Omega Y_{\nu})_{ij}$ $(i \neq j)$ are found to be

$$\left| (Y_{\nu}^{\dagger} \Omega Y_{\nu})_{12} \right| = \frac{m_{D2} m_{D3}}{\sqrt{2} v^2 \sin^2 \beta} \sin 2\zeta \ln \frac{M_2}{M_3} , \qquad (21)$$

$$\left| (Y_{\nu}^{\dagger} \Omega Y_{\nu})_{13} \right| = \frac{m_{D1} m_{D3}}{\sqrt{2} v^2 \sin^2 \beta} \sin 2\zeta \ln \frac{M_2}{M_3} , \qquad (22)$$

$$\left| (Y_{\nu}^{\dagger} \Omega Y_{\nu})_{23} \right| = \frac{m_{D1} m_{D2}}{v^2 \sin^2 \beta} \left(\ln \frac{M_1}{M_3} - \cos^2 \frac{\zeta}{2} \ln \frac{M_2}{M_3} \right) .$$
 (23)

The branching ratio of $\tau \to \mu \gamma$ is found to be smaller than those of $\mu \to e\gamma$ and $\tau \to e\gamma$ by a factor of r. All the processes $\ell_i \to \ell_j \gamma$ are maximally suppressed for $|\cos(\alpha_0/2)| \to 1$ because of $M_1 \simeq M_2$. From $m_{D1} \simeq m_{D2}$, we have the relation among the branching ratios of the LFV processes as

$$Br(\tau \to e\gamma) \simeq Br(\tau \to \bar{\nu}_e \nu_\tau e) Br(\mu \to e\gamma) , \qquad (24)$$

where Eqs. (19), (21) and (22) are used. In Fig. 3, we show $Br(\mu \to e\gamma)$ as a function of $|\cos(\alpha_0/2)|$. We find that the value can reach 10^{-12} for the smallest value of $|\cos(\alpha_0/2)|$ under $\cos 2\theta_{\odot} \leq |\cos(\alpha_0/2)| \leq 1$ and for $m_{D1} \gtrsim 50$ GeV. In the case of quasi-degenerate light neutrinos, it is known that the lepton flavor violation processes are strongly suppressed with trivial right-handed mixings, *i.e.* the degenerate heavy neutrino mass spectrum and real mixings among right-handed neutrinos [25]. However, Pascoli et al. showed the possibility of enhancement of the lepton flavor violation processes due to the existence of new CP phases in right-handed mixing even for the case where both light and heavy neutrinos are quasidegenerate [27]. In our model, we don't introduce any CP phases other than α and β , but the lepton flavor violation processes are nevertheless enhanced because of the threshold effects of heavy neutrinos. Therefore we expect that current[18] and future experiments can test our scenario through the LFV measurement for $\mu \to e\gamma$. As compared to the hierarchical case, larger branching ratios for the LFV processes can be obtained in the quasi-degenerate case because of the mixing among right-handed neutrinos. In Fig. 4, we show the ratio of $Br(\tau \to \mu \gamma)$ to $Br(\mu \to e\gamma)$ as a function of $|\cos(\alpha_0/2)|$. In a wide range of the parameter space, $Br(\tau \to \mu \gamma)$ is smaller than $Br(\mu \to e\gamma)$. This is a striking feature of our scenario.

4. CONCLUSION

We have studied the quasi-degenerate scenario in the model in which the bi-maximal mixing solution is realized at M_X in the MSSMRN. By using the low energy neutrino data, we have shown that our result is consistent with the WMAP data assuming leptogenesis. Furthermore, it has been found that the LFV process $\mu \to e\gamma$ can be large enough to be detected at forthcoming experiments such as MEG. We also have found that the branching ratios of $\tau \to e\gamma$ and $\tau \to \mu\gamma$ are smaller than that of $\mu \to e\gamma$. The prediction of sizable LFV processes is a discriminative feature of the quasi-degenerate scenario as compared to the hierarchical one.

It should be noted that our results strongly depend on the Majorana phases α_0 and β_0 . In particular, the processes $\ell_i \to \ell_j \gamma$ become more enhanced for smaller values of $|\cos(\alpha_0/2)|$. We also note that the assumption of diagonal $Y^{\dagger}_{\nu}Y_{\nu}$ is crucial for our results in the quasi-degenerate scenario. The tau associated LFV processes might become significant when a non-zero 2-3 element of $Y^{\dagger}_{\nu}Y_{\nu}$ would be taken into account, by relaxing the



FIG. 3: The branching ratio of $\mu \to e\gamma$ as a function of $|\cos(\alpha_0/2)|$ for $m_{D1} = 10, 50$ and 100 GeV. The SUSY parameters are taken to be $\tan \beta = 5, m_0 = 200$ GeV, $A_0 = 100$ GeV and $m_S = 200$ GeV.

assumption of diagonal $Y^{\dagger}_{\nu}Y_{\nu}$.

Acknowledgments

This work is supported in part by Japan Society for Promotion of Science (Nos. 15-03693, 15-03700 and 15-03927), and also by the Japanese Grant-in-Aid for Scientific Research of Ministry of Education, Science, Sports and Culture, No. 12047218.

APPENDIX A: RGE ANALYSIS

The RGE for the neutrino mass matrix is given by

$$\frac{\mathrm{d}m_{\nu}}{\mathrm{d}\ln\mu} = \frac{1}{16\pi^2} \left\{ [(Y_{\nu}^{\dagger}Y_{\nu})^T + (Y_e^{\dagger}Y_e)^T]m_{\nu} + m_{\nu}[(Y_{\nu}^{\dagger}Y_{\nu}) + (Y_e^{\dagger}Y_e)] \right\},\tag{A.1}$$



FIG. 4: The ratio of $Br(\tau \to \mu \gamma)$ to $Br(\mu \to e\gamma)$ as a function of $|\cos(\alpha_0/2)|$ for $m_{D1} = 10$, 50 and 100 GeV. We take $\tan \beta = 5$.

at the scale between M_X and the typical mass scale of right-handed neutrinos \overline{M}_R aside from the terms proportional to the unit matrix[28]. In the region below \overline{M}_R , we use Eq. (A.1) but without the terms for $Y_{\nu}^{\dagger}Y_{\nu}$. The Yukawa matrix for charged leptons is given by $Y_e \simeq \text{diag}(0, 0, y_{\tau})$ with $y_{\tau} = \sqrt{2}m_{\tau}/(v\cos\beta)$. We concentrate on the case where $Y_{\nu}^{\dagger}Y_{\nu} = \text{diag}(y_1^2, y_2^2, y_3^2)$ for simplicity. We neglect the case in which the 2-3 element may affect on the physics as discussed in Ref. [29]. The solution of Eq. (A.1) can be expressed as[6, 19]

$$m_{\nu}(m_Z) \simeq m_{\nu}(M_X) + K m_{\nu}(M_X) + m_{\nu}(M_X) K$$
, (A.2)

where $K = \text{diag}(\epsilon_e, 0, \epsilon_{\tau})$. The ϵ_e and ϵ_{τ} are given by [12]

$$\epsilon_{e} = \frac{y_{1}^{2} - y_{2}^{2}}{16\pi^{2}} \ln \frac{M_{X}}{\overline{M}_{R}} ,$$

$$\epsilon_{\tau} = \frac{y_{3}^{2} - y_{2}^{2}}{16\pi^{2}} \ln \frac{M_{X}}{\overline{M}_{R}} + \frac{y_{\tau}^{2}}{16\pi^{2}} \ln \frac{M_{X}}{m_{Z}} .$$
(A.3)

The experimental value θ_{\odot} can be reproduced from the bi-maximal mixing solution at M_X by the running effects. The solar mixing angle θ_{\odot} is given in terms of $\epsilon_{e,\tau}$, α_0 and m_1

by

$$\tan^2 \theta_{\odot} = \frac{1 + 2(\epsilon_{\tau} - 2\epsilon_e) \cos^2(\alpha_0/2) (m_1^2 / \Delta m_{\odot}^2)}{1 - 2(\epsilon_{\tau} - 2\epsilon_e) \cos^2(\alpha_0/2) (m_1^2 / \Delta m_{\odot}^2)},$$
(A.4)

where Δm_{\odot}^2 is the experimental value for the mass-squared difference of solar neutrinos. From Eq. (A.4) the allowed region of α_0 is obtained; $\cos 2\theta_{\odot} \leq |\cos(\alpha_0/2)|$. In addition, the following condition among the Yukawa coupling constants is found;

$$2y_1^2 > y_2^2 + y_3^2 + y_\tau^2. (A.5)$$

There are two possibilities under the condition Eq. (A.5) for the pattern of neutrino Yukawa couplings, i.e., the hierarchical case $(y_3^2 < y_2^2 < y_1^2)$ and the quasi-degenerate case $(y_3^2 \simeq y_2^2 < y_1^2)$. As the order of m_{Di} is defined as $m_{D1} \leq m_{D2} \leq m_{D3}$, y_i (i = 1, 2, 3) are assigned as

$$y_1 = \frac{\sqrt{2}m_{D3}}{v\sin\beta}, \quad y_2 = \frac{\sqrt{2}m_{D2}}{v\sin\beta}, \quad y_3 = \frac{\sqrt{2}m_{D1}}{v\sin\beta}, \quad (A.6)$$

which correspond that V_L in Eq. (3) is given by P_{ex} in Eq. (7).

APPENDIX B: DERIVATIONS FOR V_R **AND** M_i

We rotate \widetilde{M}_R^{-1} in Eq. (8) by $P_{ex}O_B$, and find

$$(P_{ex}O_B)^T \widetilde{M}_R^{-1} P_{ex}O_B \simeq \frac{m}{4m_{D1}^2} \begin{pmatrix} [(1+r)^2 + (1-r)^2 e^{i\alpha_0}] & 2(1-r^2)\cos\frac{\alpha_0}{2}e^{i\frac{\alpha_0}{2}} & 0\\ 2(1-r^2)\cos\frac{\alpha_0}{2}e^{i\frac{\alpha_0}{2}} & [(1-r)^2 + (1+r)^2 e^{i\alpha_0}] & 0\\ 0 & 0 & 4e^{i\beta_0} \end{pmatrix}.$$
(B.1)

This matrix can be diagonalized by using the unitary matrix

$$V_x = \begin{pmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}}e^{i\zeta} & 0\\ -\frac{1}{\sqrt{2}}e^{-i\zeta} & \frac{1}{\sqrt{2}} & 0\\ 0 & 0 & 1 \end{pmatrix} , \qquad (B.2)$$

where ζ is defined in Eq. (10). Consequently, we obtain

$$V_R = P_{ex} O_B V_x P_{ex}^T \text{diag}(e^{-i\frac{\beta_0}{2}}, e^{-i\frac{\alpha_0/2+\zeta}{2}}, e^{-i\frac{\alpha_0/2-\zeta}{2}}).$$
(B.3)

The explicit form is shown in Eq. (9), and those for M_i are given in Eq. (11).

- P. Minkowski, Phys. Lett. B 67, 421 (1977); M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, Proceedings of the Workshop, Stony Brook, New York, 1979, edited by P. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979), p. 315; T. Yanagida, in *Proceedings of the Workshop on the Unified Theories and Baryon Number in the Universe*, Tsukuba, Japan, 1979, edited by O. Sawada and A. Sugamoto(KEK Report No. 79-18,Tsukuba, 1979), p. 95; Prog. Theor. Phys. 64, 1103 (1980); R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44, 912 (1980).
- B. Pontecorvo, Zh. Eksp. Teor. Fiz. 53, 1717 (1967) [Sov. Phys. JETP 26, 984 (1968)]; Z. Maki,
 M. Nakagawa and S. Sakata, Prog. Theor. Phys. 28, 870 (1962).
- [3] Y. Nomura and T. Yanagida, Phys. Rev. D 59, 017303 (1999); Q. Shafi and Z. Tavartkiladze, Phys. Lett. B 451, 129 (1999); Phys. Lett. B 482, 145 (2000); Phys. Lett. B 487, 145 (2000);
 K. Choi, E. J. Chun, K. Hwang and W. Y. Song, Phys. Rev. D 64, 113013 (2001); A. Aranda, C. D. Carone and P. Meade, Phys. Rev. D 65, 013011 (2002).
- [4] S. M. Bilenky, J. Hosek and S. T. Petcov, Phys. Lett. B 94, 495 (1980).
 M. Doi, T. Kotani, H. Nishiura, K. Okuda and E. Takasugi, Phys. Lett. B 102, 323 (1981).
 J. Schechter and J. W. Valle, Phys. Rev. D 22, 2227 (1980); Phys. Rev. D 23, 1666 (1981).
- [5] J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. B 573, 652 (2000)
- [6] T. Miura, T. Shindou and E. Takasugi, Phys. Rev. D. 68, 093009 (2003).
- [7] S. Fukuda et al. [SuperKamiokande Collaboration], Phys. Rev. Lett. 86, 5651 (2001).
- [8] Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **87**, 071301 (2001); Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **89**, 011302 (2002); S. N. Ahmed *et al.* [SNO Collaboration], Phys. Rev. Lett. **92**, 181301 (2004).
- [9] K. Eguchi et al. [KamLAND Collaboration], Phys. Rev. Lett. 90, 021802 (2003).
- [10] T. Miura, T. Shindou and E. Takasugi, Phys. Rev. D. 66, 093002 (2002).
- [11] S. Antusch, J. Kersten, M. Lindner and M. Ratz, Phys. Lett. B 544, 1 (2002).
- [12] T. Shindou and E. Takasugi, Phys. Rev. D 70, 013005 (2004).
- [13] M. Fukugita and T. Yanagida, Phys. Lett. B 174, 45 (1986); M. Flanz, E. A. Paschos and U. Sarkar, Phys. Lett. B 345, 248 (1995) [Erratum-ibid. B 382, 447 (1996)]; L. Covi, E. Roulet and F. Vissani, Phys. Lett. B 384, 169 (1996); W. Buchmuller and M. Plumacher, Phys. Lett.

B **431**, 354 (1998).

- [14] W. Buchmuller, P. Di Bari and M. Plumacher, Phys. Lett. B 547, 128 (2002); Nucl. Phys. B 665, 445 (2003); New J. Phys. 6, 105 (2004); G. F. Giudice, A. Notari, M. Raidal, A. Riotto and A. Strumia, Nucl. Phys. B 685, 89 (2004).
- [15] A. Pilaftsis, Phys. Rev. D 56, 5431 (1997); A. Pilaftsis and T. E. J. Underwood, Nucl. Phys. B 692, 303 (2004).
- [16] E. K. Akhmedov, M. Frigerio and A. Y. Smirnov, JHEP 0309, 021 (2003).
- [17] M. Plumacher, Nucl. Phys. B 530, 207 (1998); A. Pilaftsis, Int. J. Mod. Phys. A 14, 1811 (1999); M. Flanz and E. A. Paschos, Phys. Rev. D 58, 113009 (1998); W. Buchmuller and M. Plumacher, Int. J. Mod. Phys. A 15, 5047 (2000); H. B. Nielsen and Y. Takanishi, Phys. Lett. B 507, 241 (2001); A. S. Joshipura, E. A. Paschos and W. Rodejohann, JHEP 0108, 029 (2001).
- [18] MEG collaboration, web site, http://meg.psi.ch/
- [19] N. Haba and N. Okamura, Eur. Phys. J. C 14, 347 (2000).
- [20] Y. Fukuda et al. [Super-Kamiokande Collaboration], Phys. Rev. Lett. 81, 1562 (1998).
- [21] M. Kawasaki, K. Kohri and T. Moroi; Phys. Lett. B 625, 7 (2005); hep-ph/0410287.
- [22] R. Allahverdi, S. Hannestad, A. Jokinen, A. Mazumdar and S. Pascoli, hep-ph/0504102.
- [23] D. N. Spergel et al. [WMAP Collaboration], Astrophys. J. Suppl. 148, 175 (2003).
- [24] F. Borzumati and A. Masiero, Phys. Rev. Lett. 57, 961 (1986); J. Hisano, T. Moroi, K. Tobe,
 M. Yamaguchi and T. Yanagida, Phys. Lett. B 357, 579 (1995); J. Hisano, T. Moroi, K. Tobe and M. Yamaguchi, Phys. Rev. D 53, 2442 (1996).
- [25] J. A. Casas and A. Ibarra, Nucl. Phys. B 618, 171 (2001).
- [26] S. T. Petcov, S. Profumo, Y. Takanishi and C. E. Yaguna, Nucl. Phys. B 676, 453 (2004).
- [27] S. Pascoli, S. T. Petcov and C. E. Yaguna, Phys. Lett. B 564, 241 (2003).
- [28] K. S. Babu, C. N. Leung and J. Pantaleone, Phys. Lett. B **319**, 191 (1993).
- [29] J. R. Ellis, J. Hisano, M. Raidal and Y. Shimizu, Phys. Rev. D 66, 115013 (2002); S. Pascoli,
 S. T. Petcov and W. Rodejohann, Phys. Rev. D 68, 093007 (2003).