

Some remarks on the behaviour of the finite element solution in nonsmooth domains

Nakao, Mitsuhiro T.
Faculty of Mathematics, Kyushu University

Kinoshita, Takehiko
Faculty of Mathematics, Kyushu University

<http://hdl.handle.net/2324/6098>

出版情報 : MHF Preprint Series. 2007-14, 2007-06-19. Faculty of Mathematics, Kyushu University
バージョン : draft
権利関係 :



MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

Some remarks on the behaviour of the finite element solution in nonsmooth domains

M. T. Nakao & T. Kinoshita

MHF 2007-14

(Received June 19, 2007)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

Some remarks on the behaviour of the finite element solution in nonsmooth domains

Mitsuhiro T. Nakao and Takehiko Kinoshita

Faculty of Mathematics, Kyushu University, Fukuoka 812-8581, Japan

Abstract. In this paper, we consider on the behaviour of the residual error by a smooth finite element solution for elliptic problems on nonconvex and nonsmooth domains. Against expectations, it is proved that the residual error is unbounded and actually diverges to infinity as the mesh size goes to zero. A numerical example which illustrates this phenomena will be presented for the Poisson equation on L-shaped domain using C^1 -Hermite element as well as the similar results will be shown for a C^0 element with a posteriori smoothing.

Keywords. Poisson equation, Non-smooth domain, Residual error.

1 Introduction

In this paper, we consider a finite element solution u_h of the following Poisson equation.

$$\begin{cases} -\Delta u &= f & \text{in } \Omega, \\ u &= 0 & \text{on } \partial\Omega. \end{cases} \quad (1)$$

Here, Ω is a bounded polygonal domain in \mathbb{R}^2 and f is a function in $L^2(\Omega)$. Then, in case of $u_h \in H^2(\Omega) \cap H_0^1(\Omega)$, the residual error $\|f + \Delta u_h\|_{L^2(\Omega)}$ plays an important role in the numerical enclosure methods of solutions for nonlinear elliptic problems (see, e.g., [3], [4], [7], [5] etc.). Let S_h be a finite dimensional subspace of $H_0^1(\Omega)$ dependent on the mesh size parameter h . Usually, u_h is defined as an element of S_h such that

$$(\nabla u_h, \nabla v_h)_{L^2(\Omega)} = (f, v_h)_{L^2(\Omega)}, \quad \forall v_h \in S_h. \quad (2)$$

In the below, we assume that, for solutions to (1) and (2),

$$u_h \rightarrow u \quad (h \rightarrow 0) \quad \text{in } H^1(\Omega), \quad (3)$$

which would be a natural condition for usual finite element subspaces.

If S_h is a C^0 element, since the residue $f + \Delta u_h$ no longer belongs to $L^2(\Omega)$, we need some smoothing procedure to get the residual estimation[6].

In this and the next sections, we assume that S_h is a C^1 finite element. For the convex domain Ω , assuming an inverse inequality for S_h , we easily have the following estimates:

$$\begin{aligned}
\|f + \Delta u_h\|_{L^2(\Omega)} &= \|\Delta(u - u_h)\|_{L^2(\Omega)} \\
&\leq Ch^{-1} \|\nabla(u - u_h)\|_{L^2(\Omega)} \\
&\leq C \|\Delta u\|_{L^2(\Omega)} \\
&= C \|f\|_{L^2(\Omega)},
\end{aligned} \tag{4}$$

where C is a general constant independent of h . Hence, Δu_h is bounded in h . There is, rather, a possibility to get some positive order estimates for $\|f + \Delta u_h\|_{L^2(\Omega)}$ in h provided that we use higher order polynomials. Therefore, we will naturally expect that Δu_h should also be bounded even if Ω is nonconvex, because the approximation scheme (2) is equivalent to

$$(-\Delta u_h, v_h)_{L^2(\Omega)} = (f, v_h)_{L^2(\Omega)}, \quad \forall v_h \in S_h, \tag{5}$$

which strongly suggests that Δu_h seems to be determined only by the function f . However, it is shown that this expectation is actually wrong. Namely, in Section 2, when u does not have H^2 smoothness, we prove that the concerning residual error is unbounded. In Section 3, we will present some computational results of these errors for the Poisson equation on L-shaped domain using C^1 Hermite functions, which confirm our theoretical assertion. Furthermore, we will show the similar result even for the case that we use some a posteriori smoothing technique with C^0 piecewise linear element. These should be interesting and rather surprising facts which is beyond our intuitive observation.

2 Unboundedness of residual error

In this section, let Ω be a nonconvex polygonal domain. Then, as well known, the weak solution u of (1) uniquely exists in $H_0^1(\Omega)$, and not necessarily belongs to $H^2(\Omega)$ (see [1]). We now describe the main result in this paper.

Theorem 2.1 *Let S_h be an C^1 finite element subspace on Ω , i.e., $S_h \subset H_0^1(\Omega) \cap H^2(\Omega)$, and let u_h be a solution of (2), or equivalently defined by (5). Then, the residual error has the following property.*

$$\lim_{h \rightarrow 0} \|f + \Delta u_h\|_{L^2(\Omega)} = \infty.$$

Proof. Let assume that the set $\left\{ \|\Delta u_h\|_{L^2(\Omega)} \right\}_{0 < h < 1}$ is bounded in \mathbb{R} . First, by (2) and the Poincaré inequality, we have

$$\begin{aligned}
\|\nabla u_h\|_{L^2(\Omega)}^2 &\leq \|f\|_{L^2(\Omega)} \|u_h\|_{L^2(\Omega)} \\
&\leq C_p \|f\|_{L^2(\Omega)} \|\nabla u_h\|_{L^2(\Omega)},
\end{aligned}$$

where C_p is a Poincaré constant. Therefore, $\left\{ \|u_h\|_{H_0^1(\Omega)} \right\}_{0 < h < 1}$ is also bounded. Next, since Ω is a polygon, we have the following well known result

$$\|\Delta u_h\|_{L^2(\Omega)} = |u_h|_{H^2(\Omega)},$$

which implies that $\{u_h\}_{0 < h < 1}$ is a bounded set in $H^2(\Omega)$.

Thus, by the weak compactness of $H^2(\Omega)$, there exists a subsequence $\{u_{h_n}\}_{n=1}^\infty$ in $\{u_h\}_{0 < h < 1}$, which weakly converge to some $\hat{u} \in H^2(\Omega)$. By the compactness of the embedding $H^2(\Omega) \hookrightarrow H^1(\Omega)$, we have the strong convergence:

$$u_{h_n} \rightarrow \hat{u} \quad (n \rightarrow \infty) \quad \text{in } H^1(\Omega).$$

On the other hand, u_{h_n} converges to the solution u of (1) in $H^1(\Omega)$ by the assumption (3). Therefore, by the uniqueness of the limit, we have $u = \hat{u}$, which implies that u has to be an element of $H^2(\Omega)$. This is a contradiction. \square

3 Numerical Examples

In this section, we show some numerical evidences for the actual divergent situations in the previous section, which suggest the difficulty to construct the approximate solution with convergent residual error, provided that the corresponding exact solution has no sufficient smoothness.

3.1 Smooth basis

We considered (1) with $f \equiv 1$ and Ω as a L-shaped domain in Fig. 1. We used the bi-cubic Hermite function as the basis of S_h with a uniform mesh in Fig.1. In this case S_h is a subspace of $H^2(\Omega)$.




Figure 1: domain Ω (mesh size $h = 1/5$)




Figure 2: approximate solution u_h

Fig.3 shows the computed results of residual errors, in which the horizontal axis means mesh size h , the vertical axis residual error $\|1 + \Delta u_h\|_{L^2(\Omega)}$ with logarithmic scale. By considering the results in Fig.3, we obtained the residual error with negative order in h , i.e., approximately $1.23h^{-0.33}$, which confirms us the divergence property proved in Theorem 2.1.




Figure 3: residual error $\|1 + \Delta u_h\|_{L^2(\Omega)}$

3.2 A posteriori smoothing by piecewise linear element

In [6], some a posteriori smoothing techniques were used to get the residual error for the C^0 element. We considered the same problem in the previous subsection by using the piecewise bilinear polynomial functions for the same mesh. Naturally, the finite element solution u_h of (2) is almost same contour as Fig.2. Since the direct calculation Δu_h is not possible, some smoothing procedure are taken as in [6]. Namely, we provided a piecewise bilinear finite element subspace S_h^* of $H^1(\Omega)$ which is constituted by adding the bases corresponding the boundary nodes to S_h . And define the vector function \bar{u}_h , denoted as $\bar{\nabla} u_h$, which means a smoothing of ∇u_h in $S_h^* \times S_h^*$ by

$$(\bar{u}_h, \bar{v}_h)_{L^2(\Omega)^2} = (\nabla u_h, \bar{v}_h)_{L^2(\Omega)^2}, \quad \forall \bar{v}_h \in S_h^* \times S_h^*. \quad (6)$$

Then define $\bar{\Delta} u_h \equiv \text{div } \bar{\nabla} u_h$.




Figure 4: smoothing error $\|\bar{\nabla} u_h - \nabla u_h\|_{L^2(\Omega)^2}$




Figure 5: residual error $\|1 + \bar{\Delta} u_h\|_{L^2(\Omega)}$

The smoothing error $\|\bar{\nabla} u_h - \nabla u_h\|_{L^2(\Omega)^2}$ and the residual error $\|1 + \bar{\Delta} u_h\|_{L^2(\Omega)}$ are shown in Fig.4 and in Fig.5, respectively, with the same scale as before. According to these computations, we observed that $\|\bar{\nabla} u_h - \nabla u_h\|_{L^2(\Omega)^2} \approx 0.35h^{0.80}$ and $\|1 + \bar{\Delta} u_h\|_{L^2(\Omega)} \approx 0.52h^{-0.33}$. These results suggest that it should be not possible to improve the residual error by this kind of a posteriori smoothing.

4 Concluding remarks

By Theorem 2.1 and our numerical experiments above, we could say that

1. When we use smooth approximate method by C^1 element it is proved that we can't constitute sufficient approximation to Δu .
2. Even if we take a smoothing method by some a posteriori techniques for the C^0 element, it could not be possible to improve the approximate property for Δu .
3. As an alternative approach, the singular function method, e.g., [2], might have the desired property for this problem, although it should be a little bit of unusual finite element method.

References

- [1] Grisvard, P., Singularities In Boundary Value Problems, Springer-Verlag(1992) ISBN : 3-540-55450-5.
- [2] Kobayashi, K., A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions, MHF Preprint Series, Kyushu University, MHF **2007-9**, 24 pages.
- [3] Nakao, M.T., Numerical verification methods for solutions of ordinary and partial differential equations, Numerical Functional Analysis and Optimization **22(3&4)** (2001), 321-356.
- [4] Nakao, M.T., Hashimoto, K. & Watanabe, Y., A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems, Computing **75** (2005), 1-14.
- [5] Plum, M., Computer-assisted enclosure methods for elliptic differential equations, Journal of Linear Algebra and its Applications **327** (2001), 147-187.
- [6] Yamamoto, N & Nakao, M.T., Numerical verifications of solutions for elliptic equations in nonconvex polygonal domains, Numer. Math. **65** (1993) 503-521.
- [7] Yamamoto, N. & Nakao, M.T., Numerical verifications for solutions to elliptic equations using residual iterations with higher order finite element, Journal of Computational and Applied Mathematics **60** (1995), 271-279.

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift
- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations

- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA
Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS
Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI
Nonlinear regression modeling via regularized wavelets and smoothing parameter selection
- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO, Yoshihiro MIZOGUCHI & Yasuo KAWAHARA
On reversible cellular automata with finite cell array

- MHF2005-28 Toru KOMATSU
Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU
A computational approach to constructive a priori and a posteriori error estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO
Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI
A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI
A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI
An explicit formula of the Shapley value for a cooperative game induced from the conjugate point
- MHF2005-34 Hideki MURAKAWA
A regularization of a reaction-diffusion system approximation to the two-phase Stefan problem
- MHF2006-1 Masahisa TABATA
Numerical simulation of Rayleigh-Taylor problems by an energy-stable finite element scheme
- MHF2006-2 Ken-ichi MARUNO & G R W QUISPEL
Construction of integrals of higher-order mappings
- MHF2006-3 Setsuo TANIGUCHI
On the Jacobi field approach to stochastic oscillatory integrals with quadratic phase function
- MHF2006-4 Kouji HASHIMOTO, Kaori NAGATOU & Mitsuhiro T. NAKAO
A computational approach to constructive a priori error estimate for finite element approximations of bi-harmonic problems in nonconvex polygonal domains
- MHF2006-5 Hidefumi KAWASAKI
A duality theory based on triangular cylinders separating three convex sets in R^n
- MHF2006-6 Raimundas VIDŪNAS
Uniform convergence of hypergeometric series
- MHF2006-7 Yuji KODAMA & Ken-ichi MARUNO
N-Soliton solutions to the DKP equation and Weyl group actions

- MHF2006-8 Toru KOMATSU
Potentially generic polynomial
- MHF2006-9 Toru KOMATSU
Generic sextic polynomial related to the subfield problem of a cubic polynomial
- MHF2006-10 Shu TEZUKA & Anargyros PAPAGEORGIOU
Exact cubature for a class of functions of maximum effective dimension
- MHF2006-11 Shu TEZUKA
On high-discrepancy sequences
- MHF2006-12 Raimundas VIDŪNAS
Detecting persistent regimes in the North Atlantic Oscillation time series
- MHF2006-13 Toru KOMATSU
Tamely Eisenstein field with prime power discriminant
- MHF2006-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the Hankel determinant formula for the solutions of the Painlevé IV equation
- MHF2006-15 Raimundas VIDŪNAS
Darboux evaluations of algebraic Gauss hypergeometric functions
- MHF2006-16 Masato KIMURA & Isao WAKANO
New mathematical approach to the energy release rate in crack extension
- MHF2006-17 Toru KOMATSU
Arithmetic of the splitting field of Alexander polynomial
- MHF2006-18 Hiroki MASUDA
Likelihood estimation of stable Lévy processes from discrete data
- MHF2006-19 Hiroshi KAWABI & Michael RÖCKNER
Essential self-adjointness of Dirichlet operators on a path space with Gibbs measures via an SPDE approach
- MHF2006-20 Masahisa TABATA
Energy stable finite element schemes and their applications to two-fluid flow problems
- MHF2006-21 Yuzuru INAHAMA & Hiroshi KAWABI
Asymptotic expansions for the Laplace approximations for Itô functionals of Brownian rough paths
- MHF2006-22 Yoshiyuki KAGEI
Resolvent estimates for the linearized compressible Navier-Stokes equation in an infinite layer

- MHF2006-23 Yoshiyuki KAGEI
Asymptotic behavior of the semigroup associated with the linearized compressible Navier-Stokes equation in an infinite layer
- MHF2006-24 Akihiro MIKODA, Shuichi INOKUCHI, Yoshihiro MIZOGUCHI & Mitsuhiro FUJIO
The number of orbits of box-ball systems
- MHF2006-25 Toru FUJII & Sadanori KONISHI
Multi-class logistic discrimination via wavelet-based functionalization and model selection criteria
- MHF2006-26 Taro HAMAMOTO, Kenji KAJIWARA & Nicholas S. WITTE
Hypergeometric solutions to the q -Painlevé equation of type $(A_1 + A'_1)^{(1)}$
- MHF2006-27 Hiroshi KAWABI & Tomohiro MIYOKAWA
The Littlewood-Paley-Stein inequality for diffusion processes on general metric spaces
- MHF2006-28 Hiroki MASUDA
Notes on estimating inverse-Gaussian and gamma subordinators under high-frequency sampling
- MHF2006-29 Setsuo TANIGUCHI
The heat semigroup and kernel associated with certain non-commutative harmonic oscillators
- MHF2006-30 Setsuo TANIGUCHI
Stochastic analysis and the KdV equation
- MHF2006-31 Masato KIMURA, Hideki KOMURA, Masayasu MIMURA, Hidenori MIYOSHI, Takeshi TAKAISHI & Daishin UEYAMA
Quantitative study of adaptive mesh FEM with localization index of pattern
- MHF2007-1 Taro HAMAMOTO & Kenji KAJIWARA
Hypergeometric solutions to the q -Painlevé equation of type $A_4^{(1)}$
- MHF2007-2 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Verified numerical computation of solutions for the stationary Navier-Stokes equation in nonconvex polygonal domains
- MHF2007-3 Kenji KAJIWARA, Marta MAZZOCCO & Yasuhiro OHTA
A remark on the Hankel determinant formula for solutions of the Toda equation
- MHF2007-4 Jun-ichi SATO & Hidefumi KAWASAKI
Discrete fixed point theorems and their application to Nash equilibrium
- MHF2007-5 Mitsuhiro T. NAKAO & Kouji HASHIMOTO
Constructive error estimates of finite element approximations for non-coercive elliptic problems and its applications

- MHF2007-6 Kouji HASHIMOTO
A preconditioned method for saddle point problems
- MHF2007-7 Christopher MALON, Seiichi UCHIDA & Masakazu SUZUKI
Mathematical symbol recognition with support vector machines
- MHF2007-8 Kenta KOBAYASHI
On the global uniqueness of Stokes' wave of extreme form
- MHF2007-9 Kenta KOBAYASHI
A constructive a priori error estimation for finite element discretizations in a non-convex domain using singular functions
- MHF2007-10 Myoungnyoun KIM, Mitsuhiro T. NAKAO, Yoshitaka WATANABE & Takaaki NISHIDA
A numerical verification method of bifurcating solutions for 3-dimensional Rayleigh-Bénard problems
- MHF2007-11 Yoshiyuki KAGEI
Large time behavior of solutions to the compressible Navier-Stokes equation in an infinite layer
- MHF2007-12 Takashi YANAGAWA, Satoshi AOKI and Tetsuji OHYAMA
Human finger vein images are diverse and its patterns are useful for personal identification
- MHF2007-13 Masahisa TABATA
Finite element schemes based on energy-stable approximation for two-fluid flow problems with surface tension
- MHF2007-14 Mitsuhiro T. NAKAO & Takehiko KINOSHITA
Some remarks on the behaviour of the finite element solution in nonsmooth domains