

Collecting Records of *Spalangia* Latreille, 1805 (Hymenoptera: Pteromalidae) Parasitic on Pupae of *Stomoxys calcitrans* (Linnaeus, 1758) and *Musca domestica* Linnaeus, 1758 (Diptera: Muscidae) in Japan

MATSUO, Kazunori
Faculty of Social and Cultural Studies, Kyushu University

ASAMI, Koshi
School of Interdisciplinary Science and Innovation, Kyushu University

SHINDO, Makito
School of Interdisciplinary Science and Innovation, Kyushu University

ARAKI, Hiromitsu
Faculty of Economics, Kyushu University

他

<https://doi.org/10.5109/6613527>

出版情報 : ESAKIA. 55, pp.7-10, 2022-12-20. 九州大学大学院農学研究院昆虫学教室
バージョン :
権利関係 :



Collecting Records of *Spalangia* Latreille, 1805 (Hymenoptera: Pteromalidae) Parasitic on Pupae of *Stomoxys calcitrans* (Linnaeus, 1758) and *Musca domestica* Linnaeus, 1758 (Diptera: Muscidae) in Japan

Kazunori MATSUO^{1)*}, Koshi ASAMI²⁾, Makito SHINDO²⁾, Hiromitsu ARAKI³⁾, Junko KOHARA⁴⁾,
Hitoshi TSUKAHARA⁵⁾, Hiro TAKAHASHI⁶⁾, Hideki NAGAI⁷⁾, Takeshi MIZUKI⁸⁾, Kazutoshi
YOSHIMURA⁹⁾, Kazuaki KAKIUCHI^{9)**}, Kazuyuki MIZUGUCHI¹⁰⁾, Shuichi YONEDA¹⁰⁾, Takuya
ARIDOME¹¹⁾, Takahiro SAINO¹¹⁾, Naoko GUSHI¹²⁾, Kyoki TERUYA¹²⁾, Keisuke NINOMIYA¹³⁾ and
Yoshihisa ABE¹⁾

1) Faculty of Social and Cultural Studies, Kyushu University, Fukuoka, 819-0395 Japan

2) School of Interdisciplinary Science and Innovation, Kyushu University, Fukuoka, 819-0395 Japan

3) Faculty of Economics, Kyushu University, Fukuoka, 819-0395 Japan

4) Animal Research Center, Agricultural Research Department, Hokkaido Research Organization,
Hokkaido, 081-0038 Japan

5) Tochigi Prefectural Livestock & Dairy Experimental Center, Tochigi 329-2747 Japan

6) Animal Husbandry Division, Aichi Agricultural Research Center, Aichi 480-1193 Japan

7) Tanba Agriculture Extension Center, Hyogo 669-3309 Japan

8) Okayama Prefectural Technology Center for Agriculture, Forestry & Fisheries, Okayama 709-3494
Japan

9) Nansatsu Regional Affairs Department, Kagoshima 897-0031 Japan

10) Kagoshima Dairy Co-operative Association, Kagoshima 890-0073 Japan

11) Japan Agricultural Cooperatives Minami-Satsuma, Kagoshima 897-0302 Japan

12) Okinawa Prefectural Livestock and Grassland Research Center, Okinawa 905-0426 Japan

13) Okinawa Prefectural Miyako Agricultural Division, Okinawa 906-0012 Japan

* E-mail: matsuosudachi@scs.kyushu-u.ac.jp

** Current address: Livestock Division, Kagoshima Prefectural Government, Kagoshima 890-8577
Japan

Abstract. The present study provided new collecting records of the following *Spalangia* species in Japan: *S. cameroni* Perkins, 1910; *S. endius* Walker, 1839; *S. nigra* Latreille, 1805; and *S. nigroaenea* Curtis, 1839.

Key words: Distribution, pupal parasitoid, the house fly, the stable fly.

Introduction

Members of *Spalangia* Latreille, 1805 (Hymenoptera: Pteromalidae) are known as pupal parasitoids of Diptera, such as Calliphoridae, Muscidae, and Tephritidae (Gibson 2009; Noyes 2019). Until now, six species of *Spalangia* have been known to Japan: *S. cameroni* Perkins, 1910 (Matsuo 2020); *S. endius* Walker, 1839 (Tachikawa 1965); *S. gemina* Bouček, 1963 (Ogawa *et al.* 2009); *S. nigra* Latreille, 1805 (Nakasuji 1963; Kamijo & Yamamoto 2000); *S. nigroaenea* Curtis, 1839 (Ogawa *et al.* 2009); and *S. simplex* Perkins,

1910 (Nagase 2004). Because some of *Spalangia* species have been known as biological control agents against the stable fly, *Stomoxys calcitrans* (Linnaeus, 1758) and the house fly, *Musca domestica* Linnaeus, 1758 (Diptera: Muscidae) (Morgan *et al.* 1975; Morgan & Patterson 1990; Legner 1995; Skovgård 2004), we are accumulating their collecting records in Japan. Fly pupae that were collected from bovine manure were identified based on shape of the posterior spiracles mentioned in Rochon *et al.* (2021). Parasitoids were identified by using a key to species provided by Gibson (2009) and Matsuo (2020)

Table. Specimens of *Spalangia* spp. examined. They have been kept in the collection of the BLKU.

Species name and No. of individuals	Host	Collecting site (collector*)	Date of collecting	Date of emergence
<i>Spalangia cameroni</i>				
1 ♀	<i>Stomoxys calcitrans</i>	Kawasaki, Fukuoka, Japan (KM)	21 November 2018	December 2018
1 ♀ 1 ♂	<i>Stomoxys calcitrans</i>	Chiran, Minamikyushu, Kagoshima, Japan (KM, HA, KA, MS)	28 October 2021	30 November 2021
2 ♀ 1 ♂	<i>Stomoxys calcitrans</i>	Chiran, Minamikyushu, Kagoshima, Japan (KM, HA, KA, MS)	28 October 2021	8 December 2021
1 ♀	<i>Musca domestica</i>	Nakijin, Kunigami, Okinawa, Japan (KM, HA, KA, MS)	18 October 2021	29 November 2021
1 ♂	<i>Musca domestica</i>	Nakijin, Kunigami, Okinawa, Japan (KM, HA, KA, MS)	18 October 2021	30 November 2021
<i>Spalangia endius</i>				
2 ♀ 1 ♂	<i>Stomoxys calcitrans</i>	Miyako, Fukuoka, Japan (KM)	14 November 2018	December 2018
4 ♀	<i>Stomoxys calcitrans</i>	Yazakosagamine, Nagakute, Aichi, Japan (HT)	9 August 2021	11 August 2021
2 ♀	<i>Stomoxys calcitrans</i>	Ono, Hyogo, Japan (KM, HA)	12 October 2021	25 October 2021
1 ♂	<i>Musca domestica</i>	Ei, Minamikyushu, Kagoshima, Japan (KM, HA, KA, MS)	29 October 2021	8 December 2021
1 ♀	<i>Musca domestica</i>	Ei, Minamikyushu, Kagoshima, Japan (KM, HA, KA, MS)	29 October 2021	10 December 2021
1 ♀	<i>Musca domestica</i>	Nakijin, Kunigami, Okinawa, Japan (KM, HA, KA, MS)	18 October 2021	15 November 2021
1 ♀	<i>Musca domestica</i>	Nakijin, Kunigami, Okinawa, Japan (KM, HA, KA, MS)	18 October 2021	24 November 2021
<i>Spalangia nigra</i>				
1 ♀ 1 ♂	<i>Chrysomya pinguis</i>	Miyako, Fukuoka, Japan (KM)	17 May 2018	18 June 2018
2 ♀	<i>Stomoxys calcitrans</i>	Yazakosagamine, Nagakute, Aichi, Japan (HT)	10 June 2021	17 June 2021
<i>Spalangia nigroaenea</i>				
4 ♀ 2 ♂	<i>Stomoxys calcitrans</i>	Miyako, Fukuoka, Japan (KM)	15 May 2019	3-4 June 2019
4 ♀ 2 ♂	<i>Stomoxys calcitrans</i>	Yazakosagamine, Nagakute, Aichi, Japan (HT)	9 August 2021	11 August 2021
1 ♂	<i>Stomoxys calcitrans</i>	Ei, Minamikyushu, Kagoshima, Japan (KM, HA, KA, MS)	29 October 2021	4 November 2021
1 ♂	<i>Stomoxys calcitrans</i>	Nakijin, Kunigami, Okinawa, Japan (KM, HA, KA, MS)	18 October 2021	17 November 2021

*Names of collectors: KM Kazunori Matsuo, HA Hiromitsu Araki, HT Hiro Takahashi, KA Koshi Asami, MS Makito Shindo

as well as reference specimens kept in the Hokkaido University Museum (HUM) and the Natural History Museum, London (BMNH). Specimens obtained in this study were deposited in the collection of the Biosystematics Laboratory, Faculty of Social and Cultural Studies, Kyushu University, Japan (BLKU).

Results

Spalangia cameroni Perkins, 1910

(Table)

Distribution. Afrotropic, Australasia, Nearctic, Neotropic, Oriental, and Palaearctic (Noyes 2019). Japan (Fukuoka, Kagoshima, and Okinawa). The present study newly recorded *S. cameroni* from Kagoshima and Okinawa Prefectures, Japan.

Spalangia endius Walker, 1839

(Table)

Distribution. Afrotropic, Australasia, Nearctic, Neotropic, Oriental, and Palaearctic (Noyes 2019). Japan (Aichi, Hyogo, Fukuoka, Kagoshima, and Okinawa). The present study newly recorded *S. endius* from Aichi, Hyogo, Kagoshima, and Okinawa Prefectures, Japan.

Spalangia nigra Latreille, 1805

(Table)

Distribution. Afrotropic, Australasia, Nearctic, Neotropic, Oriental, and Palaearctic (Noyes 2019). Japan (Kanagawa, Aichi, Yamaguchi, Ehime, and Fukuoka). The present study newly recorded *S. nigra* from Aichi Prefecture, Japan.

Spalangia nigroaenea Curtis, 1839

(Table)

Distribution. Afrotropic, Australasia, Nearctic, Neotropic, Oriental, and Palaearctic (Noyes 2019). Japan (Aichi, Kochi, Fukuoka, Kagoshima, and Okinawa). The present study newly recorded *S. nigroaenea* from Aichi, Kagoshima, and Okinawa Prefectures, Japan.

Acknowledgments

We thank Messrs Kazuyuki Tada (Iwate Agricultural Research Center Animal Industry Research Institute), Tatsuya Koike (Tochigi Prefectural Livestock & Dairy Experimental Center), and Tomoki Kojima (Aichi Agricultural Research Center) for supporting field surveys. This study was supported partly by JSPS

References

- KAKENHI grant number 19H00942, JSPS KAKENHI grant number 21K05600, JST A-STEP (Adaptable and Seamless Technology transfer Program through Target-driven R&D) grant number JPMJTM20S6, and Kyushu University-initiated venture business seed development program.
- Gibson GAP, 2009. Revision of New World Spalangiinae (Hymenoptera: Pteromalidae). *Zootaxa*, **2259**: 1–159.
- Kamijo K & Yamamoto E, 2000. Chalcidoid wasps of Odamiyama and its adjacent areas. *Nature of Odamiyama*, **2**: 755–767.
- Legner EF, 1995. Biological control of Diptera of medical and veterinary importance. *Journal of Vector Ecology*, **20**: 59–120.
- Matsuo K, 2020. Detection of *Spalangia* (Hymenoptera: Pteromalidae) attacking *Stomoxys calcitrans* (Diptera: Muscidae) and their potential significance in the Japanese cattle industry. *Applied Entomology and Zoology*, **55**: 83–91.
- Morgan PB & Patterson RS, 1990. Efficiency of target formulations of pesticides plus augmentative releases of *Spalangia endius* Walker (Hymenoptera: Pteromalidae) to suppress populations of *Musca domestica* L. (Diptera: Muscidae) at poultry installations in the Southeastern United States. In: Rutz DA & Patterson RS (eds.) *Biocontrol of Arthropods Affecting Livestock and Poultry*, pp. 69–78. Westview Press, Colorado.
- Morgan PB, Patterson RS, LaBrecque GC, Weidhaas DE & Benton A. 1975. Suppression of a field population of houseflies with *Spalangia endius*. *Science*, **189**: 388–389.
- Nagase H, 2004. Hymenoptera. In: Kanagawa Konchu Danwakai (eds.) *Insect fauna of Kanagawa*, pp. 1272. Kanagawa Konchu Danwakai, Odawara.
- Nakasuji F, 1963. *Spalangia nigra* Latreille, new to Japan. *Kontyû*, **31**: 248.
- Noyes JS, 2019. Universal Chalcidoidea database. Available from URL: <http://www.nhm.ac.uk/chalcidoidea>. [Accessed 20 January 2022]
- Ogawa K, Arakawa R & Fukuda T, 2009. First records of *Spalangia gemina* Bouček and *S. nigroaenea* Curtis (Hymenoptera: Pteromalidae) in Japan. *Japanese Journal of*

- Systematic Entomology*, **15**: 277–285.
- Rochon K, Hogsette JA, Kaufman PE, Olafson PU, Swiger SL & Taylor DB, 2021. Stable fly (Diptera: Muscidae) – biology, management, and research needs. *Journal of Integrated Pest Management*, **12**: 38.
- Skovgård H, 2004. Sustained releases of the pupal parasitoid *Spalangia cameroni* (Hymenoptera: Pteromalidae) for control of house flies, *Musca domestica* and stable flies *Stomoxys calcitrans* (Diptera: Muscidae) on dairy farms in Denmark. *Biological Control*, **30**: 288–297.
- Tachikawa T, 1965. Pteromalidae. In: Asahina S, Ishihara T & Yasumatsu K (eds.) *Iconographia Insectorum Japonicorum Colore Naturali Edita III*, p. 271. Hokuryukan, Tokyo.