

Heavy Metals Contamination and Their Impacts on Fish Responses in Porong River, East Java, Indonesia

Nico Rahman Caesar

Doctoral Programme of Environment Science, Post Graduate School, Universitas Brawijaya

Yanuhar, Uun

Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya

Musa, Muhammad

Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya

Ciptadi, Gatot

Doctoral Programme of Environment Science, Post Graduate School, Universitas Brawijaya

他

<https://doi.org/10.5109/7148443>

出版情報 : Evergreen. 10 (3), pp.1218-1230, 2023-09. 九州大学グリーンテクノロジー研究教育センター

バージョン :

権利関係 : Creative Commons Attribution-NonCommercial 4.0 International



Heavy Metals Contamination and Their Impacts on Fish Responses in Porong River, East Java, Indonesia

Nico Rahman Caesar¹, Uun Yanuhar^{2*}, Muhammad Musa², Gatot Ciptadi^{1,3}, Heru Suryanto⁴, Yusuf Arif Wahyudi⁵, and Rachmat Noer Soelistyoadi⁵

¹Doctoral Programme of Environment Science, Post Graduate School, Universitas Brawijaya, Jalan Veteran, Malang, East Java, 65145, Indonesia

²Department of Aquatic Resources Management, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Jalan Veteran, Malang, East Java, 65145, Indonesia

³Faculty of Animal Husbandry, Universitas Brawijaya, Malang, Jalan Veteran, Malang – East Java, Indonesia, 65145

⁴Department of Mechanical Engineering, Faculty of Engineering, Universitas Negeri Malang, Malang, 65145, Indonesia

⁵Fish Quarantine Center for Quality Control and Safety of Fishery Products Surabaya I, Ministry of Marine Affairs and Fisheries, Surabaya, East Java, 60115, Indonesia

*corresponding author: doktoruun@ub.ac.id

(Received January 19, 2023; Revised August 2, 2023; accepted September 23, 2023).

Abstract: The presence of heavy metals and other contaminants may indicate the presence of a biological response in aquatic animals. This research investigates the heavy metal contamination and water quality on the immune response (IL-1 β) of wader fish (*Barbonymus gonionotus*) in response to the presence of contaminants in the waters of the Porong River. The location area was selected because of its diverse utilization surrounding the river, including residential areas, industrial zones, and agricultural practices. The research procedure includes water quality measurements, plankton analysis, heavy metals exposure, and immune response (IL-1 β) detection by PCR. In general, the waters of the Porong River were found to be polluted based on the assessment of water quality, as indicated by the measurement of CO₂, TDS, TSS, and salinity parameters surpassing the established threshold values. In the waters of the Porong River, the metals Cu, Hg, Cd, and Pb exhibited a descending order of penetrating power as follows: Cu > Hg > Pb > Cd. Conversely, in the sediment samples, the order of penetrating power was Cu > Pb > Cd > Hg. The PCR amplification results indicated the presence of IL-1 β response in all examined fish, suggesting that it could be attributed to water contamination throughout the entire sampling site. In addition, the heavy metals contamination in the waters of the Porong River resulted in high expression of IL-1 β in the fish sample. The findings lead to the conclusion that the waters of the Porong River are classified as polluted based on the measurements of water quality parameters, namely CO₂, TDS, TSS, and salinity, which surpass the threshold values established according to Government Regulation (PP) of the Republic of Indonesia Number 22 of 2021.

Keywords: *water quality, heavy metals, IL-1 β , PCR, polluted waters*

1. INTRODUCTION

The Porong River is a downstream area of the Brantas River which has a high potential for heavy metal pollution, among others, due to polluting activities in Tarik, Prambon, Krembung, Porong, and Jabon districts¹⁾. In addition, the Porong River is an example of a river that has experienced environmental pollution caused by the Lapindo mud overflow²⁾. Serious pollution problems can occur, including industrial wastewater containing heavy

metals, sometimes entering the environment accidentally or illegally, and sedimentation of land, rivers, and groundwater³⁾. Pollution often affects the pace of development, in particular in the industrial sector, since its creation produces waste. Sediments from the Porong River include levels of lead (Pb) and cadmium (Cd), which are two to three times passed the permissible limit at least⁴⁾. Changes in land use or physical characteristics have affected the soil's ability to retain more water during rainfall before the direct runoff. The ability to make

hydrological forecasts has also become an important part of the sustainable management of water resources, water quality, and water-related natural threats⁵. In fact, water spaces such as streams, rivers, and lakes in particular, have an important role in creating urban cooling because water spaces have a higher evaporation rate than green open spaces⁶. Besides, aquatic biota is an excellent indicator of metal contamination in the aquatic environment⁷. Metabolic activity can accumulate heavy metals. Thus, they can get into the tissues of fish organs and cause cell swelling, necrosis, fibrosis, and cirrhosis⁸. Heavy metals are a core group of water pollutants because they have bioaccumulating properties and are non-degradable. Heavy metals can be found in aquatic environments through human activities such as the mining industry⁹. Methods, such as desalination¹⁰, membrane¹¹, and bioremediation technology for sustainable wastewater treatment¹², have been developed by researchers to get clean water. Other researchers also focus on wastewater monitoring, especially detecting the heavy metals contamination in water is very important¹³.

Currently, high heavy metal levels in the river waters are generally caused by the high input of industrial, mining, agricultural, and domestic waste¹⁴. In addition, heavy metals can accumulate in the fish environment. During the last 100 years, Pb, Cr, and Cd have been used in particular in paints, preservatives, toy manufacture, pesticides, lead bullets, and as gas additives¹⁵. Concentrations of dissolved Cr, Cd, and Pb in freshwater are generally between 10 and 500 ng/l, but concentrations of more than 1 mg/l have been observed in some industrial areas¹⁶. In heavily polluted areas, Cr, Cd, and Pb contents in sediments and soil can reach more than 1 mg/kg¹⁷. The presence of heavy metals and other contaminants may indicate the presence of a biological response in aquatic animals. Fish that live in contaminated water conditions in rivers, one of which is indicated by the emergence of an infant immune response namely IL-1 β ¹⁸, is also shown in the type of Teleostei fish with the presence of IL-1 β ¹⁹. The presence of IL-1 β also showed a response due to heavy metal pollution¹⁴.

The Porong River in Sidoarjo Regency, East Java Province, has reported unexpected fish deaths in 2019²⁰. The majority of the dead species were unique freshwater fish biota in the Porong River such as Keting, Kuthuk, and Bader fish²⁰. One of the responses to fish mortality, because of the high expression of IL-1 β caused by the contamination of heavy metals and organic matter in the water¹⁹. Research by El-Boshy and Taha²¹ showed that heavy metals could impair the cellular and humoral immune responses of fish. Among cell immunological reactions, one of which can be seen by the presence of macrophage phagocytosis activity. Macrophages and monocytes produce and secrete IL-1 β . It is produced as an inactive 31 kDa precursor called Pro-IL-1 β in response to a molecular motif that acts on macrophages via pattern recognition receptors (PRR) to regulate pathways that

regulate gene expression²².

The activity of interferon-alpha cytokines (IFN- α) was the first to be studied. However, in the early 1980s, the first molecular cloning, such cytokines (IFN α , IL-1 β , IL-2, and TNF α), was discovered. Research on the regulation of fish immunity began in the early 1990s because of the importance of fish in the aquaculture industry and the need to understand the issues related to immunity and disease. The first cloned cytokine in fish was TGF- β , followed by IL-1 β and a few years later by TNF- α ²³. This specific cytokine was also the best and most studied property of the 11 IL-1 β family members. Although most research are focused on the production of innate immune cells such as monocytes and macrophages, it is produced and secreted by a wide variety of cell types. IL-1 β is a potent pro-inflammatory cytokine produced by cells of the innate immune system. It is produced without a signal sequence and does not follow conventional protein secretion pathways but instead uses one or more unconventional secretory pathways. This research aims to investigate the heavy metal contamination and water quality on the immune response (IL-1 β) of wader fish (*Barbonymus gonionotus*) in response to the presence of contaminants in the waters of the Porong River.

2. MATERIALS AND METHODS

2.1 Study Area

The location area of this study is presented in Fig. 1. The Porong watershed in Sidoarjo District in East Java Province was selected as the study site. The upstream area of the river is represented by the city of Batu, and the downstream part of the river is represented by Sidoarjo Regency. The consideration of choosing the location is due to varied utilization of land around the river, such as residential areas, industrial sites, and agricultural activities that affect pollution to the river. In these two locations, we determined five (5) points selected for the sampling location in this study.

Table 1. The Map of coordinates and elevation of the sampling site in the Porong River

Site	Latitude	Longitude	Altitude (AMSL)
1	7°32'46.0"S	112°41'55.1"E	8
2	7°32'33.7"S	112°46'26.5"E	3
3	7°33'57.0"S	112°52'11.1"E	3
4	7°45'14.0"S	112°31'35.6"E	1642




Fig. 1: Research Location in the Porong River.

2.2 Water Quality Measurements

Porong River water quality analysis with the presence of *B. gonionotus* fish consists of physical parameters (Total Dissolved Solid/TDS, Total Suspended Solid/TSS, temperature), and chemical parameters (pH, Dissolved Oxygen/DO, CO₂, nitrate, orthophosphate, Total Organic Matter/TOM, ammonia, salinity). The tools used consist of TSS meters, thermometers, pH meters, DO meters, Erlenmeyer, hot plates, cuvetts, spectrophotometers, and porcelain cups. The materials used consist of Lugol, KMnO₄, H₂SO₄, na-oxalate, distillate water, Nessler, phenolphthalein indicator, Na₂CO₃, Disulfonic phenol acid, NH₄OH, Ammonium molybdate, and SnCl₂.

2.3 Plankton Analysis

Based on Nurruhwati et al.²⁴⁾, plankton samples were taken by filtering 10 liters of water through a plankton net equipped with a storage bottle with a mesh size of 200 µm. The accommodated plankton was preserved with a 4% formalin solution of 3 drops. Calculation and identification of plankton were carried out at the Fisheries Center of the Faculty of Fisheries and Marine Sciences, University of Brawijaya, using a microscope and a plankton identification book²⁵⁾ and The Marine and Fresh Water Plankton²⁶⁾. Observations were made by means of a sample bottle containing the sample being shaken slowly until homogeneous. The sample was taken using a drip pipette and dripped on a prepare glass. Next, it was observed using a microscope with an enlargement of 100x and 400x. The obtained sample was identified using a plankton identification book, and the sample was observed as many as three repetitions. Plankton sampling was carried out at a single point, where the measurements were carried out weekly. The plankton sampling procedure is as follows: 1) Take 25 liters of pool water at five different points by 5 liters at each point; 2) Filtering pool water using plankton net number 25; 3) Precipitating plankton at the end of the planktonnet using a film bottle; 4) Adding 3-4 drops of lugol solution; 5) Plankton samples are stored and taken to the laboratory for identification.

2.4 Heavy Metals Exposure

The method of Heavy Metal Examination in the water was in accordance with the American Public Health Association (APHA), American Water Works Association (AWWA), & Water Environment Federation (WEF)²⁷⁾. Identification analysis of the heavy metal's type was used as one of the parameters of water quality. Heavy metal type analysis was carried out using the Inductively Coupled Plasma Mass Spectrometer (ICP-MS) test method. The tools used consist of electronic scales and baking sheets, microwave digestion, desiccator, and drying oven. The materials used consist of internal standard indium, HNO₃, internal standard bismuth, and pure water. The samples used were in the form of water and sediment from each sampling point. Observed heavy metal were Pb, Cd, Hg, and Cu concentrations.

2.5 Detection of Immune response (IL-1β) by PCR

DNA was extracted from tissues preserved in absolute ethanol using a silica extraction kit (Gene). Each sample of fish kidney tissue was put into a 1.5 ml microtube, added 900 µl GL buffer, grounded with a pestle grinder, and centrifuged at 12,000 rpm (3 min). As much as 600 µL of the coating solution obtained was transferred to a new 1.5 mL microfuge tube, 40 µL silica was added, vortexed until homogeneous, and washed at 12,000 rpm (15 sec). After centrifugation, the supernatant was discarded, the silica pellet was washed with 500 µl GT buffer, vortex until the silica pellet formed a suspension and centrifuged at 12,000 rpm (15 sec). Then, the supernatant was discarded again, added by 1 ml of 70% ethanol wash flint and mixed with pellets until the flint pellets formed a suspension. The solution was centrifuged again at 12,000 rpm (15 sec), the ethanol was discarded, while the pellet was collected using a micropipette and added by 1 mL of ddH₂O to suspend the silica pellet. Finally, the silica pellets was put into vortex to create a suspension.

Specific primers of IL-1β used as follows F: ACGCCACCAAGAGCCTTTTA; R: GCAGCCCATATTTGGTCAGA with a gene target of 206 bp. PCR amplification was performed using a 25 µl volume PCR reaction mixture consisting of a master mix (KAPA Biosystems, KK510) 2 µl forward primer, 2 µl reverse primer, 2 µl template DNA, and 19 µl nuclease-free water. Amplification was carried out by setting the predenaturation temperature to 95 °C (3 minutes), followed by denaturation to 95 °C (30 seconds), annealing, extension 72°C (1 min, 34 cycles) and the final stage of 72 °C (5 min). Temperature optimization was carried out to obtain the best annealing temperature by adjusting the thermal cycler gradient (T 100 Bio-Rad) at 4 different temperature levels, namely 58.0°C, 56.1°C, 53.8°C, and 51.9°C.

3. RESULTS

Based on the water quality measurement data in Table 2, the results of the water quality measurement for point 1 (upstream of the Porong River) were quite good with the parameters of TDS 87.33 mg/L, DO 16.13 mg/L, and pH 7.29, which in accordance with river water quality standards (PP. No. 22 of 2021).

The CO₂ levels at point 2 were higher above the recommended range for river water quality, at 27.72 mg/L. Point 3 had a salinity of 9.05 ppt and CO₂ concentration of 35.64 mg/L, which was extremely high for the waters. More metrics such as TDS 6792.50 mg/L, TSS 107.95 mg/L, CO₂ 23.76 mg/L, and salinity 8.60 ppt, were above

the river water quality level (PP No. 22 of 2021) at point 4.

The standard quality value for suspended residue or suspended solids (TSS) concentration in river water is not more than 50 mg/L²⁸⁾. The optimal TDS level that complies with water quality standards in river water is not more than 1000 mg/L. Inorganic substances such as minerals, metals, salts, and soluble cations in water are included in the TDS content in the water²⁹⁾. The standard for carbon dioxide concentration in water is approximately 12 mg/L. Elevated levels of carbon dioxide in water can convert it into a toxic gas for fish and fish embryos³⁰⁾.

Table 2. Results of water quality measurements in the Porong River.

Parameter	Standard	Site of Sampling			
		1	2	3	4
TDS (mg/L)	1000	87.33	305.50	672.00	6792.50*
TSS (mg/L)	50-100	0.67	4.10	16.30	107.95*
Temperature (°C)	22-28	19.33	27.00	27.50	27.50
pH	6-9	7.29	6.20	6.05	5.85
DO (mg/L)	4-3	16.13*	3.25	4.40	3.85
CO ₂ (mg/L)	10-15	9.24	27.72*	35.64*	23.76*
Nitrate (mg/L)	10-20	0.02	4.00	4.14	2.77
Orthophosphate (mg/L)	0,2-1	0.11	0.17	0.17	0.14
TOM (mg/L)	69.5 – 28.4	6.74	10.39	7.58	6.32
Ammonia (mg/L)	0.2-0.5	0.13	0.22	0.07	0.29
Salinity (ppt)	0–5	0.04	0.04	9.05*	8.60*

3.1 Plankton Analysis

The data from the test results of the identification of plankton species in the Porong River basin is attached in Appendix 1. There were several species that indicate light to heavily polluted waters. The plankton species found in the Porong watershed were *Mycrocystis* sp., *Anabaena* sp., *Lyngbya* sp., *Botryococcus brauni* sp., *Raphidiopsis* sp., *Lepocinclis* sp., *Polyartha trigla*. The types that were found are the *anabaena* sp. and *Mycrocystis* sp.

The measurement of the plankton density index is shown in Fig. 2. The result indicates that the highest plankton density level was at the first point of 6 cells/mL, while the lowest plankton density level was at the fourth point of 1 cell/mL. The plankton densities of the second and third sample were 3 and 2 cells/mL, respectively.




Fig. 2: Graph of the calculation of the plankton density index in the waters of the Porong River.

The Plankton Dominance Index, as shown in Fig. 3, indicates that the value of the dominance index (C'index) ranged from 0.11 to 0.63. The highest dominance index value at the 4th sampling point was 0.63, and the lowest dominance index value at the 1st sampling point was only 0.11. The dominance index values at the second and third sampling points were 0.15 and 0.56, respectively.




Fig. 3: The graph of the calculation of the plankton dominance index in the waters of the Porong River.

3.2 Heavy Metals of Porong River

The result of heavy metal analysis found in the Porong River basin, which are Pb, Hg, Cd, and Cu, is presented in Table 3. Analysis of heavy metals in Porong River water indicated that Pb heavy metals have the highest value at point 3, at 0.0006 mg/L. The highest Hg concentration was obtained at point 4, at 0.0009 mg/L. The Cd were not detected in all sampling points. The highest Cu concentration was found at point 4, at 0.0064 mg/L.

Table 3. Measurement results of heavy metals in water samples in the Porong River.

Parameter	Site of Sampling			
	1	2	3	4
Pb (mg/L)	ND	ND	0.0006	0.0004
Hg (mg/L)	ND	0.0005	0.0005	0.0009
Cd (mg/L)	ND	ND	ND	ND
Cu (mg/L)	ND	0.0027	0.0043	0.0064

Note: ND (Not detected).

The analysis result of heavy metal types found in Porong River sediments, including Pb, Hg, Cd, and Cu, is presented in Table 4. The highest level Pb contamination was found in sediment point 3 of 15.2 mg/L. The highest level of Hg was found in sediment point 4 of 0.109 mg/L. The highest Cd was found in the sediment point 4 of 0.203 mg/L. The highest Cu contamination was found in sediment point 4 of 67.5 mg/L. In this study, every sampling location obtained high concentrations of heavy metals contamination which exceeds river water quality standards (PP Number 22 of 2021).

Table 4. Measurement results of heavy metals in sediment samples in the Porong River.

Parameter	Site of Sampling			
	1	2	3	4
Pb (mg/L)	ND	10.3	15.2	13.6
Hg (mg/L)	ND	0.060	0.053	0.109
Cd (mg/L)	ND	0.183	0.162	0.203
Cu (mg/L)	ND	46.2	53.5	67.5

Note: ND (Not detected).

3.3 Immune response (Interleukin-1 β)

Temperature optimization was carried out to obtain the best annealing temperature. From 4 temperature gradients obtained the best temperature of 51.9°C. In addition, the amplification and electrophoresis were conducted. The electrophoresis results of IL-1 β expression appeared in the band of 206 bp, as shown in Fig. 4 below.




Fig. 4: IL-1 β immune response detection results using PCR. St0) fish samples from upstream; St1) fish samples from Downstream location point 1; St2) fish samples from downstream location point 2; St3) fish samples from downstream location point 3; K-) negative control fish samples; M) Markers.

Based on the results of PCR amplification, it is shown that there was an IL-1 β response in all fish tested. This was thought to be due to water contamination at the entire water sampling site. Various immunological cytokines were evaluated against several pathogens, indicating that the immune status of the fish was against infection. Among these cytokines, IL-1 β and TNF α were considered pro-inflammatory cytokines that are produced in the early stages of infection³¹⁾.




Fig. 5: Fish samples used for IL-1 β by PCR assay

4. DISCUSSION

In general, the waters of the Porong River were categorized as polluted waters based on the water quality test where the results of CO₂, TDS, TSS, and salinity exceeded the threshold values that had been set based on the Government Regulation (PP) of the Republic of Indonesia Number 22 of 2021. According to Astuti and Indriatmoko³²⁾, dissolved oxygen was an indicator of pollution of the aquatic environment. High oxygen indicates good waters, while low oxygen indicates water pollution. According to Adamczyk et al.³³⁾, processes that affect primary productivity could affect the rate of absorption of biological CO₂ by autotrophs. Carbon dioxide levels that are too high in water is considered as toxic to the water environment. The waters that were located nearby human activity tend to have higher levels of CO₂ compared to those facing the ocean areas³⁴⁾. According to Pratama, et al.³⁵⁾, the salinity concentration obtained through the CTD electrode sensor and the TDS concentration resulting from the water type ratio equation system had a very strong correlation relationship because both parameters were measured through the conductivity characteristics of the waters. The high salinity in the water had a major influence on the tolerance of plankton that was able to survive in that location³⁶⁾. Several researchers have analyzed the abundance and diversity of phytoplankton as biological indicators to evaluate water quality and the level of fertility in aquatic environments^{37,38)}. Plankton is used as an indicator of water quality in various ways. The quality of water is essentially determined by the abundance of plankton to determine the fertility level of the water. Plankton has a short life cycle and reacts quickly to environmental changes, making it an indicator that provides signs in determining water quality³⁹⁾. The abundance of plankton is also used to monitor the fertility level of the water²⁴⁾.

Among the mentioned heavy metals (Cu, Hg, Cd, and Pb), the successive penetrating power from large to small in the waters of the Porong River was: Cu > Hg > Pb > Cd. On the other hand, in successive sediments, it was Cu > Pb > Cd > Hg. However, the content of heavy metals in

the water was still below the threshold value determined by the Government Regulation (PP) of the Republic of Indonesia No. 22 (2021). In those heavy metal-containing waters, even if the concentrations are relatively low, the heavy metals can be absorbed biologically by aquatic biota and will be involved in the immune system. This causes a process called bioaccumulation. According to its chemical-physical properties, Pb metal is not needed at all in the life process of aquatic biota. It is classified as a chemical with high toxic potential when accumulated because it has a large affinity for the protein groups of living organisms. The result of the toxicity of this metal can be physical damage (erosion, degeneration, necrosis) and can be physiological disorders (impaired enzyme function and metabolic disorders). At a higher level of biological organization, heavy metals are capable of causing biochemical, metabolic, and physiological changes in the body and inhibiting the synthesis of proteins¹⁴⁾. Previous studies have shown that metals can also cause oxidative stress by generating free radical ions. Fish tissues such as muscles, gills, liver, kidneys, and intestines have an antioxidant defense system that protects against metal-induced oxidative stress. One of the abilities is, the body can respond to heavy metal bioaccumulation in aquatic organisms with an immune response that includes IL-1 β . It is involved in cell proliferation, differentiation, inflammation, apoptosis, also stimulates various environmental challenges in the fish immune system. Although more and more studies are focusing on heavy metal-induced immunotoxicity, fish reports are still very limited. These data indicate that the four metals individually may affect expression or function of IL-1 β , thereby reducing the level of immunotoxicity in fish. IL-1 β is a key early inflammatory cytokine, which enables organisms to respond to stress caused by the microbial invasion, tissue damage, or xenobiotics. It can activate macrophages, natural-killer cells (NK cells), lymphocytes and produce various effects that cause inflammation in various fish species⁴⁰⁾. The results of our study showed the expression of IL-1 β in the kidney organs of fish due to exposure to heavy metals of water and sediments. IL-1 β play crucial roles as pro-inflammatory cytokines in regulating both innate and adaptive immunity, as well as modulating the inflammatory response. IL-1 β , being the first identified cytokine, is secreted by various immunological cells, including endothelial cells, macrophages, T lymphocytes, and other immune cells, upon stimulation by different pathogens^{31,41)}.

5. CONCLUSION

Based on the water quality and heavy metals measurement results, the parameters of CO₂, TDS, TSS, and salinity in the waters of the Porong River exceeded the threshold values set by the Government Regulation (PP) of the Republic of Indonesia Number 22 of 2021, indicating a high level of water pollution. Among the analyzed metals, Cu, Hg, Cd, and Pb exhibited a

successive penetrating power in the following order in the waters of the Porong River: Cu > Hg > Pb > Cd. In sediments, the order was Cu > Pb > Cd > Hg. The contamination of these heavy metals has significantly increased the expression of IL-1 β in fish inhabiting the Porong River. In the future, this finding can be adopted for the consideration of stakeholders in order to make regulations to reduce water pollution in the Porong River.

Acknowledgments

Acknowledgments to the Department of Research, Technology and Community Service, Directorate General of Higher Education, Research and Technology, Ministry of Education, Culture, Research, and Technology 2022 through the "National Competitive Basic Research (PDKN) 2022" [grant number: 087/E5/PG/02.00.PT/2022]

Nomenclature

IL-1 β	Interleukin-1 β
DO	Dissolved Oxygen
COD	Chemical Oxygen Demand
BOD	Biological Oxygen Demand
ND	Not Detected

References






- 1) A. Nurry, and I.M. Anjasmara, "Kajian perubahan tutupan lahan daerah aliran sungai brantas bagian hilir menggunakan citra satelit multi temporal (studi kasus: kali porong, kabupaten sidoarjo)," *Geoid*, **10** (1) 70–74 (2014).
- 2) D. Firmansyah, "Penentuan kadar logam tembaga (cu) pada perairan dan sedimen muara sungai porong sidoarjo," *SainsTech Innov. J.*, **2** (1) 24–28 (2019).
- 3) Y. Masaki, "Characteristics of industrial wastewater discharged from industrialized provinces and specific industrial sectors in china based on the official statistical reports," *Evergreen*, **3**(2) 59-67 (2016). doi:10.5109/1800873.
- 4) F. Apriyani, "Efektivitas penyerapan logam berat hgmenggunakan kerang hijau (perna viridis) dan kerang darah (anadara granosa) pada air tandon tambak dari desa banjarpanji tanggulangun sidoarjo," (2014).
- 5) M.S. Abfertiawan, R.S. Gautama, S.B. Kusuma, and S. Notosiswoyo, "Hydrology simulation of ukud river in lati coal mine," *Evergreen*, **3**(1) 21-31 (2016). doi:10.5109/1657737.
- 6) R. Widayanti, A.D. Tohjiwa, Wijayanti, and E. Setyowati, "Functions of water bodies as mitigation of the impact of urban heat island in kampung luar batang and kampung pulo geulis," *Evergreen*, **9**(2) 484–490 (2022). doi:10.5109/4794176.
- 7) U. Suraya, "Inventarisasi dan identifikasi tumbuhan air di danau hanjalutung kota palangka raya," *Daun J. Ilm. Pertan. Dan Kehutan.*, **6** (2) 149–159 (2019).
- 8) A. Nuraeni, A. Samosir, and S. Sulistiono, "Akumulasi logam berat timbal (pb) pada hati ikan patin (pangasius djambal) di waduk saguling, jawa barat," *J. Teknol. Perikan. Dan Kelaut.*, **12** (2) 113–123 (2021).
- 9) S. Dwiki, "Development of environmental policy in indonesia regarding mining industry in comparison with the united states and australia: the lesson that can be learned," *Evergreen*, **5**(2) 50-57 (2018). doi:10.5109/1936217.
- 10) M.W. Shahzad, K. Thu, B.B. Saha, and K.C. Ng, "An emerging hybrid multi-effect adsorption desalination system," *Evergreen*, **1**(2) 30-36 (2014). doi:10.5109/1495161.
- 11) B.T. Sabarudin, and S. Kartohardjono, "The combination of coagulation-flocculation and membrane processes to minimize pollution of tofu wastewater," *Evergreen*, **7**(1) 56-60 (2020). doi:10.5109/2740942.
- 12) S. Kitjanukit, "Attitude toward bioremediation-related technology and relation with company social responsibility," *Evergreen*, **6**(3) 240-245 (2019). doi:10.5109/2349300.
- 13) M.J. Hoque, "Causes, mechanisms and outcomes of environmental degradation in bangladesh: a study in sylhet," *Evergreen*, **9**(2) 310–325 (2022). doi:10.5109/4793670.
- 14) S. Rajeshkumar, Y. Liu, J. Ma, H.Y. Duan, and X. Li, "Effects of exposure to multiple heavy metals on biochemical and histopathological alterations in common carp, cyprinus carpio l.," *Fish Shellfish Immunol.*, **70** 461–472 (2017).
- 15) N.M. Roy, S. DeWolf, A. Schutt, A. Wright, and L. Steele, "Neural alterations from lead exposure in zebrafish," *Neurotoxicol. Teratol.*, **46** 40–48 (2014).
- 16) W. Ma, L. Wang, Y. He, and Y. Yan, "Tissue-specific cadmium and metallothionein levels in freshwater crab sinopotamon henanense during acute exposure to waterborne cadmium," *Environ. Toxicol. An Int. J.*, **23** (3) 393–400 (2008).
- 17) Q. Zhang et al., "Risk assessment of polychlorinated biphenyls and heavy metals in soils of an abandoned e-waste site in china," *Environ. Pollut.*, **185** 258–265 (2014).
- 18) G. Lopez-Castejon, and D. Brough, "Understanding the mechanism of il-1 β secretion," *Cytokine Growth Factor Rev.*, **22** (4) 189–195 (2011).
- 19) K. Todoric et al., "Body mass index correlates with pollutant-induced interleukin-1 β in sputum and blood," *Ann. Allergy, Asthma Immunol.*, **114** (3) 251–253 (2015).
- 20) R. Saleh, W.E. Setiawan, D. Indriani, F. Rahman, and K. Khalid, "Ecocide: Memutus Impunitas Korporasi-Walhi Riau," Cahaya Indonesia Publisher, 2019.








- 21) M.E. El-Boshy, and R. Taha, "Effects of mercuric chloride on the immunological, hematological, biochemical parameters and diseases resistance of Nile tilapia challenged with *Aeromonas hydrophila*," *Nat. Sci.*, **9** (12) 7–15 (2011).
- 22) O. Takeuchi, and S. Akira, "Pattern recognition receptors and inflammation," *Cell*, **140** (6) 805–820 (2010).
- 23) G. Jacobson et al., "Omics and cytokine discovery in fish: presenting the yellowtail kingfish (*Seriola lalandi*) as a case study," *Dev. Comp. Immunol.*, **75** 63–76 (2017).
- 24) I. Nurruhwati, Z. Zahidah, and A. Sahidin, "Kelimpahan plankton di waduk Cirata provinsi Jawa Barat," *Akuatika Indones.*, **2** (2) 102–108 (2017).
- 25) M. Sachlan, "Planktonologi," *Corresp. Course Centre. Direktorat Jenderal Perikanan, Dep. Pertanian, Jakarta*, **141** (1982).
- 26) C.C. Davis, and C.C. Davis, "The marine and freshwater plankton," Michigan State University Press, 1955.
- 27) American Public Health Association (APHA), "GENERAL INFORMATION," in: Stand. Methods Exam. Water Wastewater, American Public Health Association, 2017. doi:doi:10.2105/SMWW.2882.219.
- 28) D. Mardhia, and V. Abdullah, "Studi analisis kualitas air sungai Brangbiji Sumbawa Besar," *J. Biol. Trop.*, **18** (2) 182–189 (2018).
- 29) D. Rosarina, and E.K. Laksanawati, "Studi kualitas air sungai Cisdane Kota Tangerang ditinjau dari parameter fisika," *J. Redoks*, **3** (2) 38–43 (2018).
- 30) F. Ustaoglu, Y. Tepe, and B. Tas, "Assessment of stream quality and health risk in a subtropical Turkey river system: a combined approach using statistical analysis and water quality index," *Ecol. Indic.*, **113** 105815 (2020).
- 31) N.A. Younis, S.E. Laban, A.K. Al-Mokaddem, and M.M. Attia, "Immunological status and histopathological appraisal of farmed *Oreochromis niloticus* exposed to parasitic infections and heavy metal toxicity," *Aquac. Int.*, **28** (6) 2247–2262 (2020).
- 32) L.P. Astuti, and I. Indriatmoko, "Kemampuan beberapa tumbuhan air dalam menurunkan pencemaran bahan organik dan fosfat untuk memperbaiki kualitas air," *J. Teknol. Lingkung.*, **19** (2) 183–190 (2018).
- 33) A. Kruk et al., "Effect of co deposition on oxidation behavior and electrical properties of ferritic steel for solid oxide fuel cell interconnects," *Thin Solid Films*, **590** 184–192 (2015).
- 34) L.I. Bernawis et al., "Estimation of sea-air CO₂ exchange at Simulue Sea during summer Asian monsoon," *J. Ilmu Dan Teknol. Kelaut. Trop.*, **11** (3) 713–720 (2019).
- 35) G.I.P. Pratama, I.G. Hendrawana, I.W.G.A. Karanga, and A. Chappuis, "Karakteristik vertikal salinitas dan tds di perairan Amed dan Tulamben, Karangasem, Bali," *J. Mar. Res. Technol.*, **3** (1) 47–58 (2020).
- 36) B.K. Yudhatama, S. Redjeki, and C.A. Suryono, "Distribusi horizontal zooplankton berdasarkan salinitas di perairan Bonang Kabupaten Demak Indonesia," *J. Mar. Res.*, **8** (3) 322–327 (2019).
- 37) Z.H. Yusuf, "Phytoplankton as bioindicators of water quality in Nasarawa Reservoir, Katsina State Nigeria," *Acta Limnol. Bras.*, **32** (2020).
- 38) V. Kamboj, N. Kamboj, A.K. Sharma, and A. Bisht, "Phytoplankton communities as bio-indicators of water quality in a mining-affected area of the River Ganga, Haridwar, India," *Energy, Ecol. Environ.*, **7** (4) 425–438 (2022).
- 39) S. Rashad, M. Abdul Moneem, G. A. El-Chaghaby, and S. F. Abdel-Kader, "Monitoring water quality and plankton distribution in the River Nile around El-Maadi area in Egypt during 2018," *Egypt. J. Aquat. Biol. Fish.*, **23** (5 (Special Issue)) 81–87 (2019).
- 40) R. Sivangala, and G. Sumanlatha, "Cytokines that mediate and regulate immune responses," *Innov. Immunol.*, 1–26 (2015).
- 41) M. Shahjahan et al., "Effects of heavy metals on fish physiology—a review," *Chemosphere*, 134519 (2022).
- 42) N.M. Sidik, J. Lim, C. Yin, K. Jusoh, and M. Idris, "Metal binding protein in *Anabaena flos-aquae* as heavy metal biosensor," in: Asian Conf. Sensors, 2003. AsiaSense 2003., IEEE, 2003: pp. 241–245.
- 43) A.S.R. GP, S. Elumalai, T. Sangeetha, and D. Roop Singh, "Botryococcus braunii as a phycoremediation tool for the domestic waste water recycling from Cooum River, Chennai, India," *J. Bioremed Biodeg.*, **6** (294) 2 (2015).
- 44) J. Huisman et al., "Cyanobacterial blooms," *Nat. Rev. Microbiol.*, **16** (8) 471–483 (2018).
- 45) V. Frišták, H.D. Laughinghouse IV, and S.M. Bell, "The use of biochar and pyrolysed materials to improve water quality through microcystin sorption separation," *Water*, **12** (10) 2871 (2020).
- 46) P.J. Oberholster, Y. Schoeman, J.C. Truter, and A.-M. Botha, "Using periphyton assemblage and water quality variables to assess the ecological recovery of an ecologically engineered wetland affected by acid mine drainage after a dry spell," *Processes*, **10** (5) 877 (2022).
- 47) L.R. Hambaryan, L.G. Stepanyan, and A.S. Mamyan, "Investigations of development specifics of phytoplankton community in polluted areas of Syunik region," *Ecology*, **1** 26 (2016).
- 48) K. Patel, and K. Manoj, "Variation in epifloral community composition according to heavy metals in sediment of Tapi, Surat, India," *Int. J. Sci. Res. Biol. Sci. Vol.*, **7** (5) (2020).
- 49) X. Bai, Y. Jiang, Z. Jiang, L. Zhu, and J. Feng, "Nutrient potentiate the responses of plankton community structure and metabolites to cadmium: a microcosm study," *J. Hazard. Mater.*, **430** 128506





- (2022).
- 50) M. Solanki, and K. Manoj, "Algal biodiversity with reference to heavy metals in tapi river from surat district, gujarat," (n.d.).
 - 51) S. Dwivedi et al., "Characterization of native microalgal strains for their chromium bioaccumulation potential: phytoplankton response in polluted habitats," *J. Hazard. Mater.*, **173** (1–3) 95–101 (2010).
 - 52) B. Kiran, and K. Thanasekaran, "Metal tolerance of an indigenous cyanobacterial strain, *lyngbya putealis*," *Int. Biodeterior. Biodegradation*, **65** (8) 1128–1132 (2011).
 - 53) M.M. Areco, E. Haug, and G. Curutchet, "Studies on bioremediation of zn and acid waters using *botryococcus braunii*," *J. Environ. Chem. Eng.*, **6** (4) 3849–3859 (2018).
 - 54) P. Rzymiski, B. Poniedzialek, P. Niedzielski, P. Tabaczewski, and K. Wiktorowicz, "Cadmium and lead toxicity and bioaccumulation in *microcystis aeruginosa*," *Front. Environ. Sci. Eng.*, **8** (3) 427–432 (2014).
 - 55) J. Shen et al., "Urbanization has changed the distribution pattern of zooplankton species diversity and the structure of functional groups," *Ecol. Indic.*, **120** 106944 (2021).
 - 56) T. Tamrin, and M. Aris, "Early warning of heavy metal pollution in the waters of obi island based on plankton elements," *J. Ilm. PLATAX*, **10** (1) 55–60 (2022).
 - 57) E. Arguelles, and M.R. Martinez-Goss, "Diversity of philippine photosynthetic euglenophytes and their potential biotechnological uses: a review," *Int J Emerg Technol*, **10** (4) 24–31 (2019).
 - 58) A.T. Nguyen et al., "Phytoplankton characterization in a tropical tidal river impacted by a megacity: the case of the saigon river (southern vietnam)," *Environ. Sci. Pollut. Res.*, **29** (3) 4076–4092 (2022).
 - 59) D.M. Arias, J. García, and E. Uggetti, "Production of polymers by cyanobacteria grown in wastewater: current status, challenges and future perspectives," *N. Biotechnol.*, **55** 46–57 (2020).

Appendix



Appendix 1. Plankton Identification Results.








LOCATION	PICTURE	SPECIES	DESCRIPTION
THE UPSTREAM			
		<i>Anabaena</i> sp.	<i>Anabaena</i> sp. flos aquae as biosensor heavy metal ⁴²⁾ .
		<i>Botryococcus brauni</i>	Microalgae are unique that they grow well in wastewater and can be used to remove toxic metals and unwanted nutrients from wastewater. <i>Botryococcus brauni</i> around domestic wastewater treatment, <i>Botryococcus brauni</i> is able to decompose and eliminate COD, BOD, electrical conductivity (EC) and ammonia up to more than 70% of domestic wastewater, especially in the Adyar River (AR) domestic wastewater when compared to other wastewater, research in India ⁴³⁾
		<i>Chroococcus</i> sp.	<i>Chroococcus</i> sp. is commonly found in freshwater. Usually living in a slightly acidic to alkaline environment, this plant causes the water to turn green. They can form dense and sometimes toxic blooms The toxic can threatening ecosystem function and degrading the quality of water for recreational purposes, drinking water, fisheries, and human health ⁴⁴⁾
		<i>Microcystis</i> sp.	<i>Microcystis</i> sp. are oxygenated photosynthetic bacteria found in nutrient-rich, warm, low-turbulence fresh water ⁴⁵⁾ .
		<i>Gyrosigma acumination</i>	<i>Gyrosigma acumination</i> was a toxicant that associated with water anthropogenically impacted by industrial pollutants ⁴⁶⁾ .

	<i>Anabaena</i> sp.	<i>Anabaena</i> sp. flos aquae as biosensor heavy metal ⁴²⁾ .
	<i>Aphanothece clathrata</i>	Domination of blue-green algae, <i>Aphanothece clathrata</i> might occur by the influence of wastewaters from urban mining, metallurgical and domestic activities ⁴⁷⁾ .
	<i>Cerataulina</i> sp.	The epifloral community found by <i>Cerataulina</i> sp. associated with heavy metals (Cu, Zn, Ni, Pb, and Cd) in sediments at several river locations ⁴⁸⁾ .
	<i>Trichocerca</i> sp.	In the context of zooplankton, Rotifers exhibit Cd tolerance. <i>Trichocerca</i> sp. proliferated and became dominant at the species level after Cd application, therefore LC50 values of the Rotifera species ranged from 90 to 39,000 g/L ⁴⁹⁾ .
	<i>Pleurosigma</i> sp.	<i>Pleurosigma</i> sp., was reported which are tolerating the heavy metal pollution and also indicated the organic pollution found in water ⁵⁰⁾ .
	<i>Oscillatoria tenuis</i>	The species <i>Oscillatoria tenuis</i> mostly grows in the surface waters of the Ganges River, which are polluted by fly ash leaching ⁵¹⁾ .
	<i>Lyngbya</i>	One of the Cyanobacteria species, <i>Lyngbya</i> sp can increase the accumulation of starch and carbohydrates in the presence of metals (copper or cobalt) ⁵²⁾ .

	<i>Botryococcus braunii</i>	<i>B. braunii</i> can grow in contaminated leachate obtained from an anaerobic process in decomposing sediments from polluted rivers while successfully remediating the presence of zinc, copper, nickel, iron, and nitrate ⁵³).
	<i>Microcystis aeruginosa</i>	<i>M. aeruginosa</i> has the potential to survive in highly metal polluted environments, become a significant source of toxic metals in the food chain, and thereby contribute to increasing heavy metal toxicity to living organisms, including humans ⁵⁴).
	<i>Polyarthra trigla</i>	The suburban waters mainly found communities similar to those found in the oligo-infested waters, namely <i>Polyarthra</i> . <i>Polyarthra</i> belongs to a very closely related and highly supported group of copepods ⁵⁵).
	<i>Sulculeolaria monoica</i>	There are 17 phytoplankton genera in two separate classes, namely Diatom (<i>Sulculeolaria monoica</i>) and Dinoflagellates, and three zooplankton genera in the zooplankton class that have been identified in plankton samples from rivers ⁵⁶).

THE DOWNSTREAM

First Location	<i>Lepocinclis</i> sp	<i>Lepocinclis</i> can thrive in water bodies with high concentrations of organic matter, such as municipal sewage, sewage ponds and dairy runoff ⁵⁷).
	<i>Myrocystis</i> sp.	<i>M. aeruginosa</i> has the potential to survive in highly metal-contaminated environments, become a significant source of toxic metals in the food chain, and cause increased toxicity of heavy metals to live organisms, including humans ⁵⁴).
	<i>Lyngbya limnetica</i>	One of the Cyanobacteria species, Lygbya sp can increase the accumulation of starch and carbohydrates in the presence of metals (copper or cobalt) ⁵²).

Second Location		<i>Anabaena</i> sp.	<i>Anabaena flos aquae</i> as biosensor heavy metal ⁴²⁾ .
		<i>Raphidiopsis</i> sp.	The emergence and presence of potentially toxic blue-green algae (<i>Raphidiopsis</i> sp) combined with high levels of TSS indicate a potential health risk to the local population ⁵⁸⁾ .
		<i>Aphanocapsa elachista</i>	Filamentous of cyanobacteria colonial <i>Aphanocapsa</i> sp. is the most dominant genus in the waste stream, surpassing the green algae genus ⁵⁹⁾ .
		<i>Botryococcus braunii</i>	<i>Botryococcus</i> sp. grows in textile industry waste contaminated with heavy metals because of its ability to biologically remove Cr, Cu, As, and Cd ⁵³⁾ .
		<i>Lyngbya limnetica</i>	Cyanobacteria species <i>Lyngbya</i> sp increased accumulation of starch and carbohydrates in presence of metal (copper or cobalt) ⁵²⁾ .
		<i>Lyngbya limnetica</i>	One of Cyanobacteria species <i>Lyngbya</i> sp increased accumulation of starch and carbohydrates in presence of metal (copper or cobalt) ⁵²⁾ .
Third Location		<i>Mycrocystis</i> sp.	<i>Mycrocystis</i> sp. has the potential to survive in highly metal-polluted environments as it is a significant source of toxic metals in the food chain, thereby increasing the toxicity of heavy metals to living organisms, including humans ⁵⁴⁾ .