Relationship between the bacterial community structures on human hair and scalp

Watanabe, Kota

Laboratory of Soil and Environmental Microbiology, Division of Systems Bioengineering, Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School, Kyushu University

Yamada, Azusa Laboratory of Soil and Environmental Microbiology, Division of Systems Bioengineering, Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School, Kyushu University

Nishi, Yuri Laboratory of Soil and Environmental Microbiology, Division of Systems Bioengineering, Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School, Kyushu University

Tashiro, Yukihiro Laboratory of Soil and Environmental Microbiology, Division of Systems Bioengineering, Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School, Kyushu University

他

https://hdl.handle.net/2324/7174510

出版情報:Bioscience, Biotechnology, and Biochemistry. 84 (12), pp.2585-2596, 2020-12-01. Oxford University Press バージョン: 権利関係:

1	Relationship between the bacterial community structures on human hair
2	and scalp
3	
4	Kota Watanabe ^a , Azusa Yamada ^a , Yuri Nishi ^a , Yukihoro Tashiro ^{a,b*} , Kenji
5	Sakai ^{a,b*}
6	
7	^a Laboratory of Soil and Environmental Microbiology, Division of Systems Bioengineering,
8	Department of Bioscience and Biotechnology, Faculty of Agriculture, Graduate School,
9	Kyushu University, Motooka 744, Nishi-ku, Fukuoka 819-0395, Japan; and ^b Laboratory of
10	Microbial Environmental Protection, Tropical Microbiology Unit, Center for International
11	Education and Research of Agriculture, Faculty of Agriculture, Kyushu University, Motooka
12	744, Nishi-ku, Fukuoka 819-0395, Japan
13	
14	*Corresponding author. E-mail: kensak@agr.kyushu-u.ac.jp; TEL: +81-92-802-4738
15	
16	Acknowledgement
17	This work was supported by JSPS under the Grant-in-Aid for JSPS Research Fellows (Grant
18	No. 20J12699).
19	

20	
21	Relationship between the bacterial community structures on human hair
22	and scalp
23	
24	Abstract
25	In this study, we investigated and compared characteristics of the bacterial
26	community structures on hair (scalp hair) and scalp in 18 individuals. Significant
27	differences were found between the sites, in terms of cell density, alpha and beta
28	diversity, and relative abundance of the phyla, Firmicutes and Proteobacteria, whereas
29	no difference was found in relative abundance of the phylum Actinobacteria. Bacteria
30	of the genus Cutibacterium showed similar relative abundance at both sites, whereas
31	those of genus Pseudomonas were highly abundant on hair, and those of genus
32	Staphylococcus were significantly lesser in abundance on hair than on scalp. Statistical
33	correlations between the sites were high for the individual relative abundance of five
34	major operational taxonomic units (OTUs). This suggests that the bacterial community
35	structure on hair is composed of hair-specific genus, Pseudomonas, and skin-derived
36	genera, Cutibacterium and Staphylococcus, and is distinguishable from other human
37	skin microbiomes.

38

Key words: human scalp hair; bacterial community structure; quantitative PCR; NGS

39

Introduction 40

41	The entire human body surface, including hair and skin, is colonized by a wide variety of
42	microorganisms, including bacteria, fungi, and viruses (1-3). Human skin, the largest organ of
43	the human body, is colonized by 10^2 to 10^7 bacteria per cm ² (4). Some of these bacteria live in
44	a symbiotic relationship with their host, and protect against invasion by pathogenic
45	microorganisms (5, 6). The physical and chemical features of various parts of the skin form
46	unique bacterial community structures, that are adapted to the niche that they inhabit (7).
47	Colonization by bacteria is dependent on the physiology of the skin site, with specific bacteria
48	being associated with their microenvironments (8).
49	The scalp surface also provides a distinct microenvironment to the microbiome,
50	primarily arising from the host skin's physiological conditions, which include sebum content,
51	moisture, pH, and topography of the hair. In general, skin sites are roughly classified into
52	three groups, based on their microenvironments, which are: moist, dry, and sebaceous groups.
53	Among these skin site groups, scalp belongs to the sebaceous group. Sebaceous glands of the
54	scalp produce a large amount of oily sebum (9). Furthermore, several studies on scalp
55	microbiome in various countries have revealed the association of dandruff with bacterial
56	community structure (10-13). These studies showed that the major bacterial genera on the
57	scalp are Cutibacterium (Propionibacterium) and Staphylococcus. When compared with

58	normal scalp, the scalp with dandruff had a decreased population of Cutibacterium and an
59	increased population of Staphylococcus. As dandruff is one of the disorders of the scalp, it is
60	also a problem of the hair. Despite these facts, the relationship between the microbiomes on
61	human hair (scalp hair) and scalp is yet unknown, and studies to elucidate this relationship
62	and would be essential for a better understanding of both hair and scalp health.
63	Several studies have reported observations on the bacteria present on hair, using
64	fluorescence light microscopy or scanning electron microscopy (SEM) (14, 15). Existence of
65	bacteria was reported not only on hair shafts but also on hair follicles (16). Analysis of
66	bacterial community structures on hair by terminal restriction fragment length polymorphism
67	(T-RFLP) (17) indicated that these bacterial community structures are specific for each
68	individual, and when collected from any human body part including hand (18), could be a tool
69	for identifying a suspect. Bacterial community structures on specific parts of hair shaft and
70	hair root were further compared by 16S rRNA gene amplicon sequencing analysis, and it was
71	reported that the bacteria on hair shaft are indigenous and derived from the hair root, having
72	similar number and structure from the top to the base parts of a long hair shaft (19). The
73	report also suggested that bacterial community structures on scalp hair are distinct from those
74	on other body sites, including various parts of the skin (20). However, the differences of
75	bacterial community structures found on scalp hair and scalp were not studied, which would
76	be essential for a better understanding of hair and scalp health, as well as bacterial ecology of
77	hair.

In this study, we analyzed bacterial community structures, by 16S rRNA gene amplicon sequencing, to elucidate characteristics of the structures and to analyze the relationship between bacterial community structures of human scalp hair and human scalp within the same individual.

83 Materials and methods

This research was performed with permission from the research ethics committee of the
Graduate School of Bioscience and Biotechnology at Kyushu University.

87 Samples and collection

86

88 Hair shaft samples and scalp swab samples were collected from 18 healthy Japanese and Chinese adults of both sexes (9 males and 9 females), ranging in age from 21 to 62 years, who 89 90 consented to take part in this study (Supplemental Table 1). None of the volunteers were 91 taking any medication during the experimental period. All of the volunteers washed their hair 92 6 hours prior to sample collection. Also, samples at both sites were collected on the same day. Samples of hair shafts and scalp swabs were collected using nitrile gloves. Scalp swab 93 94 samples were directly taken from the crown of the head using cotton swabs (Mentip for 95 hospital, Nihon Menbou Corporation, Saitama, Japan) pre-moistened with 50 µL of sterile 96 distilled water. Cotton swabs were rubbed onto the scalp surface (between the hair strands) to

- 97 cover a total surface area of 2.5 cm^2 . The head of each swab was cut from the handle and
- 98 placed into an Eppendorf tube. Samples of hair shaft were cut using sterilized scissors, and
- 99 chopped into pieces of 5 mm length with the scissors before use.
- 100

101 Extraction of bacterial DNA from hair shaft and scalp swab samples

- 102 Bacterial DNA was extracted using the NucleoSpin[®] Tissue kit (MACHEREY-NAGEL,
- 103 Düren, Germany) according to the manufacturer's instructions, with a slight modification.
- 104 First, samples of scalp swabs and hair shafts were immersed in 100 µL of lysozyme solution
- 105 (20 mg/mL lysozyme derived from egg white [Wako Pure Chemical Industries, Osaka, Japan]
- in 20 mM Tris-HCl and 0.2 mM EDTA, pH 8.0) for 30 min at 37 °C, as previously reported
- 107 (19), and the DNA extracts obtained (100 μ L) were stored at -20 °C until use.
- 108

109 Quantification of the bacterial cell number on hair and scalp by qPCR

- 110 The bacterial cell number on the hair and scalp of 18 volunteers was quantified by estimation
- 111 of 16S rRNA gene copy number using real-time PCR (CFX Connect[™] System, BIO-RAD
- 112 Laboratories, Inc., CA, USA) with universal primers for a portion of the bacterial 16S rRNA
- 113 gene. We have previously shown that the values estimated by quantitative PCR (qPCR)
- 114 correspond well with the values obtained by the direct SEM observation (19).

115	Each 10 μ L reaction mixture consisted of 2 μ L of KOD SYBR® qPCR Mix
116	(TOYOBO Co., Ltd., Osaka, Japan), 0.1 μ L of each primer [357F (5'-CCT ACG GGA GGC
117	AGC AG-3') (21) and 518R (5'-ATT ACC GCG GCT GCT GG-3') (22)], and 2 μL of
118	bacterial DNA. The amplification program included an initial denaturation step at 95 °C for 5
119	min, followed by 40 cycles each of denaturation at 95 °C for 5 s, annealing at 64 °C for 20 s,
120	and elongation at 72 °C for 20 s. DNA extract from <i>Escherichia coli</i> DH5a was used as a
121	standard to generate a calibration curve. After amplification, the copy numbers of the 16S
122	rRNA genes per hair sample were calculated per cm of hair and converted to per cm ² of hair.
123	For the calculation, the following equation was used:
124	Cells/cm ² = qPCR copies / hair length (cm) × hair diameter (cm)× π
125	The diameter of the hair was measured using a stereo microscope (Stemi 305,
126	ZEISS, Oberkochen, Germany).
127	
128	Analysis of the bacterial community structures on scalp and hair by 16S rRNA gene
129	sequencing
130	To analyze the bacterial community structures of scalp and hair from 18 volunteers using the
131	MiSeq TM platform (Illumina Inc., CA, USA), a three-step PCR method was performed using
132	the extracted DNA samples. In the first-step PCR amplification, a universal primer set for the
133	V4 region of the bacterial 16S rRNA gene (515F, 5'-GTG CCA GCM GCC GCG GTA A-3'

134	and 806R, 5'-GGA CTA CHV GGG TWT CTA AT-3') (23) was used. The 25 μL reaction
135	mixture consisted of 12.5 μ L of Kapa HiFi HotStart Ready Mix (Kapa Biosystems Inc.,
136	Wilmington, MA, USA), 0.5 μ L of each primer (10 pM), and 11.5 μ L of extracted bacterial
137	DNA. The amplification program included an initial denaturation step at 95 °C for 3 min,
138	followed by 40 cycles each of denaturation at 98 °C for 30 s, annealing at 56 °C for 30 s, and
139	elongation at 72 °C for 30 s. After electrophoresis through a 1.5% (w/v) agarose gel, the
140	targeted bands were excised from the gel with sterilized cutters, and the DNA was extracted
141	using the FastGene® Gel/PCR Extraction Kit (NIPPON Genetics Co., Ltd., Tokyo, Japan),
142	according to the manufacturer's instructions. The DNA concentration was measured using a
143	NanoDrop [™] 2000 spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA, USA).
144	In preparation of 16S rRNA amplicon sequencing with MiSeq, the templates are given tail,
145	adapter, and index sequences in a two-step PCR. Therefore, long-tailed primers are required
146	for the preparation, which makes the amplification difficult. We were unable to perform direct
147	amplification in the two-step PCR, probably because the amount of bacterial DNA obtained
148	from a 3 cm hair shaft is very small. Therefore, we first performed PCR using the universal
149	primer set without any additional sequences. As a result, we succeeded in obtaining enough
150	template fragments with a minimum number of reaction cycles.
151	For the second-step PCR, a universal primer set for the V4 region of the bacterial

153 GCA GCG TCA GAT GTG TAT AAG AGA CAG GTG CCA GCM GCC GCG GTA A-3'

152

8

16S rRNA gene and tailed sequences for MiSeq sequencing were used (1-515F, 5'- TCG TCG

154	and 1-806R, 5'-GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GGG ACT
155	ACH VGG GTW TCT AAT-3') (24). Although it is reported that this primer set would poorly
156	amplify Propionibacterium of human skin (25), the results in this study showed good
157	amplifications of Cutibacterium acnes (previous name is Propionibacterium acnes) of not
158	only scalp but also hair using this primer set as the predominant species. The 25 μ L reaction
159	mixture consisted of 1.0 μL of each primer (5 $\mu M)$ which was heat-shocked at 95 °C for 5
160	min, 12.5 μ L of Kapa HiFi HotStart Ready Mix, 12.5 ng of DNA obtained from the first-step
161	PCR amplicon, and sterilized ultrapure water. The amplification program included an initial
162	denaturation step at 95 °C for 3 min, followed by 20 cycles of denaturation at 98 °C for 30 s,
163	annealing at 55 °C for 30 s, and elongation at 72 °C for 30 s. PCR products were purified
164	using the FastGene [®] Gel/PCR Extraction Kit according to the manufacturer's instructions.
165	For the third-step PCR, a primer set with flow cell adapter sequences, index
166	sequences, and tailed sequences was used (Forward primer, 5'-AAT GAT ACG GCG ACC
167	ACC GAG ATC TAC AC-Index sequence-TCG TCG GCA GCG TC-3' and Reverse primer,
168	5'-CAA GCA GAA GAC GGC ATA CGA GAT-Index sequence-GTC TCG TGG GCT
169	CGG-3'). The third-step PCR mixture (25 μ L) was composed of 12.5 μ L of Kapa HiFi
170	HotStart Ready Mix, 0.5 μ L of each primer (10 pM), 11.5 μ L of the second-step PCR
171	amplicon. The amplification program included an initial denaturation step at 95 °C for 3 min,
172	followed by 8 cycles of denaturation at 98 °C for 30 s, annealing at 55 °C for 30 s, and
173	elongation at 72 °C for 30 s. After electrophoresis in a 1.5% (w/v) agarose gel, the target

174 bands were excised with sterilized cutters, and the DNA was extracted using the FastGene® 175 Gel/PCR Extraction Kit as described above. The DNA concentration of the third-step PCR amplicons was quantified by using a Qubit[™] dsDNA HS Assay Kit (Thermo Fisher Scientific 176 177 Inc., Waltham, MA, USA) according to the manufacturer's instructions. The purified PCR 178 products from each sample were mixed, denatured, and sequenced with an MiSeq System 179 (Illumina) using MiSeq Reagent Kit v3 (300 bp ×2 cycles with pair-end; Illumina), according 180 to the manufacturer's instructions. We obtained Good's coverage values (> 95%) for all hair 181 samples using the DNA extraction kit and PCR conditions described above, indicating that 182 the results obtained contain meaningful information (Supplemental Table 2). Good's coverage 183 values were estimated using QIIMETM 1.9.1 software (26).

184

185 Bioinformatics and statistical analysis

186 The index and universal sequences of each read were checked, and reads with complete index 187 sequences were selected as valid sequences. USEARCH V8.1.1861 (27) software was used to 188 merge paired-end reads and remove chimeric sequences. After the chimera check, the reads 189 were grouped into operational taxonomic units (OTUs) at > 97% similarity. Alpha diversity 190 (observed OTUs and Shannon index) was evaluated at a 1% OTU distance using the QIIMETM software package (26). In the taxonomy-based analysis, representative sequences 191 192 for each OTU were analyzed with the EzBioCloud platform (28). Statistical analysis in 193 quantification of bacterial cell numbers was done using ANOVA, and statistical analysis in

- 194 bacterial community structure was done using Kruskal-Wallis test. Both statistical analyses
- 195 were run under XLSTAT software ver. 2014 (http://www.xlstat.com/en/).
- 196

197 Accession number

- 198 Illumina raw read sequences and the top 13 most abundant OTU sequences were deposited
- in the DDBJ/ENA/ GenBank database under accession numbers LC557428-LC557440.
- 200
- 201 **Results**
- 202 Quantification of bacterial cell number on hair and scalp
- 203 Bacterial cell number on hair and scalp from 108 samples derived from 18 volunteers was
- analyzed separately (Supplemental Table 1). The average bacterial cell number on a hair
- sample was 1.6 (\pm 1.6) × 10⁵ cells/cm², while that on a scalp sample was 3.8 (\pm 3.7) × 10⁴
- 206 cells/cm², which was lower than the number on hair by one order of magnitude (Fig. 1). In
- spite of that, analysis of correlation between these numbers, at the level of an individual,
- showed that there was statistically no correlation between the numbers on both sites
- 209 (Supplemental Fig. 1).

211 Alpha diversity of the bacterial community structures on hair and scalp

- Fig. 2 shows the two indexes of alpha diversity: observed OTUs, and the Shannon index. Both indexes were significantly higher (p < 0.0001) on hair than on scalp. It is noted that the average OTU numbers on hair and scalp were 46.2 and 19.2, respectively. On the other hand, there was weak correlation of observed OTUs, and no correlation of Shannon indexes at individual levels (Supplemental. Figs. 2A and 2B, R = 0.38, and -0.04, respectively). This suggests that the bacterial communities on hair and scalp were rather independent in alpha diversity formation.
- 219

220 Bacterial community structures on hair and scalp at phylum level

221	Three major phyla, Actinobacteria, Proteobacteria, and Firmicutes were commonly found on
222	both sites, and abundances of the three phyla added up to 97.9% and 99.4% on hair and on
223	scalp respectively (Fig. 3). The Kruskal-Wallis test revealed no statistically significant
224	difference in the abundance of Actinobacteria between the sites. On the other hand, the
225	abundance of Proteobacteria was significantly higher ($p < 0.01$) on hair (36.9 ± 20.1%) than
226	on scalp (9.5±9.4%), whereas the abundance of Firmicutes was significantly lower ($p < 0.01$)
227	on hair $(11.0 \pm 9.3\%)$ than on scalp $(33.8 \pm 12.1\%)$. These results indicate that the bacterial
228	communities on hair and on scalp had distinguishable structures at the phylum level.
229	Correlation analysis at individual level showed that abundances of Firmicutes and

230	Actinobacteria between the sites were correlated ($R = 0.58$ and 0.63 respectively), whereas
231	the abundance of Proteobacteria between the sites was not correlated ($R = 0.18$)
232	(Supplemental Fig. 3).
233	
234	Bacterial community structures on hair and scalp at OTU level
235	We extracted OTUs present in samples of all individuals, and found that 13 OTU sequences
236	existed either on hair or scalp of each individual. Of these, eight OTUs (OTU 1, 2, 3, 4, 7, 9,
237	15, 16) were present on both sites in all individuals, and the other five OTUs (OTU 5, 18, 26,
238	42, 131) were present only on hair in some individuals. The total relative abundances of the
239	eight common OTUs were 70.9% and 94.6% on hair and on scalp, respectively, being much
240	lower on hair. This observation corresponded with the results of alpha diversity analysis (Fig.
241	2).
242	These 12 OTHs were assigned to a most clearly related species (Dhylum, pairwise
242	These 13 OTOs were assigned to a most closely related species (Phylum, pairwise
243	similarity) (Fig. 4). The five major OTUs showing abundance higher than 5% were: OTU1
244	related to Cutibacterium acnes (Actinobacteria, 99.3%), OTU2 related to Staphylococcus
245	epidermidis (Firmicutes, 99.3%), OTU3 related to Lawsonella clevelandensis (Actinobacteria,
246	99.3%), OTU4 related to Pseudomonas lini (Proteobacteria, 99.3%) and OTU15 related to
247	Pseudomonas antarctica (Proteobacteria, 99.0%). Other three OTUs found in all individuals
248	were OTU 7, 9, and 16. These show rather low total abundance at 3.2% and 1.0% in total on

hair and on scalp, respectively. It is noticeable that the five OTUs (OTU 5, 18, 26, 42, 131),
found only on hair, added up to 6.0% in total abundance, and were all assigned to phylum

251 Proteobacteria.

252	Relative abundances of the five major OTUs present at both sites, and their
253	statistical correlations between hair and scalp are shown in Fig. 5 and Fig. 6, respectively.
254	OTU1 related to Cutibacterium acnes was the most predominant species on both sites. Its
255	relative abundance was not significantly different between the sites (Fig. 5A). On the other
256	hand, its individual abundance was statistically correlated between the two sites ($R = 0.66$)
257	(Fig. 6A). OTU2 related to Staphylococcus epidermidis was less abundant on average, on hair
258	than on scalp (Fig. 5B), and its individual abundance was also statistically correlated between
259	the two sites ($R = 0.69$) (Fig. 6B). OTU3 related to <i>Lawsonella clevelandensis</i> showed no
260	significantly difference in its relative abundance between the sites (Fig. 5C), while its
261	individual abundance was statistically correlated between the two sites ($R = 0.79$) (Fig. 6C).
262	On the other hand, the relative abundances of OTU4 related to <i>Pseudomonas lini</i> and OTU15
263	related to <i>Pseudomonas antarctica</i> were higher on hair than on scalp (Fig. 5D, 5E). These
264	two, OTU4 and OTU15 showed correlation between individual abundance on hair and on
265	scalp (R = 0.60, R = 0.60) (Fig. 6D, 6E). The five OTUs (OTU 5, 18, 26, 42, 131), sometimes
266	found only on hair showed relatively lower correlation between their abundance on the hair
267	and on the scalp of the same individuals (Fig. S4). These results suggest that bacterial
268	community structures on each site were partly formed with skin-resident bacteria including

269 *Cutibacterium, Staphylococcus,* and *Lawsonella*, and partly formed with some hair-specific
270 bacteria belonging to phylum Proteobacteria.

271

272 Beta diversity of the bacterial community structures on hair and scalp

273 Fig. 7A shows the beta diversity of bacterial community structures on hair and scalp samples 274 obtained by principal coordinates analysis (PCoA) based on weighted UniFrac distances. The plots were roughly grouped into two clusters on each site, although some plots were 275 276 positioned at similar areas. We further performed biplot analysis at phylum level and the 277 result corresponded well with the relative abundances of major phyla in bacterial community 278 structures on each site shown in Fig. 2 As seen in Fig. 7A, biplot of Actinobacteria was 279 located in the middle of the two clusters; biplot of Firmicutes was located in the scalp cluster. 280 On the other hand, biplot of Proteobacteria was located in the hair cluster. This clustering 281 tendency was confirmed by comparison of average weighted UniFrac distance (Fig. 7B): the 282 distances within hair and within scalp were significantly lower (p < 0.0001) than those 283 between hair and scalp. These results indicated that the bacterial community structure on hair 284 was specified by a comparably higher abundance of Proteobacteria, although major OTUs were commonly present on both sites. 285

286

287 Discussion

288 Recent studies of bacterial community structures on scalp elucidated its relationships with 289 health and disease of hair (scalp hair) and scalp (9-13). Also, relatively stable and individually 290 unique bacterial communities were found on hair (17, 19). In this study, we provide the first 291 overview of characteristics and relationship of the bacterial community structures on hair and 292 scalp using real-time PCR and 16S rRNA amplicon sequencing. As a result, it was found that 293 major bacterial species were commonly present at both sites, but the bacterial community 294 structure on hair was specific and distinguishable from that on the scalp. Important 295 relationships obtained are summarized in Table 1. Significant differences between the 296 bacterial community structures on hair and scalp were found in terms of cell density, alpha 297 diversity, and on relative abundances of Firmicutes and Proteobacteria, while no difference 298 was found in terms of relative abundance of Actinobacteria. There was correlation between 299 hair and scalp of an individual person in relative abundance of Actinobacteria and Firmicutes, 300 while there was no correlation in terms of cell density, alpha diversity, and relative abundance 301 of Proteobacteria. We will discuss each result one by one.

Average bacterial cell number on hair was significantly higher (*p*<0.05) than on scalp (Fig. 1). Bacterial cell number on hair was similar to that seen in previous reports (19, 29). On the other hand, there were no correlation between hair and scalp on individual bacterial cell number. It should be notable that there was almost no variation in the three samples per person, showing that error due to sampling method was relatively low. In spite of

307	that, taking into account that the sampling methods to collect bacterial DNA were different at
308	each site (whole extraction for hair shaft and swabbing for scalp), further structural analysis
309	was conducted by comparing relative abundances of a highlighted bacterial group, instead of
310	the cell densities. Grice et al. (30) reported that major OTUs were overlapping (97.2%) in
311	bacterial skin samples collected using three different methods: swab, scrape, and punch
312	biopsy.

313 We showed two indexes of alpha diversity on each site: observed OTUs and 314 Shannon Index. Both indexes were significantly higher on hair than on scalp (Fig. 2). This 315 could explain why the eight OTUs, commonly present on both sites of all individuals, show 316 higher abundance on scalp (94.6%) than on hair (70.9%) (Fig. 4). These results suggest that 317 various kinds of minor bacterial species inhabited only hair. It was reported that physical 318 conditions affect the formation of skin bacterial communities, and thus, alpha diversity of 319 bacterial community structures on sebaceous skin was lower than that on dry skin (1, 12, 31). 320 In a previous study, we also reported that alpha diversity was higher on dry hair shaft than on 321 sebaceous hair root (19). Higher hydrophobicity of hair would cause its higher alpha 322 diversity, compared to a sebaceous scalp. On the other hand, there was no correlation between 323 hair and scalp in terms of the individual alpha diversity of their bacterial community 324 structures. This suggests that the bacterial community structures on the two sites are 325 independent in alpha diversity formation.

326

At phylum level, average relative abundance of Actinobacteria were similar on hair

327	and scalp ($50.0 \pm 22.5\%$ and $56.1 \pm 21.0\%$, respectively), but not similar for Firmicutes (11.0
328	\pm 9.3% and 33.8 \pm 12.1%), and Proteobacteria (36.9 \pm 20.1% and 9.5 \pm 9.4%) (Fig. 3). On the
329	contrary, correlation between hair and scalp on individual relative abundances was found both
330	in Actinobacteria and Firmicutes, but not in Proteobacteria. For more detailed analysis of
331	bacterial community structure, we assigned the OTU sequences to the most closely related
332	bacterial species. Four genera - Cutibacterium, Staphylococcus, Lawsonella, and
333	Pseudomonas were present with more than 5% abundance in both sites for every sample (Fig.
334	4). These bacterial genera are also reported to be present in other human body sites.
335	Cutibacterium is the major bacterial genus present in areas with sebaceous glands, such as the
336	forehead (32). <i>Staphylococcus</i> widely inhabits various parts of human skin and the nasal area
337	(33). In case of Lawsonella, the genus has been isolated from various kinds of human
338	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is
338 339	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i> , were also found other studies, as
338 339 340	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i> , were also found other studies, as you indicated. These bacteria protect the host from pathogenic bacteria (32). They are
338339340341	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i> , were also found other studies, as you indicated. These bacteria protect the host from pathogenic bacteria (32). They are considered to proliferate using sebum and sweat as nutrient, and playing role of barrier
 338 339 340 341 342 	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i> , were also found other studies, as you indicated. These bacteria protect the host from pathogenic bacteria (32). They are considered to proliferate using sebum and sweat as nutrient, and playing role of barrier function on hair and scalp against pathogenic bacteria (32). On the other hand, the genus
 338 339 340 341 342 343 	 abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i>, were also found other studies, as you indicated. These bacteria protect the host from pathogenic bacteria (32). They are considered to proliferate using sebum and sweat as nutrient, and playing role of barrier function on hair and scalp against pathogenic bacteria (32). On the other hand, the genus <i>Proteobacteria</i> generally inhabits the natural environment such as soil and river water, and
 338 339 340 341 342 343 344 	abscesses (34). Some of the common OTUs, Cutibacterium acnes (previous name isPropionibacterium acnes) and Staphylococcus epidermidis, were also found other studies, asyou indicated. These bacteria protect the host from pathogenic bacteria (32). They areconsidered to proliferate using sebum and sweat as nutrient, and playing role of barrierfunction on hair and scalp against pathogenic bacteria (32). On the other hand, the genusProteobacteria generally inhabits the natural environment such as soil and river water, andhas been recognized as being transient on human skin (35). Correlations between hair and
 338 339 340 341 342 343 344 345 	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i> , were also found other studies, as you indicated. These bacteria protect the host from pathogenic bacteria (32). They are considered to proliferate using sebum and sweat as nutrient, and playing role of barrier function on hair and scalp against pathogenic bacteria (32). On the other hand, the genus <i>Proteobacteria</i> generally inhabits the natural environment such as soil and river water, and has been recognized as being transient on human skin (35). Correlations between hair and scalp were found for individual relative abundances of five major OTUs (Fig. 6) but not in
 338 339 340 341 342 343 344 345 346 	abscesses (34). Some of the common OTUs, <i>Cutibacterium acnes</i> (previous name is <i>Propionibacterium acnes</i>) and <i>Staphylococcus epidermidis</i> , were also found other studies, as you indicated. These bacteria protect the host from pathogenic bacteria (32). They are considered to proliferate using sebum and sweat as nutrient, and playing role of barrier function on hair and scalp against pathogenic bacteria (32). On the other hand, the genus <i>Proteobacteria</i> generally inhabits the natural environment such as soil and river water, and has been recognized as being transient on human skin (35). Correlations between hair and scalp were found for individual relative abundances of five major OTUs (Fig. 6) but not in some secondary OTUs related to Proteobacteria including <i>Pseudomonas</i> (Fig. S4). Our results

348	transient habitation on hair. In a previous study, we reported that major bacterial OTUs
349	including Pseudomonas were common on hair shaft and hair root, and it was possible that
350	these bacteria on hair shaft were indigenous and not transient due to circumstances (19).
351	Cutibacterium was the most abundant bacterial genus on hair, even though the physical
352	condition of hair seems dry and different from other sebaceous human skin zones.
353	Cutibacterium acnes carries the genes for biosynthesis of biotin, which is an essential nutrient
354	for hair growth and scalp health (11, 36, 37). On the other hand, Pseudomonas, the second
355	major bacterial genus present on hair, was not identified in a key role. Further detailed studies
356	are required to clarify the bacterial ecology of hair.
357	Finally, we analyzed beta diversity and biplots at phylum level, to compare
358	bacterial community structures between hair and scalp (Fig. 7A). It was confirmed that
359	phylum Proteobacteria (to which Pseudomonas lini and Pseudomonas antarctica belong)
360	contributed to the formation of bacterial community structure specific for hair, and phylum
361	Firmicutes (to which Staphylococcus epidermidis belongs) contributed to that on scalp.
362	Although some reports analyzed bacterial community structures on scalp (9-13), or on hair (1,
363	19, 38), there was no study on the correlation between these sites at the same individual level.
364	Also, Klerk et al. (39) evaluated bacterial adherence and colonization on hair by SEM
365	observation and reported that Pseudomonas aeruginosa (belonging to Proteobacteria) adhered
366	to and colonized hair surfaces, while Staphylococcus epidermidis (belonging to Firmicutes)
367	showed inhibited growth. This inhibition of S. epidermidis may be attributable to hair-derived

368	antimicrobial proteins or peptides (40). Therefore, it was suggested that hair contained
369	specific bacterial community structures, in that the relative abundances of Pseudomonas
370	(Proteobacteria) was higher on hair than on other skin sites.
371	In conclusion, it was suggested that bacterial community structure on hair was
372	formed from both hair-specific and skin-derived bacteria, which were different from other
373	skin microbiomes, including forearms, nostrils, and forehead (20). In particular, <i>Pseudomonas</i>
374	was one of the most hair-specific bacteria, but the genus has been recognized as being
375	transient on skin and not much attention has been paid to it. Further studies on hair-specific
376	bacteria, particularly genus <i>Pseudomonas</i> would clarify their roles and interaction with hair.
377	

378 Acknowledgements

379 The authors would like to acknowledge Center for Advanced Instrumental and Educational Supports,

380 Faculty of Agriculture, Kyushu University for their support to use MiSeq. This work was supported by

381 JSPS under the Grant-in-Aid for JSPS Research Fellows (Grant No. 20J12699).

382

383 Author contribution

384 Kota Watanabe processed the experimental data, performed the analysis and wrote the manuscript.

385 Azusa Yamada and Yuri Nishi carried out the part of experiments. Yukihiro Tashiro and Kenji Sakai

386	supervised the p	roject from ex	perimental des	ign to submissi	ion of the manus	cript. All authors agree
				-0		

and approved the manuscript to be published.

389	References

- 390 [1] Costello EK, Lauber CL, Hamady M, et al. Bacterial community variation in human body
- habitats across space and time. Science. 2009;326:1694-1697.
- 392 [2] Findley K, Oh J, Yang J, et al. Topographic diversity of fungal and bacterial communities
- in human skin. Nature. 2013;498:367-370.
- 394 [3] Wylie KM, Mihindukulasuriya KA, Zhou Y, et al. Metagenomic analysis of double-
- 395 stranded DNA viruses in healthy adults. BMC Biol. 2014;12:71.
- 396 [4] Fredricks DN. Microbial ecology of human skin in health and disease. J. Invest. Dermatol.
- 397 Symp. Proc. Proc. 2001;6:167-169.
- 398 [5] Montes LF, Wilborn WH. Anatomical location of normal skin flora. Arch derm.
- 399 1970;101:145-159.
- 400 [6] Lange-Asschenfeldt B, Marenbach D, Lang C, et al. Distribution of bacteria in the
- 401 epidermal layers and hair follicles of the human skin. Skin Pharmacol Physiol.
- 402 2011;24:305-311.
- 403 [7] Oh J, Byrd AL, Deming C, et al. Biogeography and individuality shape function in the
- 404 human skin metagenome. Nature. 2014;514:59-64.

405	[8] Kong HH, Oh J, Deming C, et al. Temporal shifts in the skin microbiome associated with
406	disease flares and treatment in children with atopic dermatitis. Genome Res.
407	2012;22:850-859.
408	[9] Clavaud C, Jourdain R, Bar-hen A, et al. Dandruff is associated with disequilibrium in the
409	proportion of the major bacterial and fungal populations colonizing the scalp. Plos
410	One. 2013;8: e58203.
411	[10] Wang L, Clavaud C, Bar-Hen A, et al. Characterization of the major bacterial-fungal
412	populations colonizing dandruff scalps in Shanghai, China, shows microbial
413	disequilibrium. Exp Dermatol. 2015;24:381-400.
414	[11] Saxena R, Mittal P, Clavaud C, et al. Comparison of Healthy and Dandruff Scalp
415	Microbiome Reveals the Role of Commensals in Scalp Health. Front Cell Infect
416	Microbiol. 2018;8:346.
417	[12] Xu Z, Wang Z, Yuan C, et al. Dandruff is associated with the conjoined interactions
418	between host and microorganisms. Sci. Rep. 2016;6:24877.
419	[13] Park T, Kim HJ, Myeong NR, et al. Collapse of human scalp microbiome network in
420	dandruff and seborrheic dermatitis. Exp Dermatol. 2017;26:835-838.
421	[14] Mase K, Hasegawa T, Horii T, et al. Firm adherence of Staphylococcus aureus and
422	Staphylococcus epidermidis to human hair and effect of detergent treatment.
423	Microbiol. Immunol. 2000:44:653-656.
424	[15] Yun A, Yang EJ, Lee YM, et al. Quantitative and qualitative estimation of bacteria
425	contaminating human hairs. J. Bacteriol. Virol. 2010;40:11-18.
	22

426	[16] Matard B, Meylheuc T, Briandet R, et al. First evidence of bacterial biofilms in the
427	anaerobe part of scalp hair follicles: a pilot comparative study in folliculitis decalvans.
428	J. Eur. Acad. Dermatol. Venereol. 2013;27:853-860.
429	[17] Nishi E, Watanabe K, Tashiro Y, et al. Terminal restriction fragment length
430	polymorphism profiling of bacterial flora derived from single human hair shafts can
431	discriminate individuals. Leg Med(Tokyo). 2017;25:75-82.
432	[18] Nishi E, Tashiro Y, Sakai K. Discrimination among individuals using terminal restriction
433	fragment length polymorphism profiling of bacteria derived from forensic evidence.
434	Int. J. Legal Med. 2015;129:425-433.
435	[19] Watanabe K, Nishi E, Tashiro Y, et al. Mode and Structure of the Bacterial Community
436	on Human Scalp Hair. Microbes Environ. 2019;34:252-259.
437	[20] Grice EA, Segre JA. The skin microbiome. Nat. Rev. Microbiol. 2011;9:244-253.
438	[21] Fierer N, Jackson J. Assessment of soil microbial community structure by use of taxon-
439	specific quantitative PCR assays. Appl. Environ. Microbiol. 2005;71:4117-4120.
440	[22] Turner S, Pryer KM, Miao VPW, et al. Investigating deep phylogenetic relationships
441	among cyanobacteria and plastids by small subunit rRNA sequence analysis. J.
442	Eukaryot. Microbiol. 1999;46:327-338.
443	[23] Caporaso JG, Lauber CL, Walters WA, et al. Global patterns of 16S rRNA diversity at a
444	depth of millions of sequences per sample. Proc. Natl. Acad. Sci. U.S.A.
445	2011;108:4516-4522.

- 446 [24] Brown CT, Hug LA, Thomas BC, et al. Unusual biology across a group comprising more
- 447 than 15% of domain bacteria. Nature. 2015;523:208-211.
- 448 [25] Kuczynski J, Lauber CL, Walters WA, et al. Experimental and analytical tools for
- 449 studying the human microbiome. Nat Rev Genet. 2011;13:47-58.
- 450 [26] Caporaso JG, Kuczynski J, Stombaugh J, et al. QIIME allows analysis of high-
- 451 throughput community sequencing data. Nat. Methods. 2010;7:33-336.
- 452 [27] Edgar RC. Search and clustering orders of magnitude faster than BLAST.
- 453 Bioinformatics. 2010;26:2460-2461.
- 454 [28] Chun J, Lee JH, Jung Y, et al. EzTaxon: a web-based tool for the identification of
- 455 prokaryotes based on 16S ribosomal RNA gene sequences. Int. J. Syst. Evol.
- 456 Microbiol. 2007;57:2259-2261
- 457 [29] Brinkac L, Clarke TH, Singh H, et al. Spatial and Environmental Variation of the Human
- 458 Hair Microbiota. Sci Rep. 2018;8:9017.
- 459 [30] Grice EA, Kong HH, Renaud G, et al. A diversity profile of the human skin microbiota.
- 460 Genome Res. 2008;18:1043-1050
- 461 [31] Gao Z, Tseng CH, Pei Z, et al. Molecular analysis of human forearm superficial skin
- 462 bacterial biota. Proc. Natl Acad. Sci. USA. 2007;104:2927-2932.
- 463 [32] Chiller K, Selkin BA, Murakawa GJ. Skin Microflora and Bacterial Infections of the
- 464 Skin. J. Investig. Dermatol. Symp. Proc. 2001;6:170-174
- 465 [33] Eiff CV, Becker K, Machka K, et al. Nasal carriage as a source of Staphylococcus aureus
- 466 bacteremia. N Engl J Med. 2001;344:11-16

467	[34] Bell ME, Bernard KA, Harrington SM, et al. Lawsonella clevelandensis gen. nov., sp.
468	nov., a new member of the suborder Corynebacterineae isolated from human
469	abscesses. Int. J. Syst. Evol. Microbiol. 2016;66:2929-2935.
470	[35] Cogen AL, Nizet V, Gallo RL. Skin microbiota: A source of disease or defence? Br. J.
471	Dermatol. 2008;158: 442-455.
472	[36] Patel DP, Swink SM, Castelo-Soccio L. A review of the use of biotin for hair loss. Skin
473	Appendage Disord. 2017;3:166-169.
474	[37] Chiu CH, Huang SH, Wang HM. A review: hair health, concerns of shampoo ingredients
475	and scalp nourishing treatments. Curr. Pharm. Biotechnol. 2015;16:1045-1052.
476	[38] Tridico SR, Murray DC, Addison J, et al. Metagenomic analyses of bacteria on human
477	hairs: a qualitative assessment for applications in forensic science. Invest. Genet.
478	2014;5:16.
479	[39] Klerk SK, Lai HY, Sze SK, et al. Bacteria display differential growth and adhesion
480	characteristics on human hair shafts. Front Microbiol. 2018;7(9):2145.
481	[40] Adav SS, Subbaiaih RS, Kerk SK, et al. Studies on the proteome of human hair-
482	identification of histones and deamidated keratins. Sci. Rep. 2018;8:1599.
483	
484	

	Significant difference between hair and scalp (p<0.05)	Hair	Scalp	Correlation between hair and scalp on individual abundance (R≧0.4)
Cells /cm ²	Yes	High 1.6(±1.6)×10 ⁵	Low 3.8(3.7) ×10 ⁴	No
Alpha diversity	6			
Observed OTUs	Yes	High 46.2(±15.9)	Low 19.2(±8.8)	No
Shannon Index	Yes	High 3.0(±0.8)	Low 1.8(±0.5)	No
Relative abundance (%)				
Actinobacteria	No	50.5(±22.5)	56.1(±21.0)	Yes
Firmicutes	Yes	Low 11.0(±9.3)	High 33.8(±12.1)	Yes
Proteobacteria	Yes	High 36.9(±20.1)	Low 9.5(±9.4)	No

485 <u>Table 1</u>: Correlation of bacterial community structures between hair and scalp

- 486 <u>Supplemental Table 1</u>: Details of hair and scalp samples in 18 volunteers and quantification
- 487 of bacterial cell number at each site by qPCR of the bacterial 16S rRNA gene
- 488

Volunteer	Gender	Аде	Hair	Hair	Cell number /	Cell number / cm ²	
Volunteer	Genuer		(mm) (µm)		of hair	of scalp	
F001	Female	22	263(±57)	89	5.7(±2.4)×10 ⁴	3.2(±2.5)×10 ⁴	
F002	Female	22	417(±31)	63.3	1.7(±0.4)×10 ⁵	1.1(±0.3)×10 ⁵	
F003	Female	21	287(±40)	88.9	5.5(±2.7)×10 ⁴	4.5(±3.1)×10 ⁴	
F004	Female	27	335(±12)	87.2	7.1(±1.9)×10 ⁴	3.4(±0.7)×10 ⁴	
F005	Female 23 332(±20) 75.6		75.6	1.1(±0.3)×10 ⁵	4.2(±1.3)×10 ⁴		
F006	Female	22	371(±27)	73.9	3.0(±0.5)×10 ⁴	5.1(±1.5)×10 ⁴	
F007	Female	41	193(±1)	88	5.9(±1.4)×10 ⁴	1.2(±0.4)×10 ⁴	
F008	Female	27	402(±19)	79.7	1.4(±0.07)×10 ⁵	4.3(±1.1)×10 ³	
F009	Female	23	236(±32)	99.7	6.0(±0.7)×10 ⁴	1.9(±0.8)×10 ⁴	
M001	Male	25	83(±2)	98.4	5.6(±2.9)×10 ⁵	2.6(±0.3)×10 ⁴	
M002	Male	26	29(±5)	70.8	4.1(±1.0)×10 ⁵	3.0(±0.7)×10 ⁴	
M003	Male	22	88(±7)	68.7	1.3(±0.6)×10 ⁵	9.2(±9.2)×10 ⁴	
M004	Male	22	46(±4)	76.6	1.6(±0.6)×10 ⁵	2.2(±0.4)×10 ⁴	
M005	Male	22	111(±10)	102	1.0(±0.1)×10 ⁵	2.6(±0.4)×10 ⁴	

M006	Male	29	74(±11)	109.8	2.6(±0.3)×10 ⁵	5.9(±2.2)×10 ⁴
M007	Male	62	65(±3)	61.8	3.5(±0.5)×10 ⁵	4.9(±1.8)×10 ⁴
M008	Male	40	72(±2)	111.4	1.0(±0.3)×10 ⁵	1.5(±1.0)×10 ⁴
M009	Male	23	65(±2)	104	2.2(±0.6)×10 ⁵	1.5(±0.1)×10 ⁴

489

409	
490	
491	
492	
493	
494	
495	
496	
497	
498	
499	
500	
501	
502	
503	
504	

505 <u>Supplemental Table 2</u>: Details of MiSeq analysis of 108 hair and scalp samples

		Hai	ir	Scalp				
Gender	Volunteer ID	Hair length (mm)	Row read number	Good's coverage value (%)	Gender	Volunteer ID	Row read number	Good's coverage value (%)
	F001	180	18,918	99.1		F001	34,119	99.5
	F001	290	7,085	99.4		F001	24,464	99.3
	F001	316	16,292	99.0		F001	13,093	99.3
	F002	373	15,218	98.8	_	F002	12,473	98.9
	F002	437	23,647	98.2	Female	F002	21,808	99.2
	F002	442	24,693	98.0		F002	26,899	98.5
Female	F003	258	4,753	98.8		F003	36,584	99.4
	F003	260	15,589	98.4		F003	24,193	99.5
	F003	343	42,776	98.7	_	F003	30,594	99.8
	F004	320	31,116	98.5		F004	22,270	99.3
	F004	335	36,188	98.8		F004	27,600	99.2
	F004	350	37,979	98.8		F004	14,740	99.5
	F005	315	25,346	97.8		F005	31,619	99.4

	F005	320	23,023	98.0		F005	19,455	99.4
	F005	360	22,073	97.9		F005	8,328	99.7
	F006	386	32,067	98.3		F006	21,400	99.5
	F006	333	23,162	97.9	-	F006	29,124	99.0
	F006	394	4,283	98.1		F006	20,064	99.4
	F007	192	17,769	99.1	-	F007	17,274	99.2
	F007	193	18,219	98.9		F007	16,555	99.5
	F007	195	28,974	98.4	-	F007	23,044	99.5
	F008	378	34,007	98.6	-	F008	28,647	99.5
	F008	405	26,627	99.3		F008	6,453	99.6
	F008	424	2,780	99.2		F008	19,118	99.8
	F009	200	21,399	97.8		F009	20,378	98.2
	F009	230	35,982	97.3		F009	8,360	99.1
	F009	277	32,199	97.8	4	F009	16,522	99.0
	M001	81	47,895	98.3		M001	30,257	99.1
	M001	82	15,069	97.7		M001	29,871	99.2
Mala	M001	85	45,660	97.6	Male	M001	33,081	98.7
iviale	M002	25	31,574	99.3		M002	17,969	99.5
	M002	26	26,218	99.4		M002	25,807	99.1
	M002	36	13,497	99.2		M002	18,835	99.5

	M003	78	7,673	98.9		M003	26,571	99.3
	M003	91	27,818	99.0		M003	23,967	99.8
	M003	95	4,179	98.0		M003	22,234	99.7
	M004	42	32,069	98.8		M004	29,714	99.4
	M004	44	9,282	98.6		M004	24,510	99.2
	M004	52	4,946	98.7	-	M004	31,295	98.7
	M005	102	38,698	98.8	-	M005	23,583	99.6
	M005	105	14,144	98.8	-	M005	42,952	99.4
	M005	125	28,906	98.6	-	M005	10,726	99.5
	M006	73	19,200	98.0		M006	24,508	99.5
	M006	61	23,549	98.3		M006	23,135	99.6
	M006	88	9,293	98.1		M006	14,694	99.3
	M007	61	24,335	98.1		M007	13,234	98.6
	M007	67	22,557	98.2	4	M007	3,388	98.9
	M007	68	23,936	98.2		M007	23,693	98.7
	M008	70	16,657	99.4		M008	35,365	99.5
	M008	72	38,767	99.2		M008	20,841	99.3
	M008	74	39,555	99.2		M008	13,439	99.3
	M009	62	26,924	99.0		M009	12,713	99.6
	 M009	65	22,408	99.0		M009	19,288	99.4
- 1	 							

						•	
	M009	67	30,353	99.1	M009	31,819	99.2
507							
508							
509							
510							
511							
512							
513							
514							
515							
516							
517							
518							
519							
520							
521							
522							
523							
524							

525 Figure legends

- 526 Figure 1 Bacterial cell number (copies/ cm^2) on hair and scalp.
- 527 Figure 2 Alpha diversity of bacterial community on the hair and scalp,
- 528 based on (A) observed OTU and (B) Shannon index. The values are obtained from clustering
- 529 of 1,000 reads per sample.
- 530 Figure 3 The average relative abundances of the major phyla in bacterial community structure
- 531 on hair and scalp. The different alphabets mean that there are significant differences at
- 532 comparison of each bacterial phyla between the sites (Kruskal-wallis test, p<0.01).
- 533 Figure 4 Phylogenetic tree and abundance heatmap of the major 13 OTUs commonly found in
- all 18 individuals.
- 535 Figure 5 Comparison of relative abundance of most major five OTUs between hair and scalp.
- 536 (A) relative abundance of OTU1 related to *Cutibacterium acnes*, (B) OTU2 related to
- 537 Staphylococcus epidermidis, (C) OTU3 related to Lawsonella clevelandensis, (D) OTU4
- 538 related to *Pseudomonas lini*, and (E) OTU15 related to *Pseudomonas antarctica*.
- 539 Figure 6 Correlation of relative abundance of most major five OTUs between hair and scalp.
- 540 (A) OTU1 related to *Cutibacterium acnes*, (B) OTU2 related to *Staphylococcus epidermidis*,
- 541 (C) OTU3 related to Lawsonella clevelandensis, (D) OTU4 related to Pseudomonas lini, and
- 542 (E) OTU15 related to *Pseudomonas antarctica*.

- 543 Figure7A Beta diversity of bacterial community structure on hair and scalp in 108 samples
- 544 from eighteen volunteers and their biplot analysis
- 545 analysis at phylum level. The PCoA plots based on the weighted UniFrac analysis are shown.
- 546 (B) Comparison of weighted Unifrac distances within and between hair and scalp samples.
- 547
- 548
- 549 Suppl Figure 1 Correlation of bacterial cell number between hair and scalp
- 550 Suppl Figure 2 Correlation of alpha diversity between hair and scalp. (A) Observed OTUs,
- 551 (B) Shannon.
- 552 Suppl Figure 3 Correlation of relative abundance of major three phylum between hair and
- 553 scalp. (A) Actinobacteria, (B) Firmicutes, and (C) Proteobacteria.
- 554 Suppl Figure 4 Correlation of relative abundance of second major OTUs between hair and
- scalp. (A) relative abundance of OTU131 related to *Pseudomonas endophytica*, (B) OTU9
- 556 related to *Pseudomonas alcaliphila*, (C) OTU18 related to *Rhodanobacter ginsengiterrae*, (D)
- 557 OTU7 related to Rahnella aquatilis, (E) OTU26 related to Escherichia coli, (F) OTU5 related
- 558 to Moraxella osloensis, (G) OTU42 related to Acinetobacter bereziniae, and (H) OTU16
- 559 related to Carnobacterium gallinarum.
- 560

- 561 Table. 1 Summary of differences in abundance and correlation of bacterial community
- 562 between hair and scalp
- 563 Suppl Table1 Details of hair and scalp samples in eighteen volunteers and quantification of
- the bacteria cell number at each site by qPCR of the bacterial 16S rRNA gene. 564
- 565 Suppl Table 2 Details of MiSeq analysis of 108 hair and scalp samples.
- 566
- 567



Fig. 1 Watanabe et al.



Fig. 2 Watanabe et al.

Fig. 3 Watanabe et al.



Kruskal-wallis test, p<0.01

Fig. 4 Watanabe et al.







URL: https://mc.manuscriptcentral.com/bbb Email: TBBB-peerreview@journals.tandf.co.uk











Fig. S1 Watanabe et al.

Fig. S2 Watanabe et al.



Bioscience, Biotechnology, and Biochemistry **Fig. S3 Watanabe et al.**





URL: https://mc.manuscriptcentral.com/bbb Email: TBBB-peerreview@journals.tandf.co.uk

