

Zeolitic Imidazole Frameworks-8 (ZIF-8) Modified with Cu(II)/Ni(II)/Co(II) As Bifunctional ORR/OER Electrocatalytic Material

Annisa Nur Buana Wati

Department of Chemistry, Faculty of Mathematics and Sciences, Universitas Sebelas Maret

Ubed Sonai Fahrudin Arrozi

Department of Chemistry, Faculty of Mathematics and Sciences, State University of Malang

Abu Masykur

Department of Chemistry, Faculty of Mathematics and Sciences, Universitas Sebelas Maret

Sri Hastuti

Department of Chemistry, Faculty of Mathematics and Sciences, Universitas Sebelas Maret

他

<https://doi.org/10.5109/7432657>

出版情報 : Evergreen. 13 (2), pp.813-824, 2026-06. Transdisciplinary Research and Education
Center for Green Technologies, Kyushu University

バージョン :

権利関係 : Creative Commons Attribution 4.0 International



Zeolitic Imidazole Frameworks-8 (ZIF-8) Modified with Cu(II)/Ni(II)/Co(II) As Bifunctional ORR/OER Electrocatalytic Material

Annisa Nur Buana Wati¹, Ubed Sonai Fahrudin Arrozi², Abu Masykur¹,
Sri Hastuti¹, Atmanto Heru Wibowo^{1,*}

¹Department of Chemistry, Faculty of Mathematics and Sciences, Universitas Sebelas Maret, Jl. Ir. Sutami No. 36A. Jebres, Surakarta 57126, Central Java, Indonesia

²Department of Chemistry, Faculty of Mathematics and Sciences, State University of Malang, Jl. Semarang 5, Malang 65145, East Java, Indonesia

*Author to whom correspondence should be addressed:

E-mail: aheruwibowo@staff.uns.ac.id

(Received February 21, 2025; Revised July 24, 2025; Accepted September 10, 2025)

Supporting Information





Fig. S1: EDX mapping of 50%-Co(II)/ZIF-8







Fig. S2: Koutecky-Levich plots determined from Figure 1a-c of the modified ZIF-8 at different electrode potential values




Fig. S3: Nitrogen physisorption plots of the modified ZIF-8