

EFFECTS OF O₂ IN AIR AND NaCl IN MEDIUM ON PHOTOSYNTHESIS AND PHOTORESPIRATION IN TWO COTTON CULTIVARS

Mert, Hasan Huseyin
Biology Department Faculty of Education DokuzeYLul University

<https://hdl.handle.net/2324/8118>

出版情報 : BIOTRONICS. 15, pp.1-7, 1986. Biotron Institute, Kyushu University
バージョン :
権利関係 :

EFFECTS OF O₂ IN AIR AND NaCl IN MEDIUM ON PHOTOSYNTHESIS AND PHOTORESPIRATION IN TWO COTTON CULTIVARS

Hasan Hüseyin MERT

Biology Department, Faculty of Education, Dokuz Eylül University, Buca-Izmir, Turkey

(Received June 30, 1986; accepted August 18, 1986)

MERT H. H. *Effects of O₂ in air and NaCl in medium on photosynthesis and photorespiration in two cotton cultivars.* BIOTRONICS 15, 1-7, 1986. The measurements of photosynthesis and photorespiration in the two cultivars of cotton plant under different salt and oxygen concentrations showed that, *Gossypium hirsutum* cv. Coker 100 A/2 was less tolerant than *G. hirsutum* cv. 2421-A. There was a decrease in the photosynthesis with an increase in the salt concentrations at both 21 and 2% oxygen concentrations, however, the inhibition was higher at the former as compared to the latter oxygen concentration. The photorespiration too was low in both the cultivars but, significantly higher in normal air (21% oxygen) than in an air of low oxygen (2%) content.

Key words: *Gossypium hirsutum* L.; cotton; oxygen; salt; sodium chloride; photosynthesis; photorespiration.

INTRODUCTION

Several investigations have been carried out on the effects of salts on the photosynthetic behaviour of plants. According to Tiku (13), in the species of *Salicornia* and *Distichlis*, the CO₂ uptake increases in all the salt concentrations, however, it was observed that, in different cultivars of cotton grown from the seeds pretreated with different concentrations of NaCl for different periods, photosynthesis of these plants decreases gradually in relation to the NaCl content of the medium (8), which could be a characteristic for the glycophytes. The findings concerning the respiratory behaviour of the plants in relation to the salt content too are contradictory. Nieman (9) has observed an increase in the respiration of the plants grown in a nutrient medium with salt. On the other hand, Boyer (3) reported that salt has an inhibitory effect on this process.

The photorespiratory behaviour of the plants in relation to the oxygen, carbon dioxide, light and temperature has also been studied by many workers, such as Forrester *et al.* (5), Bjorkman *et al.* (2), Ludvig and Calvin (7) and Laing (6), however, not much work has been done on the effects of salt on this phenomenon. From this view point, the studies were carried out on the photosynthesis and photorespiration in two cultivars of cotton, designated as a semihalophyte by Waisel (16), using different concentrations of salt and oxygen.

MATERIAL AND METHODS

The two cultivars of cotton, *Gossypium hirsutum* L., cv. Coker 100 A/2 and 2421-A were used in the present studies: the former being less tolerant to the salt and the latter more tolerant (1, 4). The seeds were sown in rockwool supplied with 0.1 percent superba nutrient solution shown in Table 1 (10).

After the plant growing for 15 days, 0.3, 0.6 and 1.0 percent solutions of NaCl were supplied to seedlings in different pots. The salt solutions were applied 3 times on different days in a week to these seedlings. The experiments were carried out in a biotron at 30°C with a photoperiod of 16 h and 70% RH. The light source was 400 W Phillips HPI/T fluorescent tubes lying at a distance of 65 cm from the pots.

Table 1. Composition and percentage distribution of elements in superba

Elements	mg per g superba
N	130
P	40
K	190
S	53
Mg	15
Fe	1
Mn	0.4
Cu	0.4
B	0.1
Zn	0.2
Mo	0.03
Co	0.01




Fig. 1. Diagram of the closed system assimilation chamber.

The experiments were performed in a closed system assimilation chamber (Fig. 1). The reaction vessel was 75 mm in diameter and 20 mm deep. The cover of the vessel was fitted with a valve combination of the quickfit MAF I/75 with five outlets through which the gas could be directed. The gas leaving the chamber passed through a column of CaCl₂ for dehydration and then through an infrared CO₂ gas analyzer model Beckman 865, calibrated with N₂ and N₂ + 378 ppm CO₂. The signal from the gas analyzer was recorded with a Watanabe milivolt recorder, and the gas returned to the reaction vessel through a glass tubing. The gas could be directed alternately by manipulating valves in to one of the three loops: leading either to a solution of 10 ml of 0.5 M KOH (1), where CO₂ was absorbed and stored for scintillation measurements; to a container of soda asbestos (2), when CO₂ free air was desired; or to a third loop connecting the system to a flask (3), where ¹⁴CO₂ was produced from NaH¹⁴CO₃. The total volume of the assimilation chamber with tube connections was 0.76 l.

Two slide projectors with quantum flux densities of 1500 μE m⁻² s⁻¹ from the top and 2500 μE m⁻² s⁻¹ from below were used for irradiation and ¹⁴CO₂ fixation and photorespiration were measured under white light. The flux densities were measured with a Li-cor-modell Li-185.

About 40 days old plants were used and the leaf discs were taken from the third youngest leaf for the measurement. Thirty leaf discs (11 mm in diameter and 0.28 dm² in the total leaf area) were floated with the upper epidermis up, on 75 ml distilled water in the reaction vessel. After placing the material in the vessel, the system was flushed for 3 min with normal air, or with N₂+CO₂ gas mixture for obtaining O₂ free air. For the treatment of 2% O₂, 15 ml of O₂ was injected to the chamber by a syringe. The measurement was performed as follows: After the measurement of dark respiration for 6 min, the CO₂ was removed from the system through the column of soda asbestos for 6 min. It was then opened to the CO₂ reservoir tube containing 1 ml unlabeled 0.02 M NaHCO₃ solution and 25 μl of NaH¹⁴CO₃ (0.587 mCi μmol⁻¹), to which 4 ml of 20% H₃PO₄ was injected just prior to opening. The ¹⁴CO₂ supply to the system was stopped after a short period of equilibrium (6 min) and the leaf discs then irradiated with white light for measurement of photosynthesis and photorespiration. Any remaining ¹⁴CO₂ was removed from the system by absorption on soda asbestos. The assimilation chamber was flushed with normal air (approx. 350 ppm CO₂) or to N₂+CO₂ gas (378 ppm CO₂). The chamber was again irradiated and the system immediately opened to the KOH column in order to recover the ¹⁴CO₂ released from the leaf discs during photorespiration. After 0, 2, 6 and 8 min, 1.5 ml samples were taken from the KOH column. The system was then cleared of all CO₂ by opening it to the soda asbestos.

The KOH samples were prepared for scintillation counting, by placing 1.0 ml in a scintillation vial and adding 10 ml toluene solution, containing 5 g PPO (2,5-Diphenyloxazole), 0.05 g POPOP (1,4-bis(2-(5-Phenyloxazolyl))benzene) and 30% triton x-100 per liter. The vials were refrigerated overnight, and radioactivity was measured with Philips scintillation counter.

The results, were tested statistically using the method by Steel and Torrie (12).

RESULTS AND DISCUSSION

Table 2 shows the rate of photosynthesis in Coker 100 A/2 as well as 2421-A. The photosynthetic rate decreased with increasing salt concentrations in both air conditions, but the decrease of photosynthesis in both cultivars at 2% O₂ is smaller than that in normal air. Although, in the case of Coker 100 A/2, the decrease rate of photosynthesis in the plants treated 1% salt as compared to that in the control plants is 34.5% in normal air, it is only 26.8% in 2% O₂. In the case of 2421-A, the decrease is 24.0% in normal air and 21.4% in 2% O₂. This fact shows that there is some relation between salt tolerance and the O₂ effect on photosynthesis. The rate of photosynthesis is higher in 2421-A cv. than in Coker 100 A/2 at 1.0% salt concentration in both air conditions, which is related to the other physiological factors of these cultivars, e.g., disintegration of chlorophyll and protein content in Coker 100 A/2. It is necessary to make further analysis to solve this problem.

Table 3 shows the photorespiration data measured as the ¹⁴CO₂ release after a period of ¹⁴C-labelling in white light. Here again, the photorespiration was significantly higher in normal air than in the air with the lower O₂-content. There

Table 2. Net photosynthetic rate ($\mu\text{g CO}_2 \text{ dm}^{-2} \text{ h}^{-1}$) of the leaf discs, illuminated with white light under different air conditions (mean of eight experiments)

Air	Salt conc. (%)	Coker 100 A/2	2421-A
Normal air	Control	4202	3898
	0.3	3451 (17.9)*	3284 (15.8)
	0.6	3091 (26.4)	3072 (21.2)
	1.0	2757 (34.4)	2962 (24.0)
2% O ₂	Control	7275	6956
	0.3	6318 (13.2)	6294 (9.5)
	0.6	6029 (17.1)	5906 (15.1)
	1.0	5324 (26.8)	5465 (21.4)

LSD (5%): 0.64 at Coker 100 A/2; 0.69 at 2421-A.

* Percent reduction of photosynthetic rate against the control.

Table 3. Photorespiration rate (CPM ¹⁴CO₂), released after a period of photosynthesis in white light in normal air and in 2% O₂ under different salt concentrations in two cotton cultivars (mean of eight experiments)

Air	Salt con. (%)	Coker 100 A/2	2421-A
Normal air	Control	1228	2041
	0.3	1129	1883
	0.6	895	1663
	1.0	559	1204
2% O ₂	Control	522	798
	0.3	450	708
	0.6	398	665
	1.0	369	583

LSD (5%): 73 at Coker 100 A/2; 80 at 2421-A.




Fig. 2. The inhibitive effect of salt concentration on the photosynthesis of the two cotton cultivars as compared to the control.

is a great difference between Coker 100 A/2 and 2421-A in air. As is apparent from the table too, rates of photorespiration decreased gradually with an increase in the salt concentration of the medium in both cultivars.

The photorespiration rate in 2421-A was higher than that in Coker 100 A/2 in each salt concentration, and both in normal air and air of 2% O₂. However, photorespiration rate in air of 2% O₂ is almost half of that in normal air.

The photosynthesis in the cotton cultivars (Coker 100 A/2 and 2421-A), when measured as CO₂ uptake in the white light, seems to be dependent on oxygen as well as the salt concentrations. The percentage decrease in photosynthesis in relation to the salt in normal air is higher than that at 2% oxygen level (Table 2). No significant effect of NaCl on the photosynthesis of salt tolerant cultivar 2421-A was found in both air conditions.

This indicates that salt effect is dependent on O₂ concentrations in these different cultivars. These findings are in full agreement with those of Aydemir (1) and Emiroğlu (4). It is a well known fact that the glycolate pathway is saturated by 2% oxygen level (15) and glycolate production during photorespiration increases with oxygen concentration up to 100 percent level (14). Therefore, the effect of oxygen on the photosynthesis seems to be limited to the glycolate pathway rather than to the activity of RuDP carboxylase/oxygenase activity. According to Shomer-Ilan and Waisel (11), in the plants provided with a NaCl solution, the leaves show high activity of PEP carboxylase, as well as a significant CO₂ fixation via C₄-pathway. However, these findings are not parallel to those of ours (Table 2), which may be due to the fact that we used C₃ plants.

The photorespiration, when measured as the $^{14}\text{CO}_2$ release in air and air of low O_2 , also shows an oxygen dependence (Table 3). These observations reveal that CO_2 release increases at higher oxygen concentrations, which stimulate glycolate production from the splitting of RuDP to PGA and P-glycolate. The production of glycolate in this way shows a linear correlation to the oxygen concentration between zero and 21 percent (14). Our results fully support the findings of Tolbert (14). The salt concentration also affected the photorespiration in the two cultivars of cotton plant (Table 3). The photorespiration rate is significantly higher in normal air than in the air of low O_2 . Our findings show that, under such a condition the effect of the salt concentrations resembles that of the O_2 concentrations, as the salt concentrations stimulate glycolate production in the two cotton cultivars. But, the mechanism of the effect of the salt concentrations on the photorespiration of the plants still needs an explanation.

ACKNOWLEDGEMENTS

Author would like to express his deepest gratitude to Prof. Dr. N. Takahashi (ARI, Tohoku University) and Prof. Dr. M. Öztürk (Ege University) for their critical reading of this manuscript.

REFERENCES

1. Aydemir M. (1970) Memleketimizde ekilen pamuk çeşitleri. *Karınca Mat. ve Tic. Koll. Sti. Izmir*. Pages 84.
2. Bjorkman O., Hiesey W. M., Nobs M., Nicholson F. and Hart R. W. (1968) Effect of oxygen concentration on dry matter production in higher plants. *Cornegia Inst. Washington Year Book* **66**, 228-232.
3. Boyer J. S. (1965) Effects of osmotic water stress on metabolic rates of cotton plants with open stomata. *Plant Physiol.* **40**, 229-234.
4. Emiroğlu S. H. (1964) Gossypol glandi taşımayan pamuk islahi ve glandsiz bazı introduksiyon pamuklarının ticari çeşitlerle karşılaştırma verim ve kaliterleri üzerinde araştırmalar. *Birlik Mat. Izmir*.
5. Forrester M. L., Krotkov G. and Nelson C. D. (1966) Effect of oxygen on photosynthesis, photorespiration in detached leaves. I. *Soybean. Plant Physiol.* **41**, 422-427.
6. Laing W. A., Ogren W. L. and Hageman R. H. (1974) Regulation of soybean net photosynthetic CO_2 fixation by the interaction of CO_2 , O_2 and ribulose 1.5 diphosphate carboxylase. *Plant Physiol.* **54**, 678-685.
7. Ludwig L. J. and Calvin D. T. (1971) The rate of photorespiration during photosynthesis and the relationship of the substrate light respiration to the products of photosynthesis in sunflower leaves. *Plant Physiol.* **48**, 712-719.
8. Mert H. H. (1982) *Gossypium hirsutum* L. (Pamuk) 'da tuza dayanıklılık mekanizması, fotosentez, fotorespirasyon ve klorofil içeriği ile ilişkili olarak incelenmesi. Habilitation Thesis, Ege Univ. Izmir.
9. Nieman R. H. (1962) Some effects of sodium chloride on growth, photosynthesis and respiration of twelve crop plants. *Bot. Gazette* **123**, 279-285.
10. Nilsen S. (1977) Seedling of Norway spruce, scots pine and lodge pole pine grown in rockwool. *Nor. Inst. for skogforskning*.
11. Shomer-Ilan and Waisel Y. (1973) The effect of sodium chloride on the balance the C_3 and C_4 -carbon fixation pathways. *Physiol. Plant.* **29**, 190-193.
12. Steel R. G. and Torrie J. H. (1960) *Principles and Procedures of Statistics*. Pages 106, McGraw-Hill, New York.

13. Tiku B. L. (1976) Effect of salinity on the photosynthesis of the halophyte *Salicornia rubra* and *Distichlis stricta*. *Physiol. Plant.* **37**, 23–28.
14. Tolbert N. E. (1973) Glycolate biosynthesis. Pages 21–50 in B. L. Horecker and E. R. Stadtman (eds) *Topics on Cellular Regulation. Vol. 7*. Academic Press, New York.
15. Viil J. and Parnik T. (1978) On the control of the glycolate pathway by light and oxygen. *Z. Pflanzenphysiol.* **88**, 219–226.
16. Waisel Y. (1972) *Biology of Halophytes*. Pages 395, Academic Press, New York.