

AN INTEGRATED STATISTICAL APPROACH TO ESTIMATING PLANT RESPONSES TO SEQUENTIAL AND CONCURRENT GASEOUS POLLUTANTS

Hale, B.

Department of Horticultural Science

Ryan, D.

Department of Mathematics and Statistics

Ormrod, D. P.

Department of Horticultural Science

Allen, O. B.

Departments of Mathematics and Statistics and Animal and Poultry Science University of Guelph

<https://hdl.handle.net/2324/8187>

出版情報 : BIOTRONICS. 22, pp.35-46, 1993-12. Biotron Institute, Kyushu University
バージョン :
権利関係 :

AN INTEGRATED STATISTICAL APPROACH TO ESTIMATING PLANT RESPONSES TO SEQUENTIAL AND CONCURRENT GASEOUS POLLUTANTS

B. HALE*, D. RYAN**, D. P. ORMROD*, and O. B. ALLEN***

*Department of Horticultural Science, **Department of Mathematics and Statistics, and

***Departments of Mathematics and Statistics and Animal and Poultry Science, University of
Guelph, Guelph, Ontario, N1G 2W1, Canada.

(Received April 5, 1993; Accepted June 17, 1993)

HALE B., RYAN D., ORMROD D. P. and ALLEN O. B. *An integrated statistical approach to estimating plant responses to sequential and concurrent gaseous pollutants*. BIOTRONICS 22, 35-46, 1993. Controlled environment studies of plant response to multiple environmental stresses frequently have physical (such as chamber space) and analytical (such as experimental design and data summary) limitations. This study demonstrates the use of an efficient, integrated statistical approach in evaluating rutabaga (*Brassica napus* L. ssp. *rapifera* (Metzg.) Sinsk cv. Laurentian) and cabbage (*Brassica oleracea* L. var. *capitata* cv. Market Prize) shoot growth responses to sulphur dioxide (SO₂) followed by ozone (O₃) at acute doses. The approach combines analysis of covariance, an incomplete factorial experimental design, polynomial dose response functions and a reduced-rank regression procedure to the comparison of functions. Young plants were exposed to SO₂ on one day followed by O₃ on the next day. Growth responses to sequential exposure were compared with previously reported growth responses to concurrent exposure to the same doses. Rutabaga growth was sensitive to both gases in both exposure regimes, whereas cabbage growth was sensitive to only SO₂ in the sequential exposure. Rutabaga leaf area and shoot fresh weight responses to sequential exposures followed the same pattern as concurrent responses, and their magnitudes following sequential exposures were approximately 60% of the concurrent responses. Rutabaga shoot dry weight and cabbage shoot fresh and dry weight, and leaf area responses to the sequential exposures were different in both magnitude and pattern from responses to the concurrent exposures. The importance of this work lies in the method for quantification and comparison of plant response to pollutant mixtures, in different exposure patterns. Using a suite of off-the-shelf statistical techniques, plant response to sequential versus concurrent exposure has been mathematically generalized over a broad range of pollutant mixture concentrations. Particularly for the purpose of environmental risk assessment, this integrated statistical technique has broad application to controlled environment studies of plant response to multiple stresses.

Key words: stress, models, covariate, cabbage, rutabaga, ozone, sulphur

dioxide, exposure dynamics.

INTRODUCTION

Environmental stresses in combination may have plant effects of greater magnitude than predicted from single stress studies; a good example of this phenomenon is the phytotoxicity of gaseous pollutant mixtures (12, 18, 19, 15, 6, 3). While concurrent exposure to SO₂ and O₃ is possible in the ambient environment, particularly in less well-developed countries lacking state-of-the-art control technology, it is more likely that plants would be exposed to these gases sequentially (10, 11). Plant response to the first pollutant gas may modify plant uptake or detoxification of the second pollutant gas, thus altering response to the second gas. Several mechanisms for sequential effects of pollutant gases have been proposed, whether the second exposure is to the same or to a different gas. Pre-exposure to O₃ increased the leaf diffusive resistance, thereby reducing gas flux to the leaf interior during a subsequent O₃ episode (21). Hypothetically, pre-exposure may activate detoxification or repair mechanisms (for example, elevated oxidase enzymes), which then minimize the effect of the second episode; however, evidence of this has been difficult to establish (5).

The assessment of plant response to mixtures of pollutant gases is usually carried out with a full factorial experimental design (4). Given that each treatment combination needs a separate exposure chamber, most studies are quite restricted in either the number of discrete exposure concentrations of each pollutant, or the number of replications. Either of these limitations can reduce the strength with which the plant response is estimated: the former by limiting the range of inference, and the latter by decreasing the precision of the estimate. It is possible to counter these limitations, using experimental designs and data analyses which increase the range of inference (without increasing the number of treatments) and also increase the experimental precision. Four distinct statistical methods have been separately demonstrated to be very effective at enhancing the collection, analysis or interpretation of data describing plant responses to pollutant doses:

- 1) Incomplete factorial designs are well suited for efficient determination of pollutant dose-plant response relationships: the number of discrete exposure chambers is greatly reduced, often by more than 50%, with little loss of information (17, 1);

- 2) Analysis of covariance can increase precision in randomized experiments by reducing the contribution of pre-treatment variation to experimental error. Depending on the degree of correlation between the covariate and response variable, analysis of covariance will reduce the experimental error, thus increasing precision (16);

- 3) Multiple regression can generalize the individual treatment responses into dose-response relationships. In this form, the data may be used to predict plant response to treatment combinations not included in (but within the boundaries of) the incomplete factorial design (13);

4) Reduced-rank multivariate analysis is suitable for comparing the shapes of several dose-response relationships (22).

These four procedures can be integrated into an efficient, simple statistical protocol for the determination of plant response to multiple environmental stresses, with particular application to gaseous pollutants. This protocol is well suited for assessing the responses of plants to global climate change, particularly in controlled environments: elevated carbon dioxide, ultraviolet-B radiation and air temperature are all difficult to apply on an experimental unit basis, and a study of their interaction needs an experimental approach which maximizes resource use. The objective of this study is to describe growth responses of young cabbage and rutabaga plants to sequential exposure to SO₂ and O₃, and to determine the similarity of the responses to concurrent exposure to the same gases. These determinations were made using the previously described statistical procedures, and their effectiveness at investigating plant responses to multiple stresses is demonstrated.

MATERIALS AND METHODS

Rutabaga (*Brassica napus* L. ssp. *rapifera* (Metzg.) Sinsk cv. Laurentian) and cabbage (*B. oleracea* var. *capitata* cv. Market Prize) seeds were sown in Promix BX medium (60% peatmoss, 20% perlite and 20% vermiculite, v/v) in 10 cm diameter green plastic pots which were placed in a controlled environment chamber. The photoperiod was 16 h (0400–2000h) at $325 \pm 20 \mu\text{E} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$, measured with a Li-Cor quantum sensor and meter at canopy height. Input wattage was 77% from fluorescent and 23% from incandescent lamps. The thermoperiod was 12 h (0600–1800) with a day temperature of $25 \pm 1^\circ\text{C}$ and a night temperature of $20 \pm 1^\circ\text{C}$. Relative humidity was maintained at $75 \pm 5\%$ during both day and night. The pots were irrigated with half-strength Hoagland's complete nutrient solution (9). Seven days after sowing, the seedlings were thinned to one per pot. The two species developed at similar rates; the plants had one pair of large expanding true leaves by the day of pollutant exposure (14 d after sowing).

Pollutant exposure

On day 14 after sowing, planar leaf area (PLA) was determined by placing a transparent acetate sheet with a 1 cm² grid over each plant, and recording the number of grid intersections directly over plant tissue. Planar leaf area differs from true leaf area (LA) in that it does not include leaf area hidden under upper leaves. This number was used, untransformed, as the covariate for all response variables except visible injury (PLI). Immediately following covariate measurement, three plants of each cultivar were placed in a Continuous Stirred Tank Reactor chamber (CSTR, for details see 8) for pollutant exposure. Canopy light level was $290 \pm 20 \mu\text{E} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$, supplied by one 400 W metal halide and one 400 W high pressure sodium lamp over each CSTR chamber; the lamps were angled so that the beams intersected in the area of the plant canopy. Relative humidity

was $70 \pm 5\%$, and temperature was $25 \pm 2^\circ\text{C}$. The pollutant exposure period was 6 h (1000–1600) for each gas on two consecutive days, after which the plants were returned to the growth chamber for two days. Ozone was produced by a Grace high-voltage generator (Model LG-2-LI) and monitored with a Dasibi analyzer (Model 1003 AH). Sulphur dioxide was supplied by a pressurized cylinder containing 0.5% SO_2 in nitrogen and monitored with a Thermo Electron Series 43 pulsed fluorescent SO_2 analyzer. Sulphur dioxide exposure occurred first, followed by O_3 on the second exposure day; the treatment concentrations were (O_3/SO_2) 0/0.5, 0/1.3, 0.2/0, 0.2/0.9, 0.2/1.7, 0.4/0.5, 0.4/1.3 $\mu\text{l} \cdot \text{l}^{-1}$.

Harvesting procedures

On the second day after exposure (17 d after sowing) the plants were harvested. The shoot was designated as all parts of the plant above the junction of the cotyledons with the stem, including the cotyledons. Leaf area was determined photometrically and shoot fresh and dry (after 48 h at 60°C) weights were determined gravimetrically. Percent foliar injury was determined as that proportion of total leaf area showing chlorotic or necrotic lesions, or abaxial pitting.

Experimental design and data analysis

The experimental design was an incomplete factorial design: six pairs of gas mixture treatments were chosen such that they formed a hexagon with the seventh gas mixture in the centre of the hexagon (1). Since four CSTR's constituted a block, each replicate was divided into two blocks in such a way that the fitted response surface was orthogonal to blocks (2). Each block was repeated four times, giving eight replicates of the central treatment (0.2/0.9) and four replicates of the peripheral treatments (all others). The extra replication of the central treatment allows for estimation of lack of fit. There were three subsamples (pots) per experimental unit. The data were analyzed using the General Linear Models (GLM) and Regression (REG) procedures of SAS (Statistical Analysis System, SAS Institute Inc, Cary, NC.). The data were first analyzed using proc GLM to determine treatment, block and covariate effects, as well as the partitioning of sums of squares amongst the various error terms, represented by the simplified covariance model:

$$Y_{ijk} = \mu + R_i + T_j + RT_{ij} + B(R)_{k(i)} + a(PLA) + E_{ijk}$$

where $R = \text{rep}$ (1, ..., 4), $T = \text{treatment}$ (1, ..., 7) and $B = \text{block}$ (1, 2). Covariate adjusted treatment means were calculated for the reduced data sets and inspected for trends. For each variable, the simplified covariance model was compared (using an F test) to the full polynomial model, where treatment effects were expanded and replication effects were pooled with residual error E_{ij} :

$$Y_{ij} = \beta_0 + \beta_1(\text{SO}_2) + \beta_2(\text{O}_3) + \beta_{12}(\text{SO}_2 \times \text{O}_3) + \beta_{11}(\text{SO}_2)^2 + \beta_{22}(\text{O}_3)^2 + a(\ln PLA) + E_{ij}$$

In no case was there a significant lack of fit ($P \leq 0.1$) and most P values were greater than 0.35, suggesting that the polynomial model was an adequate

estimator of the treatment effects described by the simplified covariance model.

This full polynomial model was reduced by the method of backward selection (6). Lack of significant increase in error sum of squares (ESS) in comparison to the full model ESS, indicated the simplest model of good fit to describe the response of each growth variable of each cultivar. The first reduced model to be tested was that which did not include the interaction term ($\text{SO}_2 \times \text{O}_3$). If the error associated with this model was not greater than the full model, progressively simpler models were tested until the simplest good fitting model was arrived at. The ESS were compared by:

$$F = (\text{EMS}_r - \text{EMS}_f) / \text{EMS}_f$$

with $r-f$, f degrees of freedom. The selection of an α level for the inclusion/exclusion of model term is not highly critical (6); a value less than 0.20 is usually used.

The final reduced model was then tested for lack of fit by comparison with the full polynomial model ($P \leq 0.15$). The exclusion of important terms from the model leads to a biased estimate of the response surface, as suggested by significant lack of fit, whereas inclusion of non-significant terms increases the variability of the surface. For most of the equations, there was no significant lack of fit suggesting that the final reduced equations were appropriate descriptions of plant response to SO_2 and O_3 . If significant lack of fit was detected, a higher order model which did not have significant lack of fit was used to describe these growth responses.

The partial R^2 for each dose response relationship was calculated to evaluate the contribution of the pollutant dose factors to the fit of the data to the regression relationship. The partial R^2 was defined as $\text{RSS}/(\text{RSS} + \text{ESS})$, where RSS was the additional reduction in variation from fitting the response surface after consideration of the covariates and block/replicate effects, and ESS was the error sum of squares (4). The F test for the significance of the partial R^2 was calculated as:

$$\frac{(\text{partial } R^2)(\text{error df})}{(1 - \text{partial } R^2)(\text{regression df})}$$

and was used to evaluate the predictive strength of the relationship (23).

One of the limitations of describing plant response to pollutants in terms of dose-response relationships, is that a general comparison of responses (for example, under different exposure scenarios) is often desirable. In this study, plant responses to sequential and concurrent exposures were compared in order to test whether predisposition to or protection from O_3 (or neither) was conferred by pre-exposure to SO_2 , or whether the plant responded similarly to concurrent and sequential exposures. The shapes of these functions describing plant growth response to sequential SO_2/O_3 exposure were then compared to functions describing plant growth response to the same concentrations of SO_2/O_3 except that the pollutants were administered concurrently (13). The comparison

was accomplished using reduced-rank regression (22). This procedure compares the shapes of two or more response surfaces by testing whether the coefficients of one response surface are similar in proportion to the coefficients of another. The coefficients of each response surface are tested as a group against the coefficients of another response surface, rather than individually. The surfaces must be described by the same regression terms; this condition can be easily met by fitting the full model to the data, and not removing any terms, no matter how small the coefficient is. The data must also have been collected using the same experimental design. The elevations of the surfaces (i.e., the intercepts) are not compared by this method, so the response variables need not be measured using the same units.

In mathematical terms, this reduced-rank regression procedure arranges the coefficients of the equations to be compared into a matrix of two rows and five columns and determines whether one row can be expressed as a multiple of the other, i.e.; whether the rank (p) of the matrix is equal to 2 or 1. The general form of the matrix was as follows:

$$\begin{bmatrix} C_1 & C_2 & C_{11} & C_{22} & C_{12} \\ S_1 & S_2 & S_{11} & S_{22} & S_{12} \end{bmatrix}$$

where C_1 and S_1 are the linear SO_2 coefficients for the plant growth response functions 1 (concurrent) and 2 (sequential), C_2 and S_2 are the linear O_3 coefficients, C_{11} and S_{11} are the quadratic SO_2 coefficients, C_{22} and S_{22} are the quadratic O_3 coefficients and C_{12} and S_{12} are the interaction coefficients. Whether one row could be expressed as a multiple of the other (i.e., calculating the rank of the matrix) was tested using a likelihood ratio statistic. This is equivalent to testing whether the concurrent response was a rescaling of the sequential response,

$$H_0: S_{(i,j)} = K(C_{(i,j)}), \text{ where } K = (1, \dots, k).$$

If the null hypothesis was accepted, K was calculated by forming a matrix from the equations to be compared, and determining whether the rank of that matrix was one (indicating that one response was the same as the other) or different from one, indicating that one response was a rescaling of the other (22).

RESULTS AND DISCUSSION

An examination of the treatment means for cabbage shoot size after exposure to SO_2 followed on the next day by exposure to O_3 indicated that fresh and dry weights were strongly modified by the gases, whereas leaf area was slightly modified (Table 1). Cabbage specific leaf area and specific water

Table 1. Treatment means of cabbage and rutabaga shoot size following sequential exposure to SO₂ followed by O₃

O ₃ ($\mu\text{l.l}^{-1}$)	0			.2		.4			s.e. [†] P [‡]	
SO ₂ ($\mu\text{l.l}^{-1}$)	.5	1.3	0	.9	1.7	.5	1.3			
CABBAGE										
Leaf area (cm ²)	135 [§]	112	120	112	80.1	104	85.9	.137	.131	
Dry Weight (g)	.543	.405	.410	.389	.303	.381	.315	.116	.036	
Fresh weight (g)	9.77	4.78	5.35	4.96	3.34	4.47	3.79	.240	.099	
Specific leaf area (cm ² · g ⁻¹)	248	277	293	287	265	273	273	.114	.969	
Specific water content (g · g ⁻¹)	16.9	10.8	12.0	11.7	9.98	10.7	11.0	.150	.283	
Leaf injury (%)	.017	.782	1.18	5.98	13.0	52.5	47.4	6.01	.0001	
RUTABAGA										
Leaf area (cm ²)	162	138	133	137	92.8	111	80.3	.111	.003	
Dry weight (g)	.427	.450	.414	.438	.264	.331	.290	.106	.006	
Fresh weight (g)	5.74	5.39	5.00	5.18	3.49	3.67	2.90	.116	.003	
Specific leaf area (cm ² · g ⁻¹)	379	307	320	313	359	335	277	.089	.299	
Specific water content (g · g ⁻¹)	12.4	10.9	11.0	10.8	12.2	10.1	8.97	.104	.350	
Leaf injury (%)	.068	8.57	9.54	7.47	12.6	26.5	45.0	5.87	.004	

[†] Standard error of the transformed means; applicable to mean values after conversion to arcsine (leaf injury) or log_e (all others).

[‡] P values for F=treatment MS/error MS, where n=4.

[§] Values are the means of four replicates and three subsamples per replicate, and are adjusted to mean covariate and replicate effects.

Table 2. Polynomial dose response relationships describing cabbage and rutabaga shoot size (transformed to natural log) and leaf injury (transformed to arcsine) following sequential exposure to SO₂ and O₃.

CABBAGE	R ^{2†}	P [‡]
ln Leaf area=4.87−.242 (SO ₂)	.503	**
ln Dry weight=−0.613−.212 (SO ₂) −.756 (O ₃)	.664	**
ln Fresh weight=2.17−.347 (SO ₂) −1.29 (O ₃)	.579	**
ln Specific leaf area (n. s.) [§]		
ln Specific water content (n. s.)		
arcsine Leaf injury=−2.96+6.96 (SO ₂) −.294 (O ₃)+259 (O ₃) ²	.837	**
RUTABAGA		
ln Leaf area=5.22−.241 (SO ₂) −1.15 (O ₃)	.768	**
ln Dry weight=−.713+.266 (SO ₂) −.869 (O ₃) −.269 (SO ₂) ²	.447	**
ln Fresh weight=1.92−.208 (SO ₂) −1.33 (O ₃)	.771	**
ln specific leaf area (n. s.)		
ln Specific water content (n. s.)		
arcsine Leaf injury=7.24+68.2 (O ₃)	.701	

[†] partial R²

[‡] P of F for H₀: partial R² is equal to zero

[§] n. s. indicates that no regression relationship could be fitted

content were apparently not changed by any of the treatments, whereas leaf injury was related to the two gases. The dose response equations for cabbage shoot fresh and dry weights were described by a negative linear relationship to both O_3 and SO_2 , whereas leaf area was negatively related to SO_2 only (Table 2). Leaf injury was positively related to SO_2 (linear) and related to O_3 in a mixed quadratic function. The level of O_3 at which the minimum response occurred ($.00057 \mu l \cdot l^{-1}$) was not different from zero, so this response was essentially a positive relationship to both SO_2 and O_3 (Table 2).

The treatment means for rutabaga demonstrate a similar pattern of response. Leaf area, and shoot fresh and dry weights were altered by the pollutant gases, and the appearance of visible injury was related to SO_2 and O_3 . Specific leaf area and specific water content were not influenced by either gas (Table 1). The dose response equations for rutabaga were also generally consistent with the treatment means (Table 1). Leaf area and shoot fresh weight were negatively related to SO_2 and O_3 (linear). Shoot dry weight was negatively related to O_3 and to SO_2 with a mixed quadratic relationship (Table 2). The level of SO_2 which produced the maximum shoot dry weight was $0.49 \mu l \cdot l^{-1}$, which indicated that low levels of SO_2 likely had a positive effect on shoot dry weight. As in cabbage, specific leaf area and specific water content were not related to SO_2 nor O_3 concentration, consistent with the treatment means (Table 1). Leaf injury was positively related to O_3 only (linear) (Table 2). The F test of whether the partial R^2 was equal to zero was significant for all of the equations at $P \leq 0.05$ (Table 2).

The dose response relationship coefficients for O_3 were much larger than for SO_2 : a comparison of the linear coefficients indicates that O_3 was 3–4 fold as phytotoxic as SO_2 . The variable responses were similar for both species. The shoot size of both was generally depressed by the gases, and both SO_2 and O_3 were involved in most of the responses. There was no evidence of interaction between the gases.

The shoot growth dose response functions were compared to those for concurrent exposure (Table 3, from 13). On a qualitative basis, the responses were different. Cabbage was sensitive to SO_2 in sequential exposure only, and rutabaga response to SO_2 was different for concurrent and sequential exposures. The O_3 coefficients for cabbage growth responses were very similar for concurrent and sequential exposures, indicating some consistency between responses to the two regimes. When the full models were tested using the reduced-rank regression approach, the shoot response of cabbage to sequential exposure was a different shape compared to concurrent exposure response (Table 4). Leaf area and shoot fresh and dry weight responses were different because of the inclusion of a negative linear SO_2 term in sequential responses. This difference suggests that during concurrent exposure, elastic strain in response to SO_2 (for example, changes in leaf diffusive resistance) was not integrated into plant growth strain. However, if the gases were delivered sequentially, the plant appeared to integrate stress from both gases into growth strain. Rutabaga leaf area and shoot fresh weight responses did not differ in

Table 3. Final polynomial dose response relationships describing cabbage and rutabaga shoot size (transformed to natural log) and leaf injury (transformed to arcsine) following concurrent exposure to SO₂ and O₃ (13).

CABBAGE	R ^{2†}	P [‡]
ln Leaf area = $4.46 - 1.25 (O_3)$.653	* *
ln Dry weight = $-1.20 - .816 (O_3)$.649	* *
ln Fresh weight = $1.33 - 1.18 (O_3)$.661	* *
ln Specific leaf area = $5.66 - .433 (O_3)$.473	* *
ln Specific water content = $2.45 - .393 (O_3)$.441	* *
arcsine Leaf injury = $.813 - 47.9 (O_3) + .355 (O_3)^2$.845	* *
RUTABAGA		
ln Leaf area = $5.01 - .422 (SO_2) - 1.02 (O_3) + .191 (SO_2)^2$.704	* *
ln Dry weight = $-.663 - .606 (SO_2) - .800 (O_3) + .282 (SO_2)^2$.667	* *
ln Fresh weight = $1.79 - .501 (SO_2) - 1.13 (O_3) + .227 (SO_2)^2$.702	* *
ln Specific leaf area (n. s.) [§]		
ln Specific water content = $2.45 - .366 (O_3)$.561	* *
arcsine Leaf injury = $-2.74 + 4.18 (SO_2) - 22.8 (O_3) + 262 (O_3)^2$.721	* *

† partial R²‡ P of F for H₀: R² is equal to zero

§ n. s. indicates that no regression relationship could be fitted

Table 4. Comparison of sequential and concurrent dose response relationships (Tables 3 and 4) describing cabbage and rutabaga shoot size (transformed to natural log) and leaf injury (transformed to arcsine) following sequential and concurrent exposure to SO₂ and O₃.

Shoot size variable	X ^{2†}	df [‡]	P [§]	K [¶]
CABBAGE				
ln Leaf area	5.30	1	*	—
ln Dry weight	3.56	1	*	—
ln Fresh weight	4.97	1	*	—
RUTABAGA				
ln Leaf area	5.29	2	ns	.59
ln Dry weight	8.16	2	*	—
ln Fresh weight	3.36	2	ns	.65

† Chi-squared test for H₀: Dose response A = K (Dose response B) where K is a constant to be estimated.‡ Degree of freedom for X² = m (q₂ - p + 1) where m is the hypothesized number of restrictions (in this case m = 1), q₂ is the number of pollutant terms in the model and p is the number of response surfaces.§ P value for H₀ tested by X².

¶ Estimated value for K, where P > 0.05.

shape for concurrent versus sequential exposure, so one could be expressed as a rescaling of the other; rutabaga shoot dry weight responses were different for the two types of exposure. The similarity between the leaf area responses is clear: both sequential and concurrent functions have negative linear O_3 coefficients of approximately the same magnitude. However, SO_2 is a mixed quadratic for concurrent versus negative linear for sequential; the linear SO_2 coefficients for sequential functions were approximately one half of those for the concurrent functions.

The values for K (0.59 (leaf area) and 0.65 (shoot fresh weight)) indicate that the sequential responses were scaled down versions of the equations describing the concurrent responses. These results suggest that for rutabaga, although there was no statistical interaction between SO_2 and O_3 , the additivity of the responses to each of the two gases was somewhat dependent on the exposure timetable. In contrast, in the presence of O_3 , SO_2 clearly had a stimulatory effect on leaf area and dry weight, at concentrations above $1.0 \mu l \cdot l^{-1}$; in the absence of O_3 , SO_2 had a negative effect at all concentrations. The difference between the response relationships for dry weight was clear: the response to SO_2 was opposite for the two exposures. Although there was no statistical interaction between the two gases, the additivity was very dependent on the exposure timetable. The extrema (maxima and minima) for SO_2 were $0.49 \mu l \cdot l^{-1}$ and $1.1 \mu l \cdot l^{-1}$ for sequential and concurrent, respectively.

The response of plants to pre-exposure and exposure to the same gas may be different from plant response to different gases presented sequentially. Bean, exposed to $0.02 \mu l \cdot l^{-1}$ O_3 prior to being exposed to O_3 concentrations up to $0.4 \mu l \cdot l^{-1}$ O_3 was less injured as a result of acute exposure than as a result of pre-exposure to filtered air (20, 21). The pretreated plants had a lower stomatal conductance prior to the initiation of the acute exposure and had less than one half as much leaf injury as plants which were not pretreated with a chronic level of O_3 . Since the same gas was used for both treatments, it is possible that the pre-exposure not only closed stomates, but also initiated detoxification mechanisms, which were then well in place for the subsequent exposure. It is unlikely that a similar sequence of events would occur for two different gases, for example SO_2 and O_3 , as the proposed detoxification mechanisms are different for each gas (14).

SUMMARY

These data for cabbage and rutabaga clearly indicate the potential for interaction between gases in their effects on plant growth response to gaseous pollutant mixtures, depending on the exposure regime. Although interaction was not detectable by examining for a statistically significant interaction term ($SO_2 \times O_3$), growth response to the gases depended upon whether the gases were presented sequentially or concurrently. The nature of the joint action was species dependent: cabbage seemed to be more sensitive to sequential exposure, whereas rutabaga was more sensitive to concurrent. The integration of the four

statistical techniques (analysis of covariance, incomplete factorial experimental design, polynomial dose response functions and comparison of dose response functions using reduced-rank regression) constituted an efficient, simple approach to determination of plant response to pollutant mixtures using commercial statistical software. Analysis of covariance, the incomplete factorial design and development of dose-response functions have been proven in earlier studies to increase the precision of detecting treatment effects and improve the efficiency of data collection. Polynomial dose-response functions enable estimation of plant response to treatment concentrations not utilized in the experiment, and through differentiation, reveal treatment levels at which plant response changes. Both of these goals are very difficult to achieve using other data comparison techniques, such as multiple range tests. A reduced-rank regression approach to comparison of polynomial dose-response functions is a good method for quantitative determination of similarity between two dose responses. This latter technique can be applied to comparison of exposure regimes, and would also be suitable for comparing species responses or responses of different growth parameters.

ACKNOWLEDGMENTS

The authors wish to acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC) in the form of a Strategic Grant to D.P.O. and O.B.A. as well as the Ontario Ministry of Agriculture and Food in the form of technical support (L. Pyear). An Ontario Graduate Scholarship to B.A.H. is gratefully acknowledged.

REFERENCES

1. Allen O. B., Marie (Hale) B. A. and Ormrod, D. P. 1987. Relative efficiency of factorial designs for estimating response surfaces with reference to gaseous pollutant mixtures. *J. Environ. Qual.* **16**, 316-320.
2. Box G. E. P. and Hunter, J. S. 1957. Multi-factor experimental designs for exploring response surfaces. *Ann. Math. Stat.* **28**, 195-241.
3. Carlson R. W. 1979. Reduction in the photosynthetic rate of *Acer*, *Quercus*, and *Fraxinus* species caused by sulfur dioxide and ozone. *Environ. Poll.* **18**, 159-170.
4. Deveau J. L., Ormrod, D. P., Allen O. B. and Beckerson, D. W. 1987. Growth and foliar injury responses of maize, soybean and tomato seedlings exposed to mixtures of ozone and sulphur dioxide. *Agric. Ecosys. and Environ.* **19**, 223-240.
5. Dijak M. and Ormrod, D. P. 1982. Some physiological and anatomical characteristics associated with differential ozone sensitivity among pea cultivars. *Environ. Exp. Bot.* **22**, 395-402.
6. Draper N. R. and Smith, H. 1981. *Applied Regression Analysis*. John Wiley and Sons, New York. pp 709.
7. Heagle A. S. and Johnston, J. W. 1979. Variable responses of soybean to mixtures of ozone and sulfur dioxide. *JAPCA* **29**, 729-732.
8. Heck W. W., Philbeck R. B. and Dunning, J. A. 1978. A continuous stirred tank reactor (CSTR) system for exposing plants to gaseous air pollutants. Principles, specifications, construction and operation. *ARS, USDA ARSS-181*, pp 32.

9. Hoagland D.R. and Arnon. D.I. 1951. The water culture method for growing plants without soil. *California Agricultural Experimental Station Circ.* **34**, 2-39.
10. Lefohn A. S., Hogsett W.E. and Tingey. D.T. 1987. The development of sulfur dioxide and ozone exposure profiles that mimic ambient conditions in the rural southeastern United States. *Atmos. Environ.* **21**, 659-666.
11. Lefohn A. S., Davis, C. E. Jones, C.K. Tingey D.T. and Hogsett. W.E. 1987. Co-occurrence patterns of gaseous air pollutant pairs at different minimum concentrations in the United States. *Atmos. Environ.* **21**, 2435-2444.
12. Marie (Hale) B.A. and Ormrod. D.P. 1984. Tomato plant growth with continuous exposure to sulfur dioxide and nitrogen dioxide. *Environ. Poll.* **33**, 257-265.
13. Marie (Hale) B.A. and Ormrod. D.P. 1986. Dose response relationships of the growth and injury effects of ozone and sulphur dioxide on brassicaceae seedlings. *Can. J. Plant Sci.* **66**, 659-667.
14. Olszyk D.M. and Tingey. D.T. 1985. Metabolic basis for injury to plants from combinations of O₃ and SO₂. *Plant Physiol.* **77**, 935-939.
15. Ormrod D.P. 1982. Air pollutant interactions in mixtures. In Unsworth, M.H. and D.P. Ormrod, (eds), *Gaseous Air Pollution in Agriculture and Horticulture*. Butterworths, pp 532.
16. Ormrod D. P., Tingey D. T. and Gumpertz. M. 1983. Covariate measurements for increasing the precision of plant response to O₃ and SO₂. *HortScience* **18**, 896-898.
17. Ormrod D. P., Tingey, D. T. Gumpertz M.L. and Olszyk. D.M. 1984. Utilization of a response surface technique in the study of plant responses to ozone and sulfur dioxide mixtures. *Plant Physiol.* **75**, 43-48.
18. Petite J. M. and Ormrod. D. P. 1984. Effects of sulfur dioxide and nitrogen dioxide on four *Solanum tuberosum* L. cultivars. *Am. Potato. J.* **61**, 320-329.
19. Reinert R.A. 1984. Plant responses to air pollutant mixtures. *Ann. Rev. Phytopath.* **22**, 421-442.
20. Runeckles V.C. and Rosen. P.M. 1974. Effects of pretreatment with low ozone concentrations on ozone injury to bean and mint. *Can. J. Bot.* **52**, 2607-2610.
21. Runeckles V.C. and Rosen. P.M. 1977. Effects of ambient ozone pretreatment on transpiration and susceptibility to ozone injury. *Can. J. Bot.* **55**, 193-197.
22. Ryan D. A. J., Hubert, J. J. Carter, E.M. Sprague J.B. and Parrott. J. 1992. A reduced-rank multivariate approach to aquatic joint toxicity experiments. *Biometrics* **48**, 155-162.
23. Steel R.G.D. and Torrie. J.H. 1980. *Principles and Procedures of Statistics*. McGraw-Hill, pp 633.