

NEW SYSTEM OF HYDROPONICS FOR GROWTH ANALYSIS OF SWEET POTATO TUBER

Eguchi, Toshihiko
Biotron Institute Kyushu University

Kitano, Masaharu
Biotron Institute Kyushu University

Eguchi, Hiromi
Biotron Institute Kyushu University

<http://hdl.handle.net/2324/8222>

出版情報 : BIOTRONICS. 25, pp.85-88, 1996-12. Biotron Institute, Kyushu University
バージョン : published
権利関係 :



NEW SYSTEM OF HYDROPONICS FOR GROWTH ANALYSIS OF SWEET POTATO TUBER

T. EGUCHI, M. KITANO and H. EGUCHI

Biotron Institute, Kyushu University 12, Fukuoka 812-81, Japan

(Received August 1, 1996; accepted September 10, 1996)

EGUCHI T., KITANO M. and EGUCHI H. *New system of hydroponics for growth analysis of sweet potato tuber*. BIOTRONICS 25, 85-88, 1996. hydroponic system for sweet potato plants (*Ipomoea batatas* Lam.) was newly developed for non-destructive measurement of tuber growth under the controlled environment. In the system, tuber growth was evaluated as changes in the volume in course of time, and the tuber of the plant with three leaves appeared in source-limited growth. Thus, new hydroponic system made it possible to evaluate the tuber volume in course of time for the analysis of the sink-source relationship in sweet potato plants.

Key words: *Ipomoea batatas* Lam.; hydroponics; tuber growth analysis

INTRODUCTION

In the previous paper, a solution-air culture system (3) has been applied for hydroponics of sweet potato plants (*Ipomoea batatas* Lam.), where the tuber was grown in the air space (2). Furthermore, for evaluation of tuber growth in course of time, it is necessary to develop new hydroponic system. The present paper deals with the hydroponic system for non-destructive measurement of tuber growth in sweet in potato plants.

MATERIALS AND METHODS

Plant materials

Cut-stems of sweet potato plants (cv. Koganesengan) with three, five and seven leaves were cultured for 28 days in a phytotron glass room (an air temperature of 25°C and a relative humidity of 70%). Several number of tubers were formed in a plant, and the largest tuber was kept in the plant. Fibrous roots on the basal node of the stem were kept for uptake of nutrient solution, and fibrous roots directly extended from the largest tuber were excised.

Solution-air culture system and environment control

Figure 1 shows a schematic diagram of a hydroponic system newly developed for growth analysis of sweet potato tuber. A culture box consisted of the air space and nutrient solution, where the styrofoam plate was used to




Fig. 1. Schematic diagram of newly developed hydroponic system for sweet potato plants.

sustain the tuber in the air space and to separate the air space from nutrient solution. In the system, the environments of the air space and nutrient solution were controlled respectively: air temperature and relative humidity in the air space were controlled at $24 \pm 0.5^\circ\text{C}$ and $70 \pm 5\%$ by the air conditioning unit, and temperature of nutrient solution was controlled at $24 \pm 0.5^\circ\text{C}$ by the water bath system. The hydroponic system was installed in a growth chamber, where the environment was controlled at an air temperature of $28 \pm 0.5^\circ\text{C}$, a relative humidity of $70 \pm 3\%$ and a light intensity of $300 \mu\text{mol m}^{-2} \text{s}^{-1}$ in a photoperiod of 12 h. Those environmental conditions have been found to maximize the sink strength of the tuber (1, 2). The material plants were transplanted into the hydroponic system, and grown for 20 days. The tuber volume was measured by immersing the tuber into a measuring cylinder filled with water in course of time.

RESULTS AND DISCUSSION

Figure 2 shows photographs of tubers and fibrous roots grown in the hydroponic system. Healthy growth of the tuber was found in the air space on the styrofoam (Fig. 2a), and many fibrous roots were grown in nutrient solution (Fig. 2b). Figure 3 shows time course pattern of the tuber volume in the system.




Fig. 2. Photographs of sweet potato tubers (a. black allows) in the air space, and fibrous roots (b. white allow) in nutrient solution after 20 days cultivation in the hydroponic system.




Fig. 3. Time course pattern of the tuber volume in the hydroponic system. Means of 4 plants are plotted.

The volume was increasing during 20 days, and the tuber yield of 8.7 g fresh weight was obtained. Thus, tuber growth was evaluated as changes in the volume in course of time. Figure 4 shows tuber volumes at 0 and 20 days after transplanting and tuber growth rates during 20 days in respective plants with




Fig. 4. Tuber volumes at 0 day (TV0d) and 20 days (TV20d) after transplanting and the tuber growth rate (GR) during 20 days in respective plants with three, five and seven leaves. Means of 5 plants are plotted with 95% confidence intervals.

three, five and seven leaves. Tuber growth rates of the plants with five and seven leaves were significantly higher than the rate of the plant with three leaves (5% level). Thus, the tuber of the plant with three leaves appeared in source-limited growth under the environmental condition of this experiment.

Thus, new hydroponic system made it possible to evaluate the tuber volume in course of time for the analysis of the sink-source relationship in sweet potato plants.

REFERENCES

1. Eguchi T., Kitano M. and Eguchi H. (1994) Effect of root temperature on sink strength of tuberous root in sweet potato plants (*Ipomoea batatas* Lam.). *Biotronics* **23**, 75-80.
2. Eguchi T., Kitano M. and Eguchi H. (1995) Effect of air humidity around tuberous root on sink strength in sweet potato plants grown in a solution-air culture system. *Biotronics* **24**, 45-49.
3. Sugi J. *et al.* (1975) Artificial rhizosphere. pages 281-306. in J. Sugi and Y. Tsukamoto (eds) *Development of Organisms and Environment Control*. Japan Society for the Promotion of Science, Tokyo, Japan.