

## COMPARING STUDIES OF ARTIFICIAL AND NATURAL ECOSYSTEM RESPONSES TO CO<sub>2</sub> ENRICHMENT

Jones, T. H.

Cardiff School of Biosciences Cardiff University | NERC Centre for Population Biology Imperial College at Silwood Park | NERC Centre for Population Biology Imperial College at Silwood Park

Bezemer, T. M.

N100-CTO | NERC Centre for Population Biology Imperial College at Silwood Park | NERC Centre for Population Biology Imperial College at Silwood Park

Korner, C.

Institut of Botany University of Basel | NERC Centre for Population Biology Imperial College at Silwood Park | NERC Centre for Population Biology Imperial College at Silwood Park

Lawton, J. H.

NERC Centre for Population Biology Imperial College at Silwood Park

他

<https://hdl.handle.net/2324/8259>

---

出版情報 : BIOTRONICS. 29, pp.1-7, 2000-12. Biotron Institute, Kyushu University  
バージョン :  
権利関係 :

## COMPARING STUDIES OF ARTIFICIAL AND NATURAL ECOSYSTEM RESPONSES TO CO<sub>2</sub> ENRICHMENT

T. H. JONES<sup>2</sup>, T. M. BEZEMER<sup>3</sup>, C. KÖRNER<sup>1</sup>, J. H. LAWTON and L. J. THOMPSON

*NERC Centre for Population Biology, Imperial College at Silwood Park, Ascot,  
Berkshire SL5 7PY, UK*

<sup>1</sup>*Institut of Botany, University of Basel, Schönbeinstraße 6, CH-4056 Basel,  
Switzerland Current address:*

<sup>2</sup>*Cardiff School of Biosciences, Cardiff University, PO Box 915 CF10 3TL;*

<sup>3</sup>*N100-CTO, Boterhoeksestraat 48, PO Box 40, 6666 2G Heteren, The Netherlands*

(Received February 1, 2000; accepted March 22, 2000)

JONES T.H., BEZEMER T.M., KÖRNER C., LAWTON J.H. and THOMPSON L.J. *Comparing studies of artificial and natural ecosystem responses to CO<sub>2</sub> enrichment*. BIOTRONICS 29, 1–7, 2000. Can experimental investigations of complex systems provide a clearer basis on which to make predictions about the real world? Can model ecosystems predict, at least qualitatively, natural effects? Using two superficially very different grassland ecosystems (natural alpine grassland and grassland model ecosystems) the effects of elevated atmospheric carbon dioxide (CO<sub>2</sub>) on these systems are compared. Although very different in composition, climatic conditions and successional stage, results from the two experiments show considerable similarities. Both experiments also highlight the implications for long-term feedback processes in soil ecosystems that are subject to rising global atmospheric CO<sub>2</sub> concentrations. Finally, while studies show that small-scale, short-term single species experiments can yield misleading results we advocate the use of either field manipulations of real ecosystems or, if this is not possible for reasons of cost or time, more realistic (and complex) model systems (microcosms).

**Key words:** elevated CO<sub>2</sub>; climate change; ecosystem processes; model ecosystems; controlled environment facilities

### INTRODUCTION

We know that atmospheric carbon dioxide (CO<sub>2</sub>) concentrations are increasing through human activities, and we can make relatively good predictions of the levels we can expect to see in the near future. The ecological effects of this rising CO<sub>2</sub> are not as easy to predict. Ecologists are constantly

---

Address for correspondence: Dr. T. H. Jones  
Cardiff School Biosciences, Cardiff University, PO Box 915, Cardiff CF10 3TL, U.K.  
Tel: +44-(0)29-2087-5357 Fax: +44-(0)29-2087-4305  
e-mail: jonesth@cardiff.ac.uk

being asked to make predictions at the community, ecosystem and biosphere level using information from lower levels (e.g. individual species) (22). There are numerous reasons why this is a difficult task. For example, (i) CO<sub>2</sub>-enrichment studies are frequently conducted with a free mineral nutrient supply that is not very relevant to natural vegetation. (ii) While pot experiments have contributed greatly to our understanding of how individual plants may respond to various treatments, many responses are altered when multi-species interactions come into play – biological isolation is rare in the real world. (iii) In many cases, long-term effects have not been studied; a full life-cycle (generation) represents a useful experimental duration (10). Can experimental investigations of more complex systems provide us with clearer bases on which to make predictions about the real world?

In this paper, we compare and contrast the results obtained from two recently published studies investigating the effects of elevated CO<sub>2</sub> on two superficially very different grassland ecosystems. The first is a natural alpine grassland at 2,470 m altitude in the Swiss Central Alps exposed to elevated CO<sub>2</sub> by using open top chambers (13); the second used grassland model ecosystems maintained in the Ecotron controlled environment facility at the Natural Environment Research Council (NERC) Centre for Population Biology at Silwood Park (14, 8). We chose these two “extremes” as being representative of “natural” and “artificial” ecosystem approaches to global climate change studies. We ask whether model ecosystems can predict, at least qualitatively, the consequences of enhanced atmospheric CO<sub>2</sub> in natural systems.

### DESCRIPTIONS OF SYSTEMS

Details of both ecosystems are given in the respective papers; here we summarise the salient features.

#### *Alpine grassland*

Experiments were conducted at 200–300 m above the climatic tree-line in a sedge-dominated alpine turf (*Caricetum curvulae*) near the Furk Pass in the Swiss Central Alps (47°N). Soils are deeply weathered acid (pH 4) gleyic podzo-cambisols on glacial moraines and are strongly enriched with aeolic sediments. Mean mid-summer air temperatures vary between 7 and 10°C, but canopy temperatures regularly exceed 20°C under direct solar radiation. *Carex curvula* represents about 62% of the total above-ground biomass, *Leontodon helveticus* represents 11%, other herbs (e.g. species of *Potentilla*, *Phyteuma*, *Trifolium*) account for 25%. The remaining 2% is composed of various grass species including *Poa alpina* (see 13).

Cylindrical open top chambers made of UV-transparent Plexiglass (37 cm diameter and height, enclosing roughly 500 tillers of *C. curvula*) were placed on each of 32 plots. Sixteen plots received air with 680 μmol mol<sup>-1</sup> CO<sub>2</sub>, and 16 received ambient air (ca. 355 μmol mol<sup>-1</sup>). Treatments began with snow-melt in 1992 and ended with the 1995 harvest conducted shortly after the time of the

seasonal biomass peak at 10–12 August. (For more details see 13, 19, 20).

#### *Ecotron model grassland*

The experiment used 16 model ecosystems, each 1m<sup>2</sup>, maintained in the Ecotron controlled environment facility. Temperature, humidity, light and water were computer-controlled and a 24 h diurnal pattern without seasonality was maintained. Rainfall (reverse osmosis water) was delivered from an irrigation lance situated above the soil surface. One minute long irrigations each delivering 4.2l of water occurred three times per day. Eight chambers were maintained at ambient external atmospheric CO<sub>2</sub> concentrations, which fluctuated naturally between 350 and 400 μmol mol<sup>-1</sup>, and eight were dynamically maintained at 200 μmol mol<sup>-1</sup> above ambient.

Soil was fumigated using methyl bromide and the soil profile consisted of 0.1 m<sup>3</sup> gravel topped with 0.3 m<sup>3</sup> of 40:60 sand:Surrey loam mix placed in 0.4 m<sup>3</sup> containers. Each container received 120 ml of a standard microbial inoculum prepared from a 20–25 μm pore filtrate (Whatman No. 4) of Silwood Park soil. This treatment also introduced protists and nematodes to all chambers.

The community, established in the soil, consisted of primary producers, herbivores, secondary consumers (parasitoids) and soil micro- and macro-organisms. Four species of plants were used: *Cardamine hirsuta*, *Poa annua*, *Spergula arvensis* and *Senecio vulgaris*. Over three generations, all communities became dominated by *P. annua*. An artificial 'winter' was imposed by cutting most of the above-ground vegetation at the end of each generation and replacing it immediately as litter. All chambers were initiated with the same community and several ecosystem processes were measured over three plant generations. (For more details see 3, 8, 9.)

### COMPARISON OF RESULTS

We consider six ecosystem parameters that were measured in both systems: (i) Biomass responses (plant and microbial); (ii) Gas exchange; (iii) Fate of excess carbon under elevated CO<sub>2</sub>; (iv) Soil microbial responses; (v) Plant tissue responses; and (vi) Decomposition. Rather than repeat information already presented in detail in other publications, here we only summarise the significant results (with the corresponding reference). Table 1 brings together the general pattern of response for each ecosystem property or process and, where appropriate, indicates the direction of that response, for both systems. Note that we are interested only in general response directions and not specific quantitative comparisons.

### DISCUSSION

Although very different in composition, geographical location, climatic conditions and successional stage, results from the two experiments show interesting similarities. Other than significant differences in above-ground plant

Table 1. General pattern of response for six ecosystem parameters measured in both Swiss alpine grassland and grassland model ecosystems. References refer to primary publication of research and data.

Parameter	Swiss Alpine Grassland	Ecotron Model Grassland
Biomass: above-ground	No above-ground biomass simulation. Elevated CO <sub>2</sub> had no effect on individual species biomass. (19, 20)	No marked increases in total above-ground biomass in elevated CO <sub>2</sub> . Significant species- and generation-dependent effects. (4, 8)
Biomass: root and rhizome	Total root and rhizome biomass in the uppermost 10 cm greater in plots maintained at high CO <sub>2</sub> . (13)	Significant treatment effects in root dry weights in the top layer (0–10 cm) in elevated CO <sub>2</sub> ; higher after 4.5 months, lower after 9 months. (9)
Biomass: microbial	Elevated CO <sub>2</sub> had no effects on microbial biomass. (13)	Soil microbial biomass unaffected by elevated CO <sub>2</sub> . No consistent treatment differences in bacterial taxonomic composition. Fungal taxonomic composition differed between treatments. (8, 9)
Gas exchange: CO <sub>2</sub> uptake	Net ecosystem CO <sub>2</sub> uptake was 41% higher during peak mid-season midday hours of Year 2, in Year 3 a full season carbon balance indicated a surplus of carbon of 22%. In Year 4 there was still a stimulation of CO <sub>2</sub> uptake at the beginning of the season, but the signal disappeared by mid-season. (5, 13)	Communities growing in elevated CO <sub>2</sub> fixed more carbon (10–17%) for most of the three plant generations. (8)
Fate of excess carbon in elevated CO <sub>2</sub>	Increases in soil organic matter or allocation of dissolved organic matter into deeper soil horizons likely. Changes in soil pools impossible to detect. Results suggest that carbon was transferred to the soil carbon pool (13)	Soil concentrations of dissolved organic carbon (DOC) were higher in elevated CO <sub>2</sub> . (8)
Soil microbial responses: enzymatic activity	Soil cellulase activity measured at harvest was significantly higher. Screening tests for <i>in vitro</i> substrate utilization revealed various changes in degradative abilities of the microflora, indicating altered substrate availabilities. (13, Insam <i>et al.</i> unpublished in 13)	One fungal functional group, cellulose decomposers, had higher biomass in elevated CO <sub>2</sub> . No major treatment effects recorded for urease, xylanase, trehalase and arginine deaminase. (8, 9)
Plant tissue response: above-ground	N- and C-contents were consistently increased under elevated CO <sub>2</sub> , but differences were more pronounced in forbs than in graminoids. (20)	Leaf C- and N-contents unaffected by CO <sub>2</sub> treatment. (4)
Plant tissue response: roots	Composition of below-ground tissues showed no changes under elevated CO <sub>2</sub> . (20)	No change in root C: N ratios between treatments during the course of the experiment. (9)
Decomposition	No long-term litter accumulation but reduced decomposition of litter produced under elevated CO <sub>2</sub> . (1, 6)	Increased decomposition rates of cotton strips placed in soil under elevated CO <sub>2</sub> conditions. (8) (Unpublished data show similar reduced decomposition rates in litter produced under elevated CO <sub>2</sub> .)

tissue C- and N-contents, and decomposition rates, the patterns observed are qualitatively similar from both Swiss Alps and Ecotron grassland ecosystems. For the two variables where differences are found, neither is surprising; plants are known to respond species-specifically to elevated CO<sub>2</sub> (e.g. 7), and rates of decomposition in the two studies were evaluated on two different materials (plant litter *vs.* cotton-strip).

There is a general discrepancy common to both studies between C-assimilation (whole ecosystem C-assimilation was strongly stimulated in both the Alpine and Ecotron grasslands, and canopy photosynthesis measurements showed individual species assimilating more carbon in elevated CO<sub>2</sub>) and C-accumulation in above-ground biomass (which hardly changed in either study). These results are similar to findings in other CO<sub>2</sub> enrichment projects (e.g. 2, 11, 15, 16, 17, 18, 23) where enhancement has yielded only marginally, and often not significant, effects on overall community above-ground biomass. What is also apparent from both studies is that any significant differences that do occur in plant populations in elevated CO<sub>2</sub> are species-dependent.

Both experiments also highlight, to varying degrees of detail and with different evidence, the implications for long-term feedback processes in soil ecosystems that are subject to rising global atmospheric CO<sub>2</sub> concentrations. Both ecosystems point to potential increases in root biomass in high CO<sub>2</sub> conditions (although the direction of this change varied across plant generations in the Ecotron system (8)) and significant changes in soil processes. In the Ecotron it would appear that increases in photosynthetically fixed carbon were allocated below ground, raising concentrations of dissolved organic carbon in the soil. These effects were then transmitted up the decomposer food chain. Soil microbial biomass remained unaffected, but the composition of the soil fungal species changed, with increases in rates of cellulose decomposition. In this context, we also observe the increased cellulase activity in the Swiss grassland. In the Ecotron, there were also changes in the abundance and species-composition of Collembola, fungi-feeding arthropods.

That two such different study sites yield broadly similar messages about the impacts of elevated CO<sub>2</sub> on multi-species, interacting communities implies that it may be possible to adopt a more comparative approach, using complex intact ecosystems, to make general predictions about the ecological effects of rising atmospheric CO<sub>2</sub> concentrations. Sufficient studies have now shown that highly reductionist investigations on single, or small numbers of species, for relatively short periods of time, be they on plants or insects (2, 10, 22, 4), are unlikely to provide adequate predictions of the effects of climate change on more natural complex systems. Ecosystem interactions and feed-backs (missing from simple experimental systems (see 22), involving fungi, bacteria, meso-fauna, herbivores, nutrients and water become subject to both direct and indirect effects of CO<sub>2</sub>, leading to complex spatially and temporally variable effects. We advise that further experiments designed to predict the effects of enhanced CO<sub>2</sub> on whole communities and ecosystems should abandon the use of small-scale, short-term single species experiments and instead, go either directly to field manipulation of

real ecosystems (21, 12), or if this is not possible for reasons of cost or time, to more realistic (and complex) model systems (10, 22, 14). We believe that it is only by adopting this approach and experimentally manipulating whole ecosystems from contrasting environments that we can even begin to hope to gain consensus about the potential outcome of the current 'experiment' humans are conducting with the entire biosphere.

#### ACKNOWLEDGEMENTS

Both experiments were made possible by the commitment of teams of researchers and collaborators – we gratefully acknowledge their contributions. The Ecotron climate change experiment was partially funded by the NERC TIGER the Initiative (GTS/02/646), the British Council (VIE/891/11) and the Austrian Academy of Sciences (ICBP-11/95). The Swiss project contributes to the GCTE Core research program and was funded by the Swiss National Science Foundation (31-30048.90 and 4031-0033431 (NFP 31)).

#### REFERENCES

1. Arnone J. A. III and Hirschel G. (1997) Does fertilizer application alter the effects of elevated CO<sub>2</sub> on *Carex* leaf litter quality and *in situ* decomposition in an alpine grassland? *Acta Oecol.* **18**, 201–206.
2. Bazzaz F. A. (1990) The response of natural ecosystems to the rising global CO<sub>2</sub> levels. *Ann. Rev. Ecol. Syst.* **21**, 167–196.
3. Bezemer T. M., Jones T. H. and Knight K. J. (1998) Long-term effects of elevated CO<sub>2</sub> and temperature on populations of the peach potato aphid *Myzus persicae* and its parasitoid *Aphidius matricariae*. *Oecologia* **116**, 128–135.
4. Bezemer T. M., Thompson L. J. and Jones T. H. (1998) *Poa annua* shows inter-generational differences in response to elevated CO<sub>2</sub>. *Glob. Ch. Biol.* **4**, 687–691.
5. Diemer M. W. (1994) Mid-season gas exchange of an alpine grassland under elevated CO<sub>2</sub>. *Oecologia* **98**, 429–435.
6. Hirschel G., Körner C. and Arnone J. (1997) Will rising atmospheric CO<sub>2</sub> affect leaf litter quality and *in situ* decomposition rates in native plant communities? *Oecologia* **110**, 387–392.
7. Hunt R., Hand D. W., Hannah M. A. and Neal A. M. (1993) Further response to CO<sub>2</sub>-enrichment in British herbaceous species. *Func. Ecol.* **7**, 661–668.
8. Jones T. H., Thompson L. J., Lawton J. H., Bezemer T. M., Bardgett R. D., Blackburn T. M., Bruce K. D., Cannon P. F., Hall G. S., Hartley S. E., Howson G., Jones C. G., Kampichler C., Kandeler E. and Ritchie D. A. (1998) Impacts of rising atmospheric carbon dioxide on model terrestrial ecosystems. *Science* **280**, 441–443.
9. Kampichler C., Kandeler E., Bardgett R. D., Jones T. H. and Thompson L. J. (1998) Impact of elevated atmospheric CO<sub>2</sub> concentration on soil microbial biomass and activity in a complex, weedy field model ecosystem. *Glob. Ch. Biol.* **4**, 335–346.
10. Körner C. (1995) Towards a better experimental basis for upscaling plant responses to elevated CO<sub>2</sub> and climate warming. *Pl. Cell. Env.* **18**, 1101–110.
11. Körner C. (1996) The response of complex multispecies systems to elevated CO<sub>2</sub>. Pages 20–42 in B. H. Walker and W. L. Steffen (eds) *Global Change and Terrestrial Ecosystems*. Cambridge University Press, Cambridge.
12. Körner C., Arnone J. A. III and Hilti W. (1993) The utility of enclosed artificial ecosystems in CO<sub>2</sub> research. Pages 185–198 in E. D. Schulze and H. A. Mooney (eds)

- Design and Execution of Experiments on CO<sub>2</sub> Enrichment.* Ecosystem Research Report 6, Commission of the European Communities, Brussels.
13. Körner C., Diemer M., Schächli B., Niklaus P. and Arnone J. III. (1997) The responses of alpine grassland to four seasons of CO<sub>2</sub> enrichment: a synthesis. *Acta Oecol.* **18**, 165–175.
  14. Lawton J.H. (1996) The Ecotron facility at Silwood Park: the value of “big bottle” experiments. *Ecology* **77**, 665–669.
  15. McKane R.B., Rastetter E.B., Shaver G.R., Nadelhoffer K.J., Giblin A.E., Laundre J.A. and Chapin F.S. III. (1997) Climatic effects on tundra carbon storage inferred from experimental data and a model. *Ecology* **78**, 1170–1187.
  16. Mooney H.A., Drake B.G., Luxmoore R.J., Oechel W.C. and Pitelka L. F. (1991) Predicting ecosystem responses to elevated CO<sub>2</sub> concentrations: what has been learned from laboratory experiments on plant physiology and field observations? *Bioscience* **41**, 96–104.
  17. Oechel W.C., Cowles S., Grulke N., Hastings S.J., Lawrence B., Prudomme T., Riechers G., Strain B., Tissue D. and Vourlitis G. (1994) Transient nature of CO<sub>2</sub> fertilization in arctic tundra. *Nature* **371**, 500–503.
  18. Roy J., Guillermin J.L., Navas M.L. and Dhillon S. (1996) Responses to elevated CO<sub>2</sub> in Mediterranean old-field microcosms: species, community and ecosystem components. Pages 123–138 in C. Körner and F.A. Bazzaz (eds) *Carbon Dioxide, Populations and Communities*. Physiological Ecology Series, Academic Press, San Diego.
  19. Schächli B. (1996) Growth dynamics and population developments in an alpine grassland under elevated CO<sub>2</sub>. *Oecologia* **106**, 93–99.
  20. Schächli B. and Körner C. (1997). *In situ* effects of elevated CO<sub>2</sub> on the carbon and nitrogen status of alpine plants. *Func. Ecol.* **11**, 290–299.
  21. Tissue D.T. and Oechel W.C. (1987) Physiological responses to CO<sub>2</sub> enrichment of *Eriophorum vaginatum* to elevated CO<sub>2</sub> and temperature in Alaskan tussock tundra. *Ecology* **68**, 401–410.
  22. Weiner J. (1996) Problems in predicting the ecological effects of elevated CO<sub>2</sub>. Pages 431–441 in C. Körner and F.A. Bazzaz (eds) *Carbon Dioxide, Populations and Communities*. Academic Press, London.
  23. Woodward F.I., Thompson G.B. and McKee I.F. (1991) The effects of elevated concentration of carbon dioxide on individual plants, populations, communities, and ecosystems. *Ann. Botany* **67**, 23–38.