

## EFFECT OF ADDITIONAL LOW INTENSITY LUMINESCENCE RADIATION 625nm ON PLANT GROWTH AND PHOTOSYNTHESIS

Kosobryukhov, A. A.  
Institute of Basic Biological Problems Russian Academy of Sciences

Kreslavski, V. D.  
Institute of Basic Biological Problems Russian Academy of Sciences

Khramov, R. N.  
Institute of Theoretical and Experimental Biophysics Russian Academy of Sciences

Bratkova, L. R.  
"Ranita" LTD

他

<https://hdl.handle.net/2324/8262>

---

出版情報 : BIOTRONICS. 29, pp.23-31, 2000-12. Biotron Institute, Kyushu University  
バージョン :  
権利関係 :

## EFFECT OF ADDITIONAL LOW INTENSITY LUMINESCENCE RADIATION 625 nm ON PLANT GROWTH AND PHOTOSYNTHESIS

A. A. KOSOBRYUKHOV\*, V. D. KRESLAVSKI\*, R. N. KHRAMOV\*\*,  
L. R. BRATKOVA\*\*\* and R. N. SHCHELOKOV\*\*\*\*

\**Institute of Basic Biological Problems, Russian Academy of Sciences, Pushchino, Moscow Region, 142292, Russia.*

\*\**Institute of Theoretical and Experimental Biophysics, Russian Academy of Sciences, Pushchino, Moscow Region, 142292, Russia.*

\*\*\* *“Ranita” LTD, Burdenko 11, office 3A, 119121, Moscow, Russia.*

\*\*\*\**Institute of General and Inorganic Chemistry RAS, Leninskii 31, GSP-1, 117907, Moscow, Russia.*

(Received March 12, 2000 ; accepted April 6, 2000)

KOSOBRYUKHOV, A. A., KRESLAVSKI, V. D., KHRAMOV, R. N., BRATKOVA, L. R., and SHCHELOKOV, R. N. *Effect of additional low intensity luminescence radiation 625 nm on plant growth and photosynthesis.* BIOTRONICS 29, 23–31, 2000. Tomato (*Lycopersicon esculentum* Mill.) and cabbage (*Brassica oleracea* L. var. *Cymosa*) plants were grown in a glass greenhouse under natural radiation in summer. A half of pots with plants were placed under an ordinary (control) and the other part under the “Redlight” (experimental variant) polyethylene films, both 100  $\mu$ m thick. The Redlight film had the same transmittance but transformed 3.5% of ultraviolet light falling on a plant into fluorescent radiation with a main maximum of 625 nm. Plants grown under modified solar radiation exhibited high intensity of photosynthesis at light saturation, a shift of saturation region to the higher level of radiation, as well as high efficiency of photosynthesis under low light intensity. An appreciable increase in the CO<sub>2</sub> assimilation rate and biological productivity under modified light irradiation of plants allows recommending the method of additional plant irradiation under controlled conditions. Under natural irradiation of plants this can be achieved by the use of Redlight film.

**Key words:** light-transforming film ; solar radiation ; controlled environment ; photosynthetic activity ; biological productivity

### INTRODUCTION

Light quality is an important factor in the regulation of physiological processes in plants resulting in significant changes in growth and photosynthesis (15, 3). Investigations show that even small amounts of low intensity red or far-red light can play considerable role in physiological processes. For example, an increase in the amount of far-red light at the same background irradiance of

white light caused a decrease in the leaf size and the stem lengthening of *Chenopodium album* (14). Absorption of far-red radiation by the far-red-absorbing film caused an increase in the leaf size and stem shortening of *Antirrhinums* plants (7). Low intensity red light influences the formation of photosynthetic apparatus, synthesis of chlorophyll and other physiological responses (2, 10). Increasing irradiation in the 450–800 nm range upon simultaneous lowering of ultraviolet radiation led to an increase in the leaf surface area and accumulation of dry mass by cucumber plants (6).

An increase in productivity up to 100% under additional irradiation of plants with red light with maxima at 615 nm (12) and 625 nm (1) using narrower spectral zones was shown. For this purpose, a film with a fluorescent additive that transformed part of the ultraviolet solar light absorbed by the film into fluorescent radiation in the red spectral range was used.

Under controlled light conditions (the level of PAR=115 Wm<sup>-2</sup>) the higher rate of CO<sub>2</sub> assimilation was revealed in the case of the lamps with maxima at 620 and 670 nm in red spectral range in comparison with the lamps giving maximum irradiation at 670 nm (16). These data also confirm the significance of the spectral range around 620 nm for the growth and development of plants. At the same time physiological mechanisms, ensuring positive effects of additional irradiation at 620 nm on the growth and development of plants, remain unknown.

The principal objective of this study was to understand the peculiarities of CO<sub>2</sub> exchange, efficiency of primary photosynthetic responses, pigment accumulation and growth processes in plants grown under additional luminescent red light with a maximum at 625 nm.

#### MATERIALS AND METHODS

The Broccoli cabbage (*Brassica oleracea* L. var. *Cymosa*, cv. Tonus), and tomatoes (*Lycopersicon esculentum* Mill, cv Belyi Naliv) plants were grown for 32 days in a fixed glass greenhouse at the Institute of Basic Biological Problems. Several cycles of plant cultivation were carried out under natural radiation in summer. The light intensity in the greenhouse in the middle of the day made up  $1,000 \pm 200 \mu\text{mol m}^{-2} \text{s}^{-1}$  the temperature varied within 25–35°C in the afternoon and 20–25°C at night. The plants were grown in 0.5 L pots (one plant per pot) filled with the substrate: peat/sand 1:1. Once a week, the plants were watered with the Knope nutrient solution supplemented with microelements. The pots with plants were placed under an ordinary (control) and fluorescent additive-containing (experimental variant) polyethylene films, both 100 μm thick. Red fluorescence in the experimental Redlight film was emitted by small amounts of a rare earth's compound that transformed about 3.5% of ultraviolet radiation, falling on the plants, into the fluorescent radiation with main 625 nm maximum (12). Fig. 1 shows the fluorescence spectrum of the Redlight film.

Fluorescence induction curves were recorded by one-ray method (8) in the installation consisting of monochromator ( $\pm 1$  nm), photomultiplier, storage oscilloscope (Tektronix, USA) and an Endim 322.01 M recorder (FRG). The leaf




Fig. 1. Fluorescence emission spectrum of Redlight film. The fluorescence was excited at the wavelength 340 nm.

surface was irradiated with blue excitation light ( $\lambda_m = 480 \text{ nm}$ ;  $I = 125 \mu\text{M m}^{-2} \text{ S}^{-1}$ ). The top-third part of the leaf was used. Standard nomenclature of fluorescence parameters was followed Van Kooten and Snel Jan (18). Calculated fluorescence parameters included the variable fluorescence yield ( $F_v$ ), determined as  $F_v = F_p - F_o$ , the half-rise time  $F_v/2$ , and the  $F_v/F_p$  ratio. The size of plastoquinone pool was determined by estimating the working integral which was defined as a square above the kinetic curve during its rise, normalized to the initial fluorescent level. The leaves of all plants were dark-adapted for 15 min. The light intensity was measured by an RTN 20 radiometer (Optical-Mechanical Association, t. S.-Petersburg, Russia) with a spectral sensitivity range of 180–5,500 nm.

The intensity of  $\text{CO}_2$  exchange in the leaves was measured with the help of an infrared gas analytical installation (Ciras-2) or a system with an Infralit-4 gas analyser [4]. The content of pigments was determined using a Specord M-40 spectrophotometer (Karl Zeis Jena, FRG) in 80% acetone and calculated by relevant formulas [9]. The area of leaves and the weight of the leaves and stems were determined. All experiments were done in 4–6 repeats. The average values were calculated. The differences between the variants were estimated at the level 0.95 by the student's *t*-test. The experimental results were processed using the Excel, Statmost 3<sup>2</sup>, and Photosyn Assistant programs [11].

## RESULTS

The photosynthetic capacity was monitored by measuring the chlorophyll




Fig. 2. The chlorophyll fluorescence inductions of completely developed cabbage leaves grown under control and Redlight films. A—slow part; B—fast part. The wavelength of excitation —480 nm and intensity — $125 \mu \text{mol m}^{-2} \text{s}^{-1}$ .

fluorescence induction kinetics (Fig. 2). The fluorescence induction had the typical S-like form and the  $F_v/F_p$  ratio about  $0.74 \pm 0.03$  that characterized normal occurrence of primary photosynthetic reactions in the control and test plants. The size of the plastoquinone pool was greater in the variant with the Redlight film and was equal to  $1.80 \pm 0.05$  and  $1.65 \pm 0.04$  (rel. units) in the experiment and control respectively. The half-rise time was  $1.4 \pm 0.2$  s in the control and  $1.7 \pm 0.15$  s in the experiment.

The value of the  $F_p/F_{st}$  ratio reflecting mainly the efficiency of the processes of dark carboxylation, was also slightly higher in the experiment than in the control and the difference was reliable. These processes in the plants grown under the Redlight film seem to be more effective. At the same time, the difference observed in the PS II activity in both variants, registrated as the  $F_v/F_p$  ratio, was insignificant.

The data obtained from slow fluorescence inductions are supported by the results of analysis of light and  $\text{CO}_2$  curves of gas exchange in plants. Fig. 3 and Table 1 (calculated by the Photosyn Assistant program) present the results describing the light curves of  $\text{CO}_2$  exchange in plants. The plants grown under the conditions of modified solar irradiation are characterized by the high intensity of photosynthesis at light saturation, shift of saturation area to the higher levels of irradiation, as well as high efficiency of photosynthesis in the field of low light intensities. At light saturation, photosynthesis in cabbage made up  $12.0 \pm 0.2$  and  $16.8 \pm 0.3 \mu \text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  in tomatoes  $5.84 \pm 0.1$  and  $12.9 \pm 0.3 \mu \text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  in the control and experiment respectively (under the same conditions). The activity of Rubisco, calculated from the  $\text{CO}_2$  curves (Fig. 4, Table 2), was  $86.1 \pm 2.6$  against  $72.0 \pm 2.2 \mu \text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  in the test and control variants of cabbage, respectively. In tomato plants, the difference in the activity of Rubisco for both variants was insignificant. The insignificant rise of the Rubisco activity in cabbage leaves could not explain the significant rise of  $\text{CO}_2$  assimilation by plants, grown under the light-transforming film. The effective utilization of light by plants in experiment can be explained in many




Fig. 3. Light curves of  $\text{CO}_2$  assimilation of cabbage (1, 2) and tomato (3, 4) leaves. Plants were grown under changed solar irradiation (1, 3) and under control conditions (2, 4) at intensity of light  $1,000 \pm 200 \mu\text{mol m}^{-2} \text{s}^{-1}$ .

Table 1. Light curve parameters of plant leaves grown under ordinary polyethelen film and "Redlight" film. Each value is the mean of four-five determination. Asterisks indicate 95% confidence limits.

Parameters	Cabbage		Tomato	
	Control	"Redlight"	Control	"Redlight"
Dark respiration rate, $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	$-3.08 \pm 0.21$	$-1.62 \pm 0.18^*$	$-1.09 \pm 0.15$	$-2.33 \pm 0.12^*$
Photosynthetic efficiency $\mu\text{mol (CO}_2) / \mu\text{mol (photon)}$	$0.0602 \pm 0.0022$	$0.0644 \pm 0.0012$	$0.0429 \pm 0.0011$	$0.0835 \pm 0.0014^*$
Photosynthetic capacity at light saturation $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	$12.0 \pm 0.2$	$16.8 \pm 0.3^*$	$5.84 \pm 0.1$	$12.9 \pm 0.3^*$
Light compensation point $\mu\text{mol (photon) m}^{-2} \text{ s}^{-1}$	$51.2 \pm 3.2$	$25.2 \pm 2.1^*$	$25.4 \pm 1.5$	$27.9 \pm 1.2$
Light saturation point $\mu\text{mol (photon) m}^{-2} \text{ s}^{-1}$	$250 \pm 5.4$	$285 \pm 6.2$	$162 \pm 6.6$	$183 \pm 7.2$




Fig. 4. Carbon dioxide curves of  $\text{CO}_2$  assimilation of cabbage (1, 2) and tomato (3, 4), leaves. Plants were grown under changed solar irradiation (1, 3) and under control conditions (2, 4) at intensity  $1,000 \pm 200 \mu\text{mol m}^{-2} \text{s}^{-1}$ .

respects by changes in the structural organization of photosynthetic apparatus under additional red light. The long-term plant cultivation under conditions of additional luminescent irradiation resulted in the increased content of chlorophylls ( $a+b$ ) and in the higher chlorophyll  $a/b$  ratio in the cabbage leaves. The cabbage plants accumulated  $5.05 \pm 0.12 \text{ mg dm}^{-2}$  chlorophyll ( $a+b$ ) in comparison with  $3.32 \pm 0.10 \text{ mg dm}^{-2}$  in the control. At additional radiation the chl  $a/\text{chl } b$  ratio increased from  $2.20 \pm 0.15$  to  $2.76 \pm 0.2$ .

Along with the increased content of pigments in plant leaves, a twofold rise of stomatal conductivity was observed in experimental plants in comparison to the control. Hence, the high rate of  $\text{CO}_2$  assimilation under conditions of additional irradiation of plants by light with  $\lambda_m = 625 \text{ nm}$  can be connected with a greater accessibility of carbonic acid due to the increased stomatal conductivity.

A positive effect of additional irradiation was observed in the whole plants. Thus, the area of the leaf surface in cabbage made up  $0.85 \pm 0.14$  and  $2.22 \pm 0.16 \text{ dm}^2$  and in tomatoes  $-3.12 \pm 0.14$  and  $1.87 \pm 10 \text{ dm}^2$  in experiment and control, respectively. The relative area and the specific surface density of leaves varied insignificantly in the experiment and control. Therefore, an increase in biological productivity of plants was due to the bigger fresh mass of leaves and stem in the experiment in comparison with the control.

Table 2. A/Ci curve parameters of plant leaves grown under ordinary polyethelen film and "Redlight" film. Each value is the mean of four-five determination. Asterisks indicate 95% confidence limits.

Parameters	Cabbage		Tomato	
	Control	"Redlight"	Control	"Redlight"
Dark respiration rate $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	2.42±0.21	5.04±0.31*	2.54±0.23	0.91±0.30*
Carboxylation efficiency $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	4.32±0.01	3.25±0.01	6.01±0.26	3.28±0.21*
Maximum rate of carboxylation by Rubisko $\text{mol m}^{-2} \text{ s}^{-1}$	72.0±2.2	86.1±2.6*	68.4±1.1	67.1±1.2
PAR saturated rate of electron transport $\mu\text{mol m}^{-2} \text{ s}^{-1}$	167.0±3.2	196.0±4.5*	103.2±2.4	98.5±3.2
Rate of triosphosphate utilisation $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	9.54±0.22	10.8±0.12	5.23±0.08	6.11±0.12
Photosynthetic capacity at light saturation $\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$	49.9±1.8	50.55±2.3	33.7±1.2	32.3±2.1
CO <sub>2</sub> compensation point $\mu\text{mol mol}^{-1}$	6.92±0.25	8.60±0.27	6.92±0.23	5.82±0.12

## DISCUSSION

The changes observed in the growth rates took place under conditions of high natural irradiation of plants. The insignificant lowering of ultraviolet radiation under the Redlight film (by 3.5%) could not explain the substantial accumulation of dry mass in plants. Experimental data show that UV-A that mainly passes through the greenhouse glass (1-0.5 cm) and film (1-100  $\mu\text{m}$ ) reduces insignificantly the growth of plants and sometimes stimulates it, at least at a high level of PAR (19). Hence, the stimulatory effects of the Redlight are likely to be connected with the long-term action of the light with  $\lambda_m=625 \text{ nm}$ . The regulatory effect of the red light in artificial-light culture with the resonance radiation lines of the lamps at 620 nm and 640 nm was also demonstrated in other studies [16, 20]. The red light was shown to cause changes in the plant respiration and CO<sub>2</sub> assimilation activity as well as photochemical activity of chloroplasts [16].

The regulatory effect of the red light on plants is often associated with the change in the ratio of two phytochrome forms (13). In particular, in the range of 620-640 nm, both forms of phytochrome have high absorption coefficient (20). Therefore, upon changes in activity of a number of processes induced by red

light, variations in stomatal conductivity may take place (13). The effect of monochromatic irradiation at 600 nm on the synthesis of matrix RNA of heat shock proteins was observed (7). The authors suggest the presence of a specific photoreceptor, operating only in response to red light irradiation with the wavelength about 600 nm.

The results obtained point to significant change in physiological parameters due to the stimulating effect of low intensity luminescence ( $\lambda_m=625$  nm), with a long-term action on plants. In this connection, it is possible to present a scheme of the photoregulation of metabolic processes and plant morphogenesis through the effect of light with  $\lambda_m$  625 nm on the plant genome.

An essential increase in the CO<sub>2</sub> assimilation rate and biological productivity under conditions of modified light irradiation of plants allows us to use the method of additional plant irradiation under the controlled conditions. Upon natural irradiation of plants, this can be achieved by the use of the Redlight film or the glass with a luminescent additive emitting red light. Under conditions of artificial light culture, the use of the lamps with the resonance line of plant irradiation in the range close to 625 nm may be effective.

#### REFERENCES

1. Bratkova S.A and Shchelokov R.N. (1999) Light-transforming material and composition for its obtaining. WO 99/27006.
2. Briggs W.R., Mosinger R. and Shafer E. (1988) Phytochrome regulation of greening in barley effects on chlorophyll accumulation. *Plant Physiol.* **86**, 435-440.
3. Bukhov N.G. (1993) Spectral structure of the light as a factor for changing physiological state and efficiency in plants. *Agricultural biology.* **1**, 9-18.
4. Egorov V.P., Anan'ev G.M. and Kosobryukhov A.A. (1991) Definition of a physiological state of photosynthetic systems. Methods of research. Pages 9-18 in V.I. Kefeli, V.N. Kydeyarov (eds) *Experimental ecology*, Science Moscow.
5. Habermann H.M. (1973) Evidence for two photoreactions and possible involvement of phytochrome in light-dependent stomatal opening. *Plant Physiol.* **51**, 543-548.
6. Krizek D. T., Mireski R. N. and Baily W. A. (1998) Uniformity of Photosynthetic photon flux and growth of 'Poinsett' cucumber plants under metal halide and Micro-wavepowered sulfur lamps. *Biotronics.* **27**, 81-92.
7. Kropat J. and Beck C.F. (1998) Characterization of photoreceptor and signaling pathway for light induction of the *Chlamydomonas* heat-shock gene HSP70A. *Photochem. and Photobiol.* **68**, 414-419.
8. Kuznetsov E. D., Vasilenko V. F., and Kreslavsky V. D. (1992) Stimulation effects of short-term red light and plant growth retardant on greening and formation of photosynthetic apparatus in wheat seedlings. *Plant Physiol. Biochem.* **30**, 559-564.
9. Lichtenthaler H. K. and Wellburn A. R. (1987) Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods Enzymol* **148**, 350-382.
10. Mancinelli A.L. (1994) The physiology of phytochrome action. Pages 211-269 in: Kendrick R.E., Kronenberg G.H.M., (eds) *Photomorphogenesis in plants.*, 2nd Ed. Dordrecht-Boston-London: Kluwer Academic Publishers.
11. Parson R. and Ogstone S.A. (1998) Photosyn Assistant Windows software for Analysis of photosynthesis. Dundee Scientific, Dundee, Scotland, UK, 1-54.
12. Shchelokov R.N. (1986) Polysvetan-new trend. *Bulletin of the Russian Academy of Sciences.* **10**, 50-54.
13. Smith H. (1982) Light quality, photoreception and plant strategy. *Annu. Rev. Plant*

- Physiol.* **33**, 481–518.
14. Smith H. (1986) The light environment. The perception of light quality. *In*: Kendrick R. E., Kronenberg G. H. M., (eds) *Photomorphogenesis in plants*. M. Nijhoff Publishers, Dordrecht 187–236.
  15. Tikhomirov A. A., Lisovski G. M. and Sidko F. Ya. (1991) Pages 7–30 *in Spectral light composition and plant productivity*. Science, Novosibirsk.
  16. Ushakova S. A. Tikhomirov A. A., Volkova E. K. Aleshina E. V. and Zavorueva E. N. (1997) Photosynthesis and respiration of plants, grown up on red and white light. *Russian Physiology of Plants* **44**, 367–372.
  17. Van Haeringen C. J., West J. S., Davis F. J., Gilbert A., Hadley P., Pearson S., Whelton A. E. and Henbest R. G. C. (1998) The development of solid spectral filters for the regulation of plant growth. *Photochem. Photobiol.* **67**, 407–413.
  18. Van Kooten O. and Snel Jan. F. H. (1990) The use of chlorophyll fluorescence nomenclature in plant stress physiology. *Photosynthesis Res.* **25**, 147–150.
  19. Wellmann E. (1983) UV radiation in photomorphogenesis. *in*: W. Shropshire Jr. and Mohr H (eds) *Encycl. Plant Physiol. New Series*. Berlin. Springer-Verlag. **16B**, 745–755.
  20. Zavorueva E. I. Nesterenko T. F. Volkova E. N. and Tikhomirov A. A. (1996) The photosynthetic apparatus of cucumber and pea plants, grown on red light with different spectral lineation *Russian Physiology of Plants*. **43**, 220–229.